

#12 attachment  
09/995225  
AA  
(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/62797 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/72**,  
C12N 15/63, 15/11, C07K 1/14, C12Q 1/68, G01N 33/53,  
C07K 16/28, A61K 38/17, 39/395

(21) International Application Number: PCT/US01/05676

(22) International Filing Date: 23 February 2001 (23.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/184,305	23 February 2000 (23.02.2000)	US
60/184,304	23 February 2000 (23.02.2000)	US
60/184,303	23 February 2000 (23.02.2000)	US
60/184,397	23 February 2000 (23.02.2000)	US
60/184,247	23 February 2000 (23.02.2000)	US
60/186,457	2 March 2000 (02.03.2000)	US
60/186,810	3 March 2000 (03.03.2000)	US
60/188,064	9 March 2000 (09.03.2000)	US
60/188,880	13 March 2000 (13.03.2000)	US
60/194,344	3 April 2000 (03.04.2000)	US
60/213,861	23 June 2000 (23.06.2000)	US
60/217,370	11 July 2000 (11.07.2000)	US
60/217,369	11 July 2000 (11.07.2000)	US
60/218,337	14 July 2000 (14.07.2000)	US
60/218,492	20 July 2000 (20.07.2000)	US

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VOGELI, Gabriel** [US/US]; 3005 First Avenue, Seattle, WA 98121 (US). **WOOD, Linda, S.** [US/US]; 10193 Fox Hollow, Portage, MI 49024 (US). **PARODI, Luis, A.** [SE/SE]; Grevgafan-24, S-115 43 Stockholm (SE). **LIND, Peter** [SE/SE]; Borjegatan 31 C, S-751 29 Uppsala (SE).

(74) Agents: **DELUCA, Mark et al.**; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(71) Applicant (for all designated States except US): **PHARMACIA & UPJOHN COMPANY** [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL G PROTEIN-COUPLED RECEPTORS

(57) Abstract: The present invention provides a gene encoding a G protein-coupled receptor termed nGPCR-x; constructs and recombinant host cells incorporating the genes; the nGPCR-x polypeptides encoded by the gene; antibodies to the nGPCR-x polypeptides; and methods of making and using all of the foregoing.

WO 01/62797 A2

**NOVEL G PROTEIN-COUPLED RECEPTORS****CROSS-REFERENCE TO RELATED APPLICATIONS**

The present application claims priority to Application Serial Numbers 60/184,305 filed February 23, 2000, 60/184,304 filed February 23, 2000, 60/184,303 filed February 23, 2000, 60/184,397 filed February 23, 2000, 60/184,247 filed February 23, 2000, 60/188,880 filed March 13, 2000, 60/217,369 filed July 11, 2000, 60/217,370 filed July 11, 2000, 60/219,492 filed July 20, 2000, 60/186,810 filed March 3, 2000, 60/188,064 filed March 9, 2000, 60/186,457 filed March 2, 2000, 60/213,861 filed June 23, 2000, 60/194,344 filed April 3, 2000, and 60/218,337 filed July 14, 2000, each of which is incorporated herein by reference in its entirety.

**FIELD OF THE INVENTION**

The present invention relates generally to the fields of genetics and cellular and molecular biology. More particularly, the invention relates to novel G protein coupled receptors, to polynucleotides that encode such novel receptors, to reagents such as antibodies, probes, primers and kits comprising such antibodies, probes, primers related to the same, and to methods which use the novel G protein coupled receptors, polynucleotides or reagents.

**BACKGROUND OF THE INVENTION**

The G protein-coupled receptors (GPCRs) form a vast superfamily of cell surface receptors which are characterized by an amino-terminal extracellular domain, a carboxyl-terminal intracellular domain, and a serpentine structure that passes through the cell membrane seven times. Hence, such receptors are sometimes also referred to as seven transmembrane (7TM) receptors. These seven transmembrane domains define three extracellular loops and three intracellular loops, in addition to the amino- and carboxy- terminal domains. The extracellular portions of the receptor have a role in recognizing and binding one or more extracellular binding partners (*e.g.*, ligands), whereas the intracellular portions have a role in recognizing and communicating with downstream molecules in the signal transduction cascade.

The G protein-coupled receptors bind a variety of ligands including calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and even photons, and are important in the normal (and sometimes the aberrant) function of many cell types. (See generally Strosberg, *Eur. J. Biochem.* 196:1-10 (1991) and Bohm *et al.*, *Biochem J.* 322:1-18 (1997)). When a specific ligand binds to its corresponding receptor, the ligand typically stimulates the receptor to activate a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) that is coupled to the intracellular portion of the receptor. The G protein in turn transmits a signal to an effector molecule within the cell, by either stimulating or

inhibiting the activity of that effector molecule. These effector molecules include adenylate cyclase, phospholipases and ion channels. Adenylate cyclase and phospholipases are enzymes that are involved in the production of the second messenger molecules cAMP, inositol triphosphate and diacylglycerol. It is through this sequence of events that an extracellular ligand  
5 stimuli exerts intracellular changes through a G protein-coupled receptor. Each such receptor has its own characteristic primary structure, expression pattern, ligand-binding profile, and intracellular effector system.

Because of the vital role of G protein-coupled receptors in the communication between cells and their environment, such receptors are attractive targets for therapeutic intervention, for  
10 example by activating or antagonizing such receptors. For receptors having a known ligand, the identification of agonists or antagonists may be sought specifically to enhance or inhibit the action of the ligand. Some G protein-coupled receptors have roles in disease pathogenesis (*e.g.*, certain chemokine receptors that act as HIV co-receptors may have a role in AIDS pathogenesis), and are attractive targets for therapeutic intervention even in the absence of  
15 knowledge of the natural ligand of the receptor. Other receptors are attractive targets for therapeutic intervention by virtue of their expression pattern in tissues or cell types that are themselves attractive targets for therapeutic intervention. Examples of this latter category of receptors include receptors expressed in immune cells, which can be targeted to either inhibit autoimmune responses or to enhance immune responses to fight pathogens or cancer; and  
20 receptors expressed in the brain or other neural organs and tissues, which are likely targets in the treatment of mental disorder, depression, schizophrenia, bipolar disease, or other neurological disorders. This latter category of receptor is also useful as a marker for identifying and/or purifying (*e.g.*, via fluorescence-activated cell sorting) cellular subtypes that express the receptor. Unfortunately, only a limited number of G protein receptors from the central nervous  
25 system (CNS) are known. Thus, a need exists for G protein-coupled receptors that have been identified and show promise as targets for therapeutic intervention in a variety of animals, including humans.

## SUMMARY OF THE INVENTION

The present invention relates to an isolated nucleic acid molecule that comprises a  
30 nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, or a fragment thereof. In some embodiments, the nucleic acid molecule encodes at least a portion of SEQ ID NO:111 to SEQ ID NO:120. In some embodiments, the nucleic acid molecule comprises a sequence that encodes a polypeptide comprising a sequence selected from the group

consisting of SEQ ID NO:61 to SEQ ID NO:120, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence selected from the group  
5 consisting of SEQ ID NO:1 to SEQ ID NO:60, or fragments thereof.

According to some embodiments, the present invention provides vectors which comprise the nucleic acid molecule of the invention. In some embodiments, the vector is an expression vector.

According to some embodiments, the present invention provides host cells which  
10 comprise the vectors of the invention. In some embodiments, the host cells comprise expression vectors.

The present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, said portion comprising at least 10 nucleotides.

15 The present invention provides a method of producing a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, or a homolog or fragment thereof. The method comprising the steps of introducing a recombinant expression vector that includes a nucleotide sequence that encodes the polypeptide into a compatible host cell, growing the host cell under conditions for expression of the polypeptide  
20 and recovering the polypeptide.

The present invention provides an isolated antibody which binds to an epitope on a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, or a homolog or fragment thereof.

The present invention provides an method of inducing an immune response in a mammal  
25 against a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, or a homolog or fragment thereof. The method comprises administering to a mammal an amount of the polypeptide sufficient to induce said immune response.

The present invention provides a method for identifying a compound which binds  
30 nGPCR-x. The method comprises the steps of contacting nGPCR-x with a compound and determining whether the compound binds nGPCR-x.

The present invention provides a method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x. The method comprises the steps of contacting said



nucleic acid molecule encoding nGPCR-x with a compound and determining whether said compound binds said nucleic acid molecule.

The present invention provides a method for identifying a compound which modulates the activity of nGPCR-x. The method comprises the steps of contacting nGPCR-x with a  
5 compound and determining whether nGPCR-x activity has been modulated.

The present invention provides a method of identifying an animal homolog of nGPCR-x. The method comprises the steps screening a nucleic acid database of the animal with a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or a portion thereof and determining whether a portion of said library or database is homologous to said sequence  
10 selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or portion thereof.

The present invention provides a method of identifying an animal homolog of nGPCR-x. The methods comprises the steps screening a nucleic acid library of the animal with a nucleic acid molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or a portion thereof and determining whether a portion of said library or database is  
15 homologous to said sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or a portion thereof.

Another aspect of the present invention relates to methods of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor. The methods comprise the steps of assaying nucleic acid of a human subject to determine a presence or an  
20 absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR that is expressed in the brain. The nGPCR comprise an amino acid sequence selected from the group consisting of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120, and allelic variants thereof. A diagnosis of the disorder or predisposition is made from the presence or absence of the mutation. The presence of a mutation altering the amino acid  
25 sequence, expression, or biological activity of the nGPCR in the nucleic acid correlates with an increased risk of developing the disorder.

The present invention further relates to methods of screening for a hereditary mental disorder genotype related to nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 in a human patient. The methods comprise the steps of providing a biological sample comprising nucleic acid from the  
30 patient, in which the nucleic acid includes sequences corresponding to alleles of nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70. The presence of one or more mutations in the nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 allele is detected indicative of a hereditary mental disorder genotype.

The present invention provides kits for screening a human subject to diagnose mental disorder or a genetic predisposition therefor. The kits include an oligonucleotide useful as a

probe for identifying polymorphisms in a human nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 gene. The oligonucleotide comprises 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 gene sequence or nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 coding sequence, except for one  
5 sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution. The kit also includes a media packaged with the oligonucleotide. The media contains information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

10 The present invention further relates to methods of identifying nGPCR allelic variants that correlates with mental disorders. The methods comprise the steps of providing biological samples that comprise nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny, and detecting in the nucleic acid the presence of one or more mutations in an nGPCR that is expressed in the brain. The nGPCR comprises an  
15 amino acid sequence selected from the group consisting of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120, and allelic variants thereof. The nucleic acid includes sequences corresponding to the gene or genes encoding nGPCR. The one or more mutations detected indicate an allelic variant that correlates with a mental disorder.

The present invention further relates to purified polynucleotides comprising nucleotide  
20 sequences encoding alleles of nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 from a human with mental disorder. The polynucleotide hybridizes to the complement of SEQ ID Numbers 1, 2, 8, 31, 34, 36, 37, 39, 40, and 51-60 under the following hybridization conditions: (a) hybridization for 16 hours at 42C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60C in a wash solution  
25 comprising 0.1x SSC and 1% SDS. The polynucleotide that encodes nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 amino acid sequence of the human differs from SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 by at least one residue.

The present invention also provides methods for identifying a modulator of biological activity of nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 comprising the steps of contacting a cell  
30 that expresses nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 in the presence and in the absence of a putative modulator compound and measuring nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 biological activity in the cell. The decreased or increased nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

The present invention further provides methods to identify compounds useful for the treatment of mental disorders. The methods comprise the steps of contacting a composition comprising nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 with a compound suspected of binding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70. The binding between nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 and the compound suspected of binding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 is detected. Compounds identified as binding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 are candidate compounds useful for the treatment of mental disorder. Compounds identified as binding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, 70, or other nGPCRs can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

The present invention further provides methods for identifying a compound useful as a modulator of binding between nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 and a binding partner of nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70. The methods comprise the steps of contacting the binding partner and a composition comprising nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 in the presence and in the absence of a putative modulator compound and detecting binding between the binding partner and nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70. Decreased or increased binding between the binding partner and nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a related disease or disorder. Compounds identified as modulating binding between nGPCR-42, 46, 48, 49, 51, 52, 61, 63, 70, or other nGPCRs and an nGPCR-x binding partner can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

Another aspect of the present invention relates to methods of purifying a G protein from a sample containing a G protein. The methods comprise the steps of contacting the sample with an nGPCR for a time sufficient to allow the G protein to form a complex with the nGPCR, isolating the complex from remaining components of the sample, maintaining the complex under conditions which result in dissociation of the G protein from the nGPCR, and isolating said G protein from the nGPCR.

Another aspect of the present invention relates to methods of identifying a compound that binds to or modulates nGPCR-51. The methods comprise contacting a composition comprising nGPCR-51 and Peptide A with a test compound, or a plurality of test compounds, and determining whether the test compound or compounds compete with Peptide A for binding to nGPCR-51.

## DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

### Definitions

Various definitions are made throughout this document. Most words have the meaning that would be attributed to those words by one skilled in the art. Words specifically defined  
5 either below or elsewhere in this document have the meaning provided in the context of the present invention as a whole and as are typically understood by those skilled in the art.

“Synthesized” as used herein and understood in the art, refers to polynucleotides produced by purely chemical, as opposed to enzymatic, methods. “Wholly” synthesized DNA sequences are therefore produced entirely by chemical means, and “partially” synthesized DNAs  
10 embrace those wherein only portions of the resulting DNA were produced by chemical means.

By the term “region” is meant a physically contiguous portion of the primary structure of a biomolecule. In the case of proteins, a region is defined by a contiguous portion of the amino acid sequence of that protein.

The term “domain” is herein defined as referring to a structural part of a biomolecule  
15 that contributes to a known or suspected function of the biomolecule. Domains may be co-extensive with regions or portions thereof; domains may also incorporate a portion of a biomolecule that is distinct from a particular region, in addition to all or part of that region. Examples of GPCR protein domains include, but are not limited to, the extracellular (*i.e.*, N-terminal), transmembrane and cytoplasmic (*i.e.*, C-terminal) domains, which are co-extensive  
20 with like-named regions of GPCRs; each of the seven transmembrane segments of a GPCR; and each of the loop segments (both extracellular and intracellular loops) connecting adjacent transmembrane segments.

As used herein, the term “activity” refers to a variety of measurable indicia suggesting or revealing binding, either direct or indirect; affecting a response, *i.e.* having a measurable affect  
25 in response to some exposure or stimulus, including, for example, the affinity of a compound for directly binding a polypeptide or polynucleotide of the invention, or, for example, measurement of amounts of upstream or downstream proteins or other similar functions after some stimulus or event.

Unless indicated otherwise, as used herein, the abbreviation in lower case (gpcr) refers to  
30 a gene, cDNA, RNA or nucleic acid sequence, while the upper case version (GPCR) refers to a protein, polypeptide, peptide, oligopeptide, or amino acid sequence. The term “nGPCR-x” refers to any of the nGPCRs taught herein, while specific reference to a nGPCR (for example nGPCR-63) refers only to that specific nGPCR.

As used herein, the term "antibody" is meant to refer to complete, intact antibodies, and Fab, Fab', F(ab)2, and other fragments thereof. Complete, intact antibodies include monoclonal antibodies such as murine monoclonal antibodies, chimeric antibodies and humanized antibodies.

5 As used herein, the term "binding" means the physical or chemical interaction between two proteins or compounds or associated proteins or compounds or combinations thereof. Binding includes ionic, non-ionic, Hydrogen bonds, Van der Waals, hydrophobic interactions, etc. The physical interaction, the binding, can be either direct or indirect, indirect being through or due to the effects of another protein or compound. Direct binding refers to interactions that do  
10 not take place through or due to the effect of another protein or compound but instead are without other substantial chemical intermediates. Binding may be detected in many different manners. As a non-limiting example, the physical binding interaction between a nGPCR-x of the invention and a compound can be detected using a labeled compound. Alternatively, functional evidence of binding can be detected using, for example, a cell transfected with and expressing a  
15 nGPCR-x of the invention. Binding of the transfected cell to a ligand of the nGPCR that was transfected into the cell provides functional evidence of binding. Other methods of detecting binding are well-known to those of skill in the art.

As used herein, the term "compound" means any identifiable chemical or molecule, including, but not limited to, small molecule, peptide, protein, sugar, nucleotide, or nucleic acid,  
20 and such compound can be natural or synthetic.

As used herein, the term "complementary" refers to Watson-Crick basepairing between nucleotide units of a nucleic acid molecule.

As used herein, the term "contacting" means bringing together, either directly or indirectly, a compound into physical proximity to a polypeptide or polynucleotide of the  
25 invention. The polypeptide or polynucleotide can be in any number of buffers, salts, solutions etc. Contacting includes, for example, placing the compound into a beaker, microtiter plate, cell culture flask, or a microarray, such as a gene chip, or the like, which contains the nucleic acid molecule, or polypeptide encoding the nGPCR or fragment thereof.

As used herein, the phrase "homologous nucleotide sequence," or "homologous amino  
30 acid sequence," or variations thereof, refers to sequences characterized by a homology, at the nucleotide level or amino acid level, of at least the specified percentage. Homologous nucleotide sequences include those sequences coding for isoforms of proteins. Such isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. Homologous

nucleotide sequences include nucleotide sequences encoding for a protein of a species other than humans, including, but not limited to, mammals. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding other known GPCRs. Homologous amino acid sequences include those amino acid sequences which contain conservative amino acid substitutions and which polypeptides have the same binding and/or activity. A homologous amino acid sequence does not, however, include the amino acid sequence encoding other known GPCRs. Percent homology can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison WI), using the default settings, which uses the algorithm of Smith and Waterman (*Adv. Appl. Math.*, 1981, 2, 482-489, which is incorporated herein by reference in its entirety).

As used herein, the term "isolated" nucleic acid molecule refers to a nucleic acid molecule (DNA or RNA) that has been removed from its native environment. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules.

As used herein, the terms "modulates" or "modifies" means an increase or decrease in the amount, quality, or effect of a particular activity or protein.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues which has a sufficient number of bases to be used in a polymerase chain reaction (PCR). This short sequence is based on (or designed from) a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 50 nucleotides, preferably about 15 to 30 nucleotides. They are chemically synthesized and may be used as probes.

As used herein, the term "probe" refers to nucleic acid sequences of variable length, preferably between at least about 10 and as many as about 6,000 nucleotides, depending on use. They are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. They may be single- or double-stranded and carefully designed to have specificity in PCR, hybridization membrane-based, or ELISA-like technologies.

The term “preventing” refers to decreasing the probability that an organism contracts or develops an abnormal condition.

The term “treating” refers to having a therapeutic effect and at least partially alleviating or abrogating an abnormal condition in the organism.

5       The term “therapeutic effect” refers to the inhibition or activation factors causing or contributing to the abnormal condition. A therapeutic effect relieves to some extent one or more of the symptoms of the abnormal condition. In reference to the treatment of abnormal conditions, a therapeutic effect can refer to one or more of the following: (a) an increase in the proliferation, growth, and/or differentiation of cells; (b) inhibition (*i.e.*, slowing or stopping) of  
10 cell death; (c) inhibition of degeneration; (d) relieving to some extent one or more of the symptoms associated with the abnormal condition; and (e) enhancing the function of the affected population of cells. Compounds demonstrating efficacy against abnormal conditions can be identified as described herein.

15       The term “abnormal condition” refers to a function in the cells or tissues of an organism that deviates from their normal functions in that organism. An abnormal condition can relate to cell proliferation, cell differentiation, cell signaling, or cell survival. An abnormal condition may also include obesity, diabetic complications such as retinal degeneration, and irregularities in glucose uptake and metabolism, and fatty acid uptake and metabolism.

20       Abnormal cell proliferative conditions include cancers such as fibrotic and mesangial disorders, abnormal angiogenesis and vasculogenesis, wound healing, psoriasis, diabetes mellitus, and inflammation.

25       Abnormal differentiation conditions include, but are not limited to, neurodegenerative disorders, slow wound healing rates, and slow tissue grafting healing rates. Abnormal cell signaling conditions include, but are not limited to, psychiatric disorders involving excess neurotransmitter activity.

Abnormal cell survival conditions may also relate to conditions in which programmed cell death (apoptosis) pathways are activated or abrogated. A number of protein kinases are associated with the apoptosis pathways. Aberrations in the function of any one of the protein kinases could lead to cell immortality or premature cell death.

30       The term “administering” relates to a method of incorporating a compound into cells or tissues of an organism. The abnormal condition can be prevented or treated when the cells or tissues of the organism exist within the organism or outside of the organism. Cells existing outside the organism can be maintained or grown in cell culture dishes. For cells harbored within the organism, many techniques exist in the art to administer compounds, including (but

not limited to) oral, parenteral, dermal, injection, and aerosol applications. For cells outside of the organism, multiple techniques exist in the art to administer the compounds, including (but not limited to) cell microinjection techniques, transformation techniques and carrier techniques.

5 The abnormal condition can also be prevented or treated by administering a compound to a group of cells having an aberration in a signal transduction pathway to an organism. The effect of administering a compound on organism function can then be monitored. The organism is preferably a mouse, rat, rabbit, guinea pig or goat, more preferably a monkey or ape, and most preferably a human.

10 By "amplification" it is meant increased numbers of DNA or RNA in a cell compared with normal cells. "Amplification" as it refers to RNA can be the detectable presence of RNA in cells, since in some normal cells there is no basal expression of RNA. In other normal cells, a basal level of expression exists, therefore in these cases amplification is the detection of at least 1 to 2-fold, and preferably more, compared to the basal level.

15 As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which a probe, primer, or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, stringent conditions are selected to be about 5C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the  
20 temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH  
25 7.0 to 8.3 and the temperature is at least about 30C for short probes, primers or oligonucleotides (e.g. 10 to 50 nucleotides) and at least about 60C for longer probes, primers or oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

30 The amino acid sequences are presented in the amino to carboxy direction, from left to right. The amino and carboxy groups are not presented in the sequence. The nucleotide sequences are presented by single strand only, in the 5' to 3' direction, from left to right. Nucleotides and amino acids are represented in the manner recommended by the IUPAC-IUB Biochemical Nomenclature Commission or (for amino acids) by three letters code.

### **Polynucleotides**



The present invention provides purified and isolated polynucleotides (*e.g.*, DNA sequences and RNA transcripts, both sense and complementary antisense strands, both single- and double-stranded, including splice variants thereof) that encode unknown G protein-coupled receptors heretofore termed novel GPCRs, or nGPCRs. These genes are described herein and designated herein collectively as nGPCR-x (where x is 42, 44, 45, 46, 47, 48, 49, 50, 51, 52, 61, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2024, 2025, 2026, 2027, 2028, 2029, and 2030). That is, these genes are described herein and designated herein as nGPCR-42, nGPCR-44, etc. Table 1 below identifies the novel gene sequence nGPCR-x designation, the SEQ ID NO: of the gene sequence, the SEQ ID NO: of the polypeptide encoded thereby, and the U.S. Provisional Application in which the gene sequence has been disclosed.

Table 1

nGPCR	Nucleotide Sequence (SEQ ID NO:)	Amino acid Sequence (SEQ ID NO:)	Originally filed in:
0042	31	91	D
0042	54	114	I
0044	32	92	D
0045	33	93	D
0046	34	94	D
0046	55	115	J
0047	35	95	D
0048	36	96	D
0048	56	116	K
0049	37	97	D
0049	59	119	N
0050	38	98	D
0051	39	99	D
0051	57	117	L
0052	40	100	D
0052	58	118	M
0061	1	61	A
0061	60	120	O
0063	2	62	A
0063	53	113	H
0063	51	111	F
0064	3	63	A
0065	4	64	A
0066	5	65	A
0067	6	66	A
0068	7	67	A
0069	10	70	A
0070	8	68	A
0070	52	112	G
0071	9	69	A
0072	43	103	E
2001	21	81	C
2002	22	82	C
2003	23	83	C
2004	24	84	C
2005	25	85	C
2006	26	86	C
2007	27	87	C
2008	28	88	C

2009	29	89	C
2010	30	90	C
2011	11	71	B
2012	12	72	B
2013	13	73	B
2014	14	74	B
2015	15	75	B
2016	16	76	B
2017	17	77	B
2018	18	78	B
2019	19	79	B
2020	20	80	B
2021	41	101	E
2022	42	102	E
2024	44	104	E
2025	45	105	E
2026	46	106	E
2027	47	107	E
2028	48	108	E
2029	49	109	E
2030	50	110	E

## Legend

A= Ser. No. 60/184,305

C= Ser. No. 60/184,303

E= Ser. No. 60/184,247

G= Ser. No. 60/217,369

I= Ser. No. 60/186,810

K= Ser. No. 60/186,457

M= Ser. No. 60/194,344

O= Ser. No. 60/217,370

B= Ser. No. 60/184,304

D= Ser. No. 60/184,397

F= Ser. No. 60/188,880

H= Ser. No. 60/219,492

J= Ser. No. 60/188,064

L= Ser. No. 60/213,861

N= Ser. No. 60/218,337

When a specific nGPCR is identified (for example nGPCR-63), it is understood that only that specific nGPCR is being referred to.

As described in Example 5 below, the genes encoding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 have been detected in brain tissue indicating that these nGPCR proteins are neuroreceptors.

The invention provides purified and isolated polynucleotides (*e.g.*, cDNA, genomic DNA, synthetic DNA, RNA, or combinations thereof, whether single- or double-stranded) that comprise a nucleotide sequence encoding the amino acid sequence of the polypeptides of the invention. Such polynucleotides are useful for recombinantly expressing the receptor and also for detecting expression of the receptor in cells (*e.g.*, using Northern hybridization and *in situ* hybridization assays). Such polynucleotides also are useful in the design of antisense and other molecules for the suppression of the expression of nGPCR-x in a cultured cell, a tissue, or an animal; for therapeutic purposes; or to provide a model for diseases or conditions characterized by aberrant nGPCR-x expression. Specifically excluded from the definition of polynucleotides of the invention are entire isolated, non-recombinant native chromosomes of host cells. A preferred polynucleotide has a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, which correspond to naturally occurring nGPCR-x sequences. It will be appreciated that numerous other polynucleotide sequences exist that also encode nGPCR-x

having the sequence selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, due to the well-known degeneracy of the universal genetic code.

The invention also provides a purified and isolated polynucleotide comprising a nucleotide sequence that encodes a mammalian polypeptide, wherein the polynucleotide  
5 hybridizes to a polynucleotide having the sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or the non-coding strand complementary thereto, under the following hybridization conditions: (a) hybridization for 16 hours at 42C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate; and (b) washing 2 times for 30 minutes each at 60C in a wash solution comprising 0.1% SSC,  
10 1% SDS. Polynucleotides that encode a human allelic variant are highly preferred.

The present invention relates to molecules which comprise the gene sequences that encode the nGPCRs; constructs and recombinant host cells incorporating the gene sequences; the novel GPCR polypeptides encoded by the gene sequences; antibodies to the polypeptides and homologs; kits employing the polynucleotides and polypeptides, and methods of making  
15 and using all of the foregoing. In addition, the present invention relates to homologs of the gene sequences and of the polypeptides and methods of making and using the same.

Genomic DNA of the invention comprises the protein-coding region for a polypeptide of the invention and is also intended to include allelic variants thereof. It is widely understood that, for many genes, genomic DNA is transcribed into RNA transcripts that undergo one or more  
20 splicing events wherein intron (*i.e.*, non-coding regions) of the transcripts are removed, or "spliced out." RNA transcripts that can be spliced by alternative mechanisms, and therefore be subject to removal of different RNA sequences but still encode a nGPCR-x polypeptide, are referred to in the art as splice variants which are embraced by the invention. Splice variants comprehended by the invention therefore are encoded by the same original genomic DNA  
25 sequences but arise from distinct mRNA transcripts. Allelic variants are modified forms of a wild-type gene sequence, the modification resulting from recombination during chromosomal segregation or exposure to conditions which give rise to genetic mutation. Allelic variants, like wild type genes, are naturally occurring sequences (as opposed to non-naturally occurring variants that arise from *in vitro* manipulation).

30 The invention also comprehends cDNA that is obtained through reverse transcription of an RNA polynucleotide encoding nGPCR-x (conventionally followed by second strand synthesis of a complementary strand to provide a double-stranded DNA).

Preferred DNA sequences encoding human nGPCR-x polypeptides are selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60. A preferred DNA of the invention

comprises a double stranded molecule along with the complementary molecule (the “non-coding strand” or “complement”) having a sequence unambiguously deducible from the coding strand according to Watson-Crick base-pairing rules for DNA. Also preferred are other polynucleotides encoding the nGPCR-x polypeptide selected from the group consisting of SEQ ID NO:61 to  
5 SEQ ID NO:120, which differ in sequence from the polynucleotides selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, by virtue of the well-known degeneracy of the universal nuclear genetic code.

The invention further embraces other species, preferably mammalian, homologs of the human nGPCR-x DNA. Species homologs, sometimes referred to as “orthologs,” in general,  
10 share at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% homology with human DNA of the invention. Generally, percent sequence “homology” with respect to polynucleotides of the invention may be calculated as the percentage of nucleotide bases in the candidate sequence that are identical to nucleotides in the nGPCR-x  
15 sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity.

Polynucleotides of the invention permit identification and isolation of polynucleotides encoding related nGPCR-x polypeptides, such as human allelic variants and species homologs,  
20 by well-known techniques including Southern and/or Northern hybridization, and polymerase chain reaction (PCR). Examples of related polynucleotides include human and non-human genomic sequences, including allelic variants, as well as polynucleotides encoding polypeptides homologous to nGPCR-x and structurally related polypeptides sharing one or more biological, immunological, and/or physical properties of nGPCR-x. Non-human species genes encoding  
25 proteins homologous to nGPCR-x can also be identified by Southern and/or PCR analysis and are useful in animal models for nGPCR-x disorders. Knowledge of the sequence of a human nGPCR-x DNA also makes possible through use of Southern hybridization or polymerase chain reaction (PCR) the identification of genomic DNA sequences encoding nGPCR-x expression control regulatory sequences such as promoters, operators, enhancers, repressors, and the like.  
30 Polynucleotides of the invention are also useful in hybridization assays to detect the capacity of cells to express nGPCR-x. Polynucleotides of the invention may also provide a basis for diagnostic methods useful for identifying a genetic alteration(s) in a nGPCR-x locus that underlies a disease state or states, which information is useful both for diagnosis and for selection of therapeutic strategies.

According to the present invention, the nGPCR-x nucleotide sequences disclosed herein may be used to identify homologs of the nGPCR-x, in other animals, including but not limited to humans and other mammals, and invertebrates. Any of the nucleotide sequences disclosed herein, or any portion thereof, can be used, for example, as probes to screen databases or nucleic acid libraries, such as, for example, genomic or cDNA libraries, to identify homologs, using screening procedures well known to those skilled in the art. Accordingly, homologs having at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, and most preferably at least 100% homology with nGPCR-x sequences can be identified.

The disclosure herein of full-length polynucleotides encoding nGPCR-x polypeptides makes readily available to the worker of ordinary skill in the art every possible fragment of the full-length polynucleotide.

One preferred embodiment of the present invention provides an isolated nucleic acid molecule comprising a sequence homologous sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, and fragments thereof. Another preferred embodiment provides an isolated nucleic acid molecule comprising a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, and fragments thereof.

As used in the present invention, fragments of nGPCR-x-encoding polynucleotides comprise at least 10, and preferably at least 12, 14, 16, 18, 20, 25, 50, or 75 consecutive nucleotides of a polynucleotide encoding nGPCR-x. Preferably, fragment polynucleotides of the invention comprise sequences unique to the nGPCR-x-encoding polynucleotide sequence, and therefore hybridize under highly stringent or moderately stringent conditions only (*i.e.*, “specifically”) to polynucleotides encoding nGPCR-x (or fragments thereof). Polynucleotide fragments of genomic sequences of the invention comprise not only sequences unique to the coding region, but also include fragments of the full-length sequence derived from introns, regulatory regions, and/or other non-translated sequences. Sequences unique to polynucleotides of the invention are recognizable through sequence comparison to other known polynucleotides, and can be identified through use of alignment programs routinely utilized in the art, *e.g.*, those made available in public sequence databases. Such sequences also are recognizable from Southern hybridization analyses to determine the number of fragments of genomic DNA to which a polynucleotide will hybridize. Polynucleotides of the invention can be labeled in a manner that permits their detection, including radioactive, fluorescent, and enzymatic labeling.

Fragment polynucleotides are particularly useful as probes for detection of full-length or fragments of nGPCR-x polynucleotides. One or more polynucleotides can be included in kits

that are used to detect the presence of a polynucleotide encoding nGPCR-x, or used to detect variations in a polynucleotide sequence encoding nGPCR-x.

The invention also embraces DNAs encoding nGPCR-x polypeptides that hybridize under moderately stringent or high stringency conditions to the non-coding strand, or complement, of the polynucleotides set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60.

Exemplary highly stringent hybridization conditions are as follows: hybridization at 42C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulfate, and washing twice for 30 minutes at 60C in a wash solution comprising 0.1 X SSC and 1% SDS. It is understood in the art that conditions of equivalent stringency can be achieved through variation of temperature and buffer, or salt concentration as described Ausubel *et al.* (Eds.), *Protocols in Molecular Biology*, John Wiley & Sons (1994), pp. 6.0.3 to 6.4.10.

Modifications in hybridization conditions can be empirically determined or precisely calculated based on the length and the percentage of guanosine/cytosine (GC) base pairing of the probe.

The hybridization conditions can be calculated as described in Sambrook, *et al.*, (Eds.), *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York (1989), pp. 9.47 to 9.51.

With the knowledge of the nucleotide sequence information disclosed in the present invention, one skilled in the art can identify and obtain nucleotide sequences which encode nGPCR-x from different sources (*i.e.*, different tissues or different organisms) through a variety of means well known to the skilled artisan and as disclosed by, for example, Sambrook *et al.*, "Molecular cloning: a laboratory manual", Second Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), which is incorporated herein by reference in its entirety.

For example, DNA that encodes nGPCR-x may be obtained by screening of mRNA, cDNA, or genomic DNA with oligonucleotide probes generated from the nGPCR-x gene sequence information provided herein. Probes may be labeled with a detectable group, such as a fluorescent group, a radioactive atom or a chemiluminescent group in accordance with procedures known to the skilled artisan and used in conventional hybridization assays, as described by, for example, Sambrook *et al.*

A nucleic acid molecule comprising any of the nGPCR-x nucleotide sequences described above can alternatively be synthesized by use of the polymerase chain reaction (PCR) procedure, with the PCR oligonucleotide primers produced from the nucleotide sequences provided herein. See U.S. Patent Numbers 4,683,195 to Mullis *et al.* and 4,683,202 to Mullis. The PCR reaction provides a method for selectively increasing the concentration of a particular

nucleic acid sequence even when that sequence has not been previously purified and is present only in a single copy in a particular sample. The method can be used to amplify either single- or double-stranded DNA. The essence of the method involves the use of two oligonucleotide probes to serve as primers for the template-dependent, polymerase mediated replication of a  
5 desired nucleic acid molecule.

A wide variety of alternative cloning and *in vitro* amplification methodologies are well known to those skilled in the art. Examples of these techniques are found in, for example, Berger *et al.*, *Guide to Molecular Cloning Techniques*, Methods in Enzymology 152, Academic Press, Inc., San Diego, CA (Berger), which is incorporated herein by reference in its entirety.

10 Automated sequencing methods can be used to obtain or verify the nucleotide sequence of nGPCR-x. The nGPCR-x nucleotide sequences of the present invention are believed to be 100% accurate. However, as is known in the art, nucleotide sequence obtained by automated methods may contain some errors. Nucleotide sequences determined by automation are typically at least about 90%, more typically at least about 95% to at least about 99.9% identical to the  
15 actual nucleotide sequence of a given nucleic acid molecule. The actual sequence may be more precisely determined using manual sequencing methods, which are well known in the art. An error in a sequence which results in an insertion or deletion of one or more nucleotides may result in a frame shift in translation such that the predicted amino acid sequence will differ from that which would be predicted from the actual nucleotide sequence of the nucleic acid molecule,  
20 starting at the point of the mutation.

The nucleic acid molecules of the present invention, and fragments derived therefrom, are useful for screening for restriction fragment length polymorphism (RFLP) associated with certain disorders, as well as for genetic mapping.

The polynucleotide sequence information provided by the invention makes possible  
25 large-scale expression of the encoded polypeptide by techniques well known and routinely practiced in the art.

### Vectors

Another aspect of the present invention is directed to vectors, or recombinant expression vectors, comprising any of the nucleic acid molecules described above. Vectors are used herein  
30 either to amplify DNA or RNA encoding nGPCR-x and/or to express DNA which encodes nGPCR-x. Preferred vectors include, but are not limited to, plasmids, phages, cosmids, episomes, viral particles or viruses, and integratable DNA fragments (*i.e.*, fragments integratable into the host genome by homologous recombination). Preferred viral particles include, but are not limited to, adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-

associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses. Preferred expression vectors include, but are not limited to, pcDNA3 (Invitrogen) and pSVL (Pharmacia Biotech). Other expression vectors include, but are not limited to, pSPORT™ vectors, pGEM™ vectors (Promega), pPROEXvectors™ (LTI, Bethesda, MD), Bluescript™ vectors (Stratagene),  
5 pQE™ vectors (Qiagen), pSE420™ (Invitrogen), and pYES2™(Invitrogen).

Expression constructs preferably comprise GPCR-x-encoding polynucleotides operatively linked to an endogenous or exogenous expression control DNA sequence and a transcription terminator. Expression control DNA sequences include promoters, enhancers, operators, and regulatory element binding sites generally, and are typically selected based on the  
10 expression systems in which the expression construct is to be utilized. Preferred promoter and enhancer sequences are generally selected for the ability to increase gene expression, while operator sequences are generally selected for the ability to regulate gene expression. Expression constructs of the invention may also include sequences encoding one or more selectable markers that permit identification of host cells bearing the construct. Expression constructs may also  
15 include sequences that facilitate, and preferably promote, homologous recombination in a host cell. Preferred constructs of the invention also include sequences necessary for replication in a host cell.

Expression constructs are preferably utilized for production of an encoded protein, but may also be utilized simply to amplify a nGPCR-x-encoding polynucleotide sequence. In  
20 preferred embodiments, the vector is an expression vector wherein the polynucleotide of the invention is operatively linked to a polynucleotide comprising an expression control sequence. Autonomously replicating recombinant expression constructs such as plasmid and viral DNA vectors incorporating polynucleotides of the invention are also provided. Preferred expression vectors are replicable DNA constructs in which a DNA sequence encoding nGPCR-x is  
25 operably linked or connected to suitable control sequences capable of effecting the expression of the nGPCR-x in a suitable host. DNA regions are operably linked or connected when they are functionally related to each other. For example, a promoter is operably linked or connected to a coding sequence if it controls the transcription of the sequence. Amplification vectors do not require expression control domains, but rather need only the ability to replicate in a host, usually  
30 conferred by an origin of replication, and a selection gene to facilitate recognition of transformants. The need for control sequences in the expression vector will vary depending upon the host selected and the transformation method chosen. Generally, control sequences include a transcriptional promoter, an optional operator sequence to control transcription, a



sequence encoding suitable mRNA ribosomal binding and sequences which control the termination of transcription and translation.

Preferred vectors preferably contain a promoter that is recognized by the host organism.

The promoter sequences of the present invention may be prokaryotic, eukaryotic or viral.

- 5 Examples of suitable prokaryotic sequences include the  $P_R$  and  $P_L$  promoters of bacteriophage lambda (The bacteriophage Lambda, Hershey, A. D., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1973), which is incorporated herein by reference in its entirety; Lambda II, Hendrix, R. W., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1980), which is incorporated herein by reference in its entirety); the trp, recA, heat shock, and lacZ promoters of
- 10 *E. coli* and the SV40 early promoter (Benoist *et al.* *Nature*, 1981, 290, 304-310, which is incorporated herein by reference in its entirety). Additional promoters include, but are not limited to, mouse mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney virus, cytomegalovirus immediate early promoter, Epstein Barr virus, Rous sarcoma virus, human actin, human myosin, human hemoglobin, human muscle creatine, and
- 15 human metallothionein.

Additional regulatory sequences can also be included in preferred vectors. Preferred examples of suitable regulatory sequences are represented by the Shine-Dalgarno of the replicase gene of the phage MS-2 and of the gene cII of bacteriophage lambda. The Shine-Dalgarno sequence may be directly followed by DNA encoding nGPCR-x and result in the

20 expression of the mature nGPCR-x protein.

Moreover, suitable expression vectors can include an appropriate marker that allows the screening of the transformed host cells. The transformation of the selected host is carried out using any one of the various techniques well known to the expert in the art and described in Sambrook *et al.*, *supra*.

- 25 An origin of replication can also be provided either by construction of the vector to include an exogenous origin or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter may be sufficient. Alternatively, rather than using vectors which contain viral origins of replication, one skilled in the art can transform mammalian cells by the method of co-transformation with a
- 30 selectable marker and nGPCR-x DNA. An example of a suitable marker is dihydrofolate reductase (DHFR) or thymidine kinase (*see*, U.S. Patent No. 4,399,216).

Nucleotide sequences encoding GPCR-x may be recombined with vector DNA in accordance with conventional techniques, including blunt-ended or staggered-ended termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends

as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulation are disclosed by Sambrook et al., *supra* and are well known in the art. Methods for construction of mammalian expression vectors are disclosed in, for example, Okayama et al., *Mol. Cell. Biol.*, **1983**, *3*, 280, Cosman et al., *Mol. Immunol.*, **1986**, *23*, 935, Cosman et al., *Nature*, **1984**, *312*, 768, EP-A-0367566, and WO 91/18982, each of which is incorporated herein by reference in its entirety.

### Host cells

According to another aspect of the invention, host cells are provided, including prokaryotic and eukaryotic cells, comprising a polynucleotide of the invention (or vector of the invention) in a manner that permits expression of the encoded nGPCR-x polypeptide.

Polynucleotides of the invention may be introduced into the host cell as part of a circular plasmid, or as linear DNA comprising an isolated protein coding region or a viral vector.

Methods for introducing DNA into the host cell that are well known and routinely practiced in the art include transformation, transfection, electroporation, nuclear injection, or fusion with carriers such as liposomes, micelles, ghost cells, and protoplasts. Expression systems of the invention include bacterial, yeast, fungal, plant, insect, invertebrate, vertebrate, and mammalian cells systems.

The invention provides host cells that are transformed or transfected (stably or transiently) with polynucleotides of the invention or vectors of the invention. As stated above, such host cells are useful for amplifying the polynucleotides and also for expressing the nGPCR-x polypeptide or fragment thereof encoded by the polynucleotide.

In still another related embodiment, the invention provides a method for producing a nGPCR-x polypeptide (or fragment thereof) comprising the steps of growing a host cell of the invention in a nutrient medium and isolating the polypeptide or variant thereof from the cell or the medium. Because nGPCR-x is a seven transmembrane receptor, it will be appreciated that, for some applications, such as certain activity assays, the preferable isolation may involve isolation of cell membranes containing the polypeptide embedded therein, whereas for other applications a more complete isolation may be preferable.

According to some aspects of the present invention, transformed host cells having an expression vector comprising any of the nucleic acid molecules described above are provided. Expression of the nucleotide sequence occurs when the expression vector is introduced into an appropriate host cell. Suitable host cells for expression of the polypeptides of the invention include, but are not limited to, prokaryotes, yeast, and eukaryotes. If a prokaryotic expression vector is employed, then the appropriate host cell would be any prokaryotic cell capable of

expressing the cloned sequences. Suitable prokaryotic cells include, but are not limited to, bacteria of the genera *Escherichia*, *Bacillus*, *Salmonella*, *Pseudomonas*, *Streptomyces*, and *Staphylococcus*.

If an eukaryotic expression vector is employed, then the appropriate host cell would be any eukaryotic cell capable of expressing the cloned sequence. Preferably, eukaryotic cells are cells of higher eukaryotes. Suitable eukaryotic cells include, but are not limited to, non-human mammalian tissue culture cells and human tissue culture cells. Preferred host cells include, but are not limited to, insect cells, HeLa cells, Chinese hamster ovary cells (CHO cells), African green monkey kidney cells (COS cells), human 293 cells, and murine 3T3 fibroblasts.

Propagation of such cells in cell culture has become a routine procedure (*see*, Tissue Culture, Academic Press, Kruse and Patterson, eds. (1973), which is incorporated herein by reference in its entirety).

In addition, a yeast host may be employed as a host cell. Preferred yeast cells include, but are not limited to, the genera *Saccharomyces*, *Pichia*, and *Kluveromyces*. Preferred yeast hosts are *S. cerevisiae* and *P. pastoris*. Preferred yeast vectors can contain an origin of replication sequence from a 2T yeast plasmid, an autonomously replication sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Shuttle vectors for replication in both yeast and *E. coli* are also included herein.

Alternatively, insect cells may be used as host cells. In a preferred embodiment, the polypeptides of the invention are expressed using a baculovirus expression system (*see*, Luckow *et al.*, *Bio/Technology*, 1988, 6, 47, Baculovirus Expression Vectors: A Laboratory Manual, O'Rielly *et al.* (Eds.), W.H. Freeman and Company, New York, 1992, and U.S. Patent No. 4,879,236, each of which is incorporated herein by reference in its entirety). In addition, the MAXBAC™ complete baculovirus expression system (Invitrogen) can, for example, be used for production in insect cells.

Host cells of the invention are a valuable source of immunogen for development of antibodies specifically immunoreactive with nGPCR-x. Host cells of the invention are also useful in methods for the large-scale production of nGPCR-x polypeptides wherein the cells are grown in a suitable culture medium and the desired polypeptide products are isolated from the cells, or from the medium in which the cells are grown, by purification methods known in the art, *e.g.*, conventional chromatographic methods including immunoaffinity chromatography, receptor affinity chromatography, hydrophobic interaction chromatography, lectin affinity chromatography, size exclusion filtration, cation or anion exchange chromatography, high

pressure liquid chromatography (HPLC), reverse phase HPLC, and the like. Still other methods of purification include those methods wherein the desired protein is expressed and purified as a fusion protein having a specific tag, label, or chelating moiety that is recognized by a specific binding partner or agent. The purified protein can be cleaved to yield the desired protein, or can be left as an intact fusion protein. Cleavage of the fusion component may produce a form of the desired protein having additional amino acid residues as a result of the cleavage process.

Knowledge of nGPCR-x DNA sequences allows for modification of cells to permit, or increase, expression of endogenous nGPCR-x. Cells can be modified (*e.g.*, by homologous recombination) to provide increased expression by replacing, in whole or in part, the naturally occurring nGPCR-x promoter with all or part of a heterologous promoter so that the cells express nGPCR-x at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to endogenous nGPCR-x encoding sequences. (See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955.) It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (*e.g.*, *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamoyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the nGPCR-x coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the nGPCR-x coding sequences in the cells.

### **Knock-outs**

The DNA sequence information provided by the present invention also makes possible the development (*e.g.*, by homologous recombination or "knock-out" strategies; see Capecchi, *Science* 244:1288-1292 (1989), which is incorporated herein by reference in its entirety) of animals that fail to express functional nGPCR-x or that express a variant of nGPCR-x. Such animals (especially small laboratory animals such as rats, rabbits, and mice) are useful as models for studying the *in vivo* activities of nGPCR-x and modulators of nGPCR-x.

### **Antisense**

Also made available by the invention are anti-sense polynucleotides that recognize and hybridize to polynucleotides encoding nGPCR-x. Full-length and fragment anti-sense polynucleotides are provided. Fragment antisense molecules of the invention include (i) those that specifically recognize and hybridize to nGPCR-x RNA (as determined by sequence comparison of DNA encoding nGPCR-x to DNA encoding other known molecules). Identification of sequences unique to nGPCR-x encoding polynucleotides can be deduced

through use of any publicly available sequence database, and/or through use of commercially available sequence comparison programs. After identification of the desired sequences, isolation through restriction digestion or amplification using any of the various polymerase chain reaction techniques well known in the art can be performed. Anti-sense polynucleotides are particularly relevant to regulating expression of nGPCR-x by those cells expressing nGPCR-x mRNA.

Antisense nucleic acids (preferably 10 to 30 base-pair oligonucleotides) capable of specifically binding to nGPCR-x expression control sequences or nGPCR-x RNA are introduced into cells (e.g., by a viral vector or colloidal dispersion system such as a liposome). The antisense nucleic acid binds to the nGPCR-x target nucleotide sequence in the cell and prevents transcription and/or translation of the target sequence. Phosphorothioate and methylphosphonate antisense oligonucleotides are specifically contemplated for therapeutic use by the invention. The antisense oligonucleotides may be further modified by adding poly-L-lysine, transferrin polylysine, or cholesterol moieties at their 5' end. Suppression of nGPCR-x expression at either the transcriptional or translational level is useful to generate cellular or animal models for diseases/conditions characterized by aberrant nGPCR-x expression.

Antisense oligonucleotides, or fragments of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or sequences complementary or homologous thereto, derived from the nucleotide sequences of the present invention encoding nGPCR-x are useful as diagnostic tools for probing gene expression in various tissues. For example, tissue can be probed *in situ* with oligonucleotide probes carrying detectable groups by conventional autoradiography techniques to investigate native expression of this enzyme or pathological conditions relating thereto. Antisense oligonucleotides are preferably directed to regulatory regions of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or mRNA corresponding thereto, including, but not limited to, the initiation codon, TATA box, enhancer sequences, and the like.

### **Transcription factors**

The nGPCR-x sequences taught in the present invention facilitate the design of novel transcription factors for modulating nGPCR-x expression in native cells and animals, and cells transformed or transfected with nGPCR-x polynucleotides. For example, the Cys<sub>2</sub>-His<sub>2</sub> zinc finger proteins, which bind DNA via their zinc finger domains, have been shown to be amenable to structural changes that lead to the recognition of different target sequences. These artificial zinc finger proteins recognize specific target sites with high affinity and low dissociation constants, and are able to act as gene switches to modulate gene expression. Knowledge of the particular nGPCR-x target sequence of the present invention facilitates the engineering of zinc

finger proteins specific for the target sequence using known methods such as a combination of structure-based modeling and screening of phage display libraries (Segal *et al.*, *Proc. Natl. Acad. Sci. USA* 96:2758-2763 (1999); Liu *et al.*, *Proc. Natl. Acad. Sci. USA* 94:5525-5530 (1997); Greisman *et al.*, *Science* 275:657-661 (1997); Choo *et al.*, *J. Mol. Biol.* 273:525-532 (1997)). Each zinc finger domain usually recognizes three or more base pairs. Since a recognition sequence of 18 base pairs is generally sufficient in length to render it unique in any known genome, a zinc finger protein consisting of 6 tandem repeats of zinc fingers would be expected to ensure specificity for a particular sequence (Segal *et al.*). The artificial zinc finger repeats, designed based on nGPCR-x sequences, are fused to activation or repression domains to promote or suppress nGPCR-x expression (Liu *et al.*). Alternatively, the zinc finger domains can be fused to the TATA box-binding factor (TBP) with varying lengths of linker region between the zinc finger peptide and the TBP to create either transcriptional activators or repressors (Kim *et al.*, *Proc. Natl. Acad. Sci. USA* 94:3616-3620 (1997). Such proteins and polynucleotides that encode them, have utility for modulating nGPCR-x expression *in vivo* in both native cells, animals and humans; and/or cells transfected with nGPCR-x-encoding sequences. The novel transcription factor can be delivered to the target cells by transfecting constructs that express the transcription factor (gene therapy), or by introducing the protein. Engineered zinc finger proteins can also be designed to bind RNA sequences for use in therapeutics as alternatives to antisense or catalytic RNA methods (McColl *et al.*, *Proc. Natl. Acad. Sci. USA* 96:9521-9526 (1997); Wu *et al.*, *Proc. Natl. Acad. Sci. USA* 92:344-348 (1995)). The present invention contemplates methods of designing such transcription factors based on the gene sequence of the invention, as well as customized zinc finger proteins, that are useful to modulate nGPCR-x expression in cells (native or transformed) whose genetic complement includes these sequences.

### Polypeptides

The invention also provides purified and isolated mammalian nGPCR-x polypeptides encoded by a polynucleotide of the invention. Presently preferred is a human nGPCR-x polypeptide comprising the amino acid sequence set out in sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, or fragments thereof comprising an epitope specific to the polypeptide. By "epitope specific to" is meant a portion of the nGPCR receptor that is recognizable by an antibody that is specific for the nGPCR, as defined in detail below.

Although the sequences provided are particular human sequences, the invention is intended to include within its scope other human allelic variants; non-human mammalian forms of nGPCR-x, and other vertebrate forms of nGPCR-x.

It will be appreciated that extracellular epitopes are particularly useful for generating and screening for antibodies and other binding compounds that bind to receptors such as nGPCR-x. Thus, in another preferred embodiment, the invention provides a purified and isolated polypeptide comprising at least one extracellular domain (e.g., the N-terminal extracellular domain or one of the three extracellular loops) of nGPCR-x. Purified and isolated polypeptides comprising the N-terminal extracellular domain of nGPCR-x are highly preferred. Also preferred is a purified and isolated polypeptide comprising a nGPCR-x fragment selected from the group consisting of the N-terminal extracellular domain of nGPCR-x, transmembrane domains of nGPCR-x, an extracellular loop connecting transmembrane domains of nGPCR-x, an intracellular loop connecting transmembrane domains of nGPCR-x, the C-terminal cytoplasmic region of nGPCR-x, and fusions thereof. Such fragments may be continuous portions of the native receptor. However, it will also be appreciated that knowledge of the nGPCR-x gene and protein sequences as provided herein permits recombining of various domains that are not contiguous in the native protein. Using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, *Comput. Appl. Biosci.* 5:527-535 (1994)), nGPCR-x was shown to contain transmembrane-spanning domains.

The invention also embraces polypeptides that have at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55% or at least 50% identity and/or homology to the preferred polypeptide of the invention.

Percent amino acid sequence "identity" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning both sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Percent sequence "homology" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and also considering any conservative substitutions as part of the sequence identity.

In one aspect, percent homology is calculated as the percentage of amino acid residues in the smaller of two sequences which align with identical amino acid residue in the sequence being compared, when four gaps in a length of 100 amino acids may be introduced to maximize alignment (Dayhoff, in *Atlas of Protein Sequence and Structure*, Vol. 5, p. 124, National

Biochemical Research Foundation, Washington, D.C. (1972), incorporated herein by reference in its entirety).

Polypeptides of the invention may be isolated from natural cell sources or may be chemically synthesized, but are preferably produced by recombinant procedures involving host cells of the invention. Use of mammalian host cells is expected to provide for such post-translational modifications (*e.g.*, glycosylation, truncation, lipidation, and phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. Glycosylated and non-glycosylated forms of nGPCR-x polypeptides are embraced by the invention.

The invention also embraces variant (or analog) nGPCR-x polypeptides. In one example, insertion variants are provided wherein one or more amino acid residues supplement a nGPCR-x amino acid sequence. Insertions may be located at either or both termini of the protein, or may be positioned within internal regions of the nGPCR-x amino acid sequence. Insertional variants with additional residues at either or both termini can include, for example, fusion proteins and proteins including amino acid tags or labels.

Insertion variants include nGPCR-x polypeptides wherein one or more amino acid residues are added to a nGPCR-x acid sequence or to a biologically active fragment thereof.

Variant products of the invention also include mature nGPCR-x products, *i.e.*, nGPCR-x products wherein leader or signal sequences are removed, with additional amino terminal residues. The additional amino terminal residues may be derived from another protein, or may include one or more residues that are not identifiable as being derived from specific proteins. nGPCR-x products with an additional methionine residue at position -1 (Met<sup>-1</sup>-nGPCR-x) are contemplated, as are variants with additional methionine and lysine residues at positions -2 and -1 (Met<sup>-2</sup>-Lys<sup>-1</sup>-nGPCR-x). Variants of nGPCR-x with additional Met, Met-Lys, Lys residues (or one or more basic residues in general) are particularly useful for enhanced recombinant protein production in bacterial host cells.

The invention also embraces nGPCR-x variants having additional amino acid residues that result from use of specific expression systems. For example, use of commercially available vectors that express a desired polypeptide as part of a glutathione-S-transferase (GST) fusion product provides the desired polypeptide having an additional glycine residue at position -1 after cleavage of the GST component from the desired polypeptide. Variants that result from expression in other vector systems are also contemplated.

Insertional variants also include fusion proteins wherein the amino terminus and/or the carboxy terminus of nGPCR-x is/are fused to another polypeptide.



In another aspect, the invention provides deletion variants wherein one or more amino acid residues in a nGPCR-x polypeptide are removed. Deletions can be effected at one or both termini of the nGPCR-x polypeptide, or with removal of one or more non-terminal amino acid residues of nGPCR-x. Deletion variants, therefore, include all fragments of a nGPCR-x

5 polypeptide.

The invention also embraces polypeptide fragments of sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, wherein the fragments maintain biological (*e.g.*, ligand binding and/or intracellular signaling) immunological properties of a nGPCR-x polypeptide.

10 In one preferred embodiment of the invention, an isolated nucleic acid molecule comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, and fragments thereof, wherein the nucleic acid molecule encoding at least a portion of nGPCR-x. In a more preferred embodiment, the isolated nucleic acid molecule comprises a

15 sequence that encodes a polypeptide comprising sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, and fragments thereof.

As used in the present invention, polypeptide fragments comprise at least 5, 10, 15, 20, 25, 30, 35, or 40 consecutive amino acids of sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120. Preferred polypeptide fragments display antigenic

20 properties unique to, or specific for, human nGPCR-x and its allelic and species homologs. Fragments of the invention having the desired biological and immunological properties can be prepared by any of the methods well known and routinely practiced in the art.

In still another aspect, the invention provides substitution variants of nGPCR-x polypeptides. Substitution variants include those polypeptides wherein one or more amino acid

25 residues of a nGPCR-x polypeptide are removed and replaced with alternative residues. In one aspect, the substitutions are conservative in nature; however, the invention embraces substitutions that are also non-conservative. Conservative substitutions for this purpose may be defined as set out in Tables 2, 3, or 4 below.

Variant polypeptides include those wherein conservative substitutions have been

30 introduced by modification of polynucleotides encoding polypeptides of the invention. Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative

substitutions are set out in Table 2 (from WO 97/09433, page 10, published March 13, 1997 (PCT/GB96/02197, filed 9/6/96), immediately below.

Table 2

## Conservative Substitutions I

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Aliphatic	
Non-polar	G A P I L V
Polar - uncharged	C S T M N Q
Polar - charged	D E K R
Aromatic	H F W Y
Other	N Q D E

Alternatively, conservative amino acids can be grouped as described in Lehninger, (*Biochemistry*, Second Edition; Worth Publishers, Inc. NY, NY (1975), pp.71-77) as set out in Table 3, below.

Table 3

## Conservative Substitutions II

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Non-polar (hydrophobic)	
A. Aliphatic:	A L I V P
B. Aromatic:	F W
C. Sulfur-containing:	M
D. Borderline:	G
Uncharged-polar	
A. Hydroxyl:	S T Y
B. Amides:	N Q
C. Sulfhydryl:	C
D. Borderline:	G
Positively Charged (Basic):	K R H
Negatively Charged (Acidic):	D E

As still another alternative, exemplary conservative substitutions are set out in Table 4,

below.

Table 4

## Conservative Substitutions III

<b>Original Residue</b>	<b>Exemplary Substitution</b>
Ala (A)	Val, Leu, Ile

Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg
Ile (I)	Leu, Val, Met, Ala, Phe,
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	Gly
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

It should be understood that the definition of polypeptides of the invention is intended to include polypeptides bearing modifications other than insertion, deletion, or substitution of amino acid residues. By way of example, the modifications may be covalent in nature, and include for example, chemical bonding with polymers, lipids, other organic, and inorganic moieties. Such derivatives may be prepared to increase circulating half-life of a polypeptide, or may be designed to improve the targeting capacity of the polypeptide for desired cells, tissues, or organs. Similarly, the invention further embraces nGPCR-x polypeptides that have been covalently modified to include one or more water-soluble polymer attachments such as polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol. Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

In a related embodiment, the present invention provides compositions comprising purified polypeptides of the invention. Preferred compositions comprise, in addition to the polypeptide of the invention, a pharmaceutically acceptable (*i.e.*, sterile and non-toxic) liquid, semisolid, or solid diluent that serves as a pharmaceutical vehicle, excipient, or medium. Any diluent known in the art may be used. Exemplary diluents include, but are not limited to, water, saline solutions, polyoxyethylene sorbitan monolaurate, magnesium stearate, methyl- and

propylhydroxybenzoate, talc, alginates, starches, lactose, sucrose, dextrose, sorbitol, mannitol, glycerol, calcium phosphate, mineral oil, and cocoa butter.

Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in assays of the invention and in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

The G protein-coupled receptor functions through a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) coupled to the intracellular portion of the G protein-coupled receptor molecule. Accordingly, the G protein-coupled receptor has a specific affinity to G protein. G proteins specifically bind to guanine nucleotides. Isolation of G proteins provides a means to isolate guanine nucleotides. G Proteins may be isolated using commercially available anti-G protein antibodies or isolated G protein-coupled receptors. Similarly, G proteins may be detected in a sample isolated using commercially available detectable anti-G protein antibodies or isolated G protein-coupled receptors.

According to the present invention, the isolated n-GPCR-x proteins of the present invention are useful to isolate and purify G proteins from samples such as cell lysates. Example 14 below sets forth an example of isolation of G proteins using isolated n-GPCR-x proteins. Such methodology may be used in place of the use of commercially available anti-G protein antibodies which are used to isolate G proteins. Moreover, G proteins may be detected using n-GPCR-x proteins in place of commercially available detectable anti-G protein antibodies. Since n-GPCR-x proteins specifically bind to G proteins, they can be employed in any specific use where G protein specific affinity is required such as those uses where commercially available anti-G protein antibodies are employed.

## **Antibodies**

Also comprehended by the present invention are antibodies (*e.g.*, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention) specific for nGPCR-x or fragments thereof. Preferred antibodies of the invention are human antibodies that are produced and identified according to methods described in WO93/11236, published June 20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')<sub>2</sub>, and F<sub>v</sub>, are also provided by the invention. The term "specific for," when used to

describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind nGPCR-x polypeptides exclusively (*i.e.*, are able to distinguish nGPCR-x polypeptides from other known GPCR polypeptides by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity between nGPCR-x and such polypeptides). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and, in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow *et al.* (Eds.), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6. Antibodies that recognize and bind fragments of the nGPCR-x polypeptides of the invention are also contemplated, provided that the antibodies are specific for nGPCR-x polypeptides. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

The invention provides an antibody that is specific for the nGPCR-x of the invention. Antibody specificity is described in greater detail below. However, it should be emphasized that antibodies that can be generated from polypeptides that have previously been described in the literature and that are capable of fortuitously cross-reacting with nGPCR-x (*e.g.*, due to the fortuitous existence of a similar epitope in both polypeptides) are considered "cross-reactive" antibodies. Such cross-reactive antibodies are not antibodies that are "specific" for nGPCR-x. The determination of whether an antibody is specific for nGPCR-x or is cross-reactive with another known receptor is made using any of several assays, such as Western blotting assays, that are well known in the art. For identifying cells that express nGPCR-x and also for modulating nGPCR-x-ligand binding activity, antibodies that specifically bind to an extracellular epitope of the nGPCR-x are preferred.

In one preferred variation, the invention provides monoclonal antibodies. Hybridomas that produce such antibodies also are intended as aspects of the invention. In yet another variation, the invention provides a humanized antibody. Humanized antibodies are useful for *in vivo* therapeutic indications.

In another variation, the invention provides a cell-free composition comprising polyclonal antibodies, wherein at least one of the antibodies is an antibody of the invention specific for nGPCR-x. Antisera isolated from an animal is an exemplary composition, as is a

composition comprising an antibody fraction of an antisera that has been resuspended in water or in another diluent, excipient, or carrier.

In still another related embodiment, the invention provides an anti-idiotypic antibody specific for an antibody that is specific for nGPCR-x.

5 It is well known that antibodies contain relatively small antigen binding domains that can be isolated chemically or by recombinant techniques. Such domains are useful nGPCR-x binding molecules themselves, and also may be reintroduced into human antibodies, or fused to toxins or other polypeptides. Thus, in still another embodiment, the invention provides a polypeptide comprising a fragment of a nGPCR-x-specific antibody, wherein the fragment and  
10 the polypeptide bind to the nGPCR-x. By way of non-limiting example, the invention provides polypeptides that are single chain antibodies and CDR-grafted antibodies.

Non-human antibodies may be humanized by any of the methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to  
15 modulate affinity or immunogenicity.

Antibodies of the invention are useful for, *e.g.*, therapeutic purposes (by modulating activity of nGPCR-x), diagnostic purposes to detect or quantitate nGPCR-x, and purification of nGPCR-x. Kits comprising an antibody of the invention for any of the purposes described herein are also comprehended. In general, a kit of the invention also includes a control antigen for  
20 which the antibody is immunospecific.

### Compositions

Mutations in the nGPCR-x gene that result in loss of normal function of the nGPCR-x gene product underlie nGPCR-x-related human disease states. The invention comprehends gene therapy to restore nGPCR-x activity to treat those disease states. Delivery of a functional  
25 nGPCR-x gene to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (*e.g.*, adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (*e.g.*, liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, *Science*, 244: 1275-1281 (1989); Verma, *Scientific American*: 68-84 (1990); and Miller, *Nature*, 357: 455-460 (1992). Alternatively, it is  
30 contemplated that in other human disease states, preventing the expression of, or inhibiting the activity of, nGPCR-x will be useful in treating disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of nGPCR-x.

Another aspect of the present invention is directed to compositions, including pharmaceutical compositions, comprising any of the nucleic acid molecules or recombinant expression vectors described above and an acceptable carrier or diluent. Preferably, the carrier or diluent is pharmaceutically acceptable. Suitable carriers are described in the most recent  
5 edition of *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is incorporated herein by reference in its entirety. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The formulations are sterilized by commonly used techniques.

10 Also within the scope of the invention are compositions comprising polypeptides, polynucleotides, or antibodies of the invention that have been formulated with, *e.g.*, a pharmaceutically acceptable carrier.

The invention also provides methods of using antibodies of the invention. For example, the invention provides a method for modulating ligand binding of a nGPCR-x comprising the  
15 step of contacting the nGPCR-x with an antibody specific for the nGPCR-x, under conditions wherein the antibody binds the receptor.

GPCRs that may be expressed in the brain, such as nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70, provide an indication that aberrant nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 signaling activity may correlate with one or more neurological or psychological disorders. The  
20 invention also provides a method for treating a neurological or psychiatric disorder comprising the step of administering to a mammal in need of such treatment an amount of an antibody-like polypeptide of the invention that is sufficient to modulate ligand binding to a nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 in neurons of the mammal. nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 may also be expressed in other tissues, including but not limited to, peripheral blood  
25 lymphocytes, pancreas, ovary, uterus, testis, salivary gland, thyroid gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues. Within the brain, nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 mRNA transcripts may be found in many tissues, including, but not limited to, frontal lobe, hypothalamus, pons, cerebellum, caudate nucleus, and medulla. Tissues and brain regions where  
30 specific nGPCRs of the present invention are expressed are identified in the Examples below.

#### Kits

The present invention is also directed to kits, including pharmaceutical kits. The kits can comprise any of the nucleic acid molecules described above, any of the polypeptides described above, or any antibody which binds to a polypeptide of the invention as described above, as well

as a negative control. The kit preferably comprises additional components, such as, for example, instructions, solid support, reagents helpful for quantification, and the like.

In another aspect, the invention features methods for detection of a polypeptide in a sample as a diagnostic tool for diseases or disorders, wherein the method comprises the steps of:

- 5 (a) contacting the sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a polypeptide having sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and (b) detecting the presence or amount of the probe:target region hybrid as an  
10 indication of the disease.

In preferred embodiments of the invention, the disease is selected from the group consisting of thyroid disorders (*e.g.* thyreotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain  
15 including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including attention deficit disorder (ADD) and attention deficit-hyperactivity disorder (ADHD), and degenerative  
20 disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*,  
25 different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); and sexual dysfunction, among others.

As described above and in Example 5 below, the genes encoding nGPCR-42, 46, 48, 49,  
30 51, 52, 61, 63, and 70 have been detected in brain tissue indicating that these n-GPCR-x proteins are neuroreceptors. Kits may be designed to detect either expression of polynucleotides encoding these proteins or the proteins themselves in order to identify tissue as being neurological. For example, oligonucleotide hybridization kits can be provided which include a container having an oligonucleotide probe specific for the n-GPCR-x-specific DNA and



optionally, containers with positive and negative controls and/or instructions. Similarly, PCR kits can be provided which include a container having primers specific for the n-GPCR-x-specific sequences, DNA and optionally, containers with size markers, positive and negative controls and/or instructions.

5 Hybridization conditions should be such that hybridization occurs only with the genes in the presence of other nucleic acid molecules. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such conditions prevent hybridization of nucleic acids having 1 or 2 mismatches out of 20 contiguous nucleotides. Such conditions are defined supra.

10 The diseases for which detection of genes in a sample could be diagnostic include diseases in which nucleic acid (DNA and/or RNA) is amplified in comparison to normal cells. By "amplification" is meant increased numbers of DNA or RNA in a cell compared with normal cells.

The diseases that could be diagnosed by detection of nucleic acid in a sample preferably  
15 include central nervous system and metabolic diseases. The test samples suitable for nucleic acid probing methods of the present invention include, for example, cells or nucleic acid extracts of cells, or biological fluids. The samples used in the above-described methods will vary based on the assay format, the detection method and the nature of the tissues, cells or extracts to be assayed. Methods for preparing nucleic acid extracts of cells are well known in the art and can  
20 be readily adapted in order to obtain a sample that is compatible with the method utilized.

Alternatively, immunoassay kits can be provided which have containers container having antibodies specific for the n-GPCR-x-protein and optionally, containers with positive and negative controls and/or instructions.

Kits may also be provided useful in the identification of GPCR binding partners such as  
25 natural ligands or modulators (agonists or antagonists). Substances useful for treatment of disorders or diseases preferably show positive results in one or more *in vitro* assays for an activity corresponding to treatment of the disease or disorder in question. Substances that modulate the activity of the polypeptides preferably include, but are not limited to, antisense oligonucleotides, agonists and antagonists, and inhibitors of protein kinases.

### 30 **Methods of inducing immune response**

Another aspect of the present invention is directed to methods of inducing an immune response in a mammal against a polypeptide of the invention by administering to the mammal an amount of the polypeptide sufficient to induce an immune response. The amount will be

dependent on the animal species, size of the animal, and the like but can be determined by those skilled in the art.

### Methods of identifying ligands

The invention also provides assays to identify compounds that bind nGPCR-x. One such assay comprises the steps of: (a) contacting a composition comprising a nGPCR-x with a compound suspected of binding nGPCR-x; and (b) measuring binding between the compound and nGPCR-x. In one variation, the composition comprises a cell expressing nGPCR-x on its surface. In another variation, isolated nGPCR-x or cell membranes comprising nGPCR-x are employed. The binding may be measured directly, *e.g.*, by using a labeled compound, or may be measured indirectly by several techniques, including measuring intracellular signaling of nGPCR-x induced by the compound (or measuring changes in the level of nGPCR-x signaling). Following steps (a) and (b), compounds identified as binding nGPCR-x can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate binding to nGPCR-x.

Specific binding molecules, including natural ligands and synthetic compounds, can be identified or developed using isolated or recombinant nGPCR-x products, nGPCR-x variants, or preferably, cells expressing such products. Binding partners are useful for purifying nGPCR-x products and detection or quantification of nGPCR-x products in fluid and tissue samples using known immunological procedures. Binding molecules are also manifestly useful in modulating (*i.e.*, blocking, inhibiting or stimulating) biological activities of nGPCR-x, especially those activities involved in signal transduction.

The DNA and amino acid sequence information provided by the present invention also makes possible identification of binding partner compounds with which a nGPCR-x polypeptide or polynucleotide will interact. Methods to identify binding partner compounds include solution assays, *in vitro* assays wherein nGPCR-x polypeptides are immobilized, and cell-based assays. Identification of binding partner compounds of nGPCR-x polypeptides provides candidates for therapeutic or prophylactic intervention in pathologies associated with nGPCR-x normal and aberrant biological activity.

The invention includes several assay systems for identifying nGPCR-x binding partners. In solution assays, methods of the invention comprise the steps of (a) contacting a nGPCR-x polypeptide with one or more candidate binding partner compounds and (b) identifying the compounds that bind to the nGPCR-x polypeptide. Identification of the compounds that bind the nGPCR-x polypeptide can be achieved by isolating the nGPCR-x polypeptide/binding partner complex, and separating the binding partner compound from the nGPCR-x polypeptide.

An additional step of characterizing the physical, biological, and/or biochemical properties of the binding partner compound is also comprehended in another embodiment of the invention, wherein compounds identified as binding nGPCR-x can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate binding to

5 nGPCR-x. In one aspect, the nGPCR-x polypeptide/binding partner complex is isolated using an antibody immunospecific for either the nGPCR-x polypeptide or the candidate binding partner compound.

In still other embodiments, either the nGPCR-x polypeptide or the candidate binding partner compound comprises a label or tag that facilitates its isolation, and methods of the invention to identify binding partner compounds include a step of isolating the nGPCR-x polypeptide/binding partner complex through interaction with the label or tag. An exemplary tag of this type is a poly-histidine sequence, generally around six histidine residues, that permits isolation of a compound so labeled using nickel chelation. Other labels and tags, such as the FLAG® tag (Eastman Kodak, Rochester, NY), well known and routinely used in the art, are

15 embraced by the invention.

In one variation of an *in vitro* assay, the invention provides a method comprising the steps of (a) contacting an immobilized nGPCR-x polypeptide with a candidate binding partner compound and (b) detecting binding of the candidate compound to the nGPCR-x polypeptide. In an alternative embodiment, the candidate binding partner compound is immobilized and binding of nGPCR-x is detected. Immobilization is accomplished using any of the methods well known in the art, including covalent bonding to a support, a bead, or a chromatographic resin, as well as non-covalent, high affinity interactions such as antibody binding, or use of streptavidin/biotin binding wherein the immobilized compound includes a biotin moiety. Detection of binding can be accomplished (i) using a radioactive label on the compound that is not immobilized, (ii) using

20 of a fluorescent label on the non-immobilized compound, (iii) using an antibody immunospecific for the non-immobilized compound, (iv) using a label on the non-immobilized compound that excites a fluorescent support to which the immobilized compound is attached, as well as other techniques well known and routinely practiced in the art.

The invention also provides cell-based assays to identify binding partner compounds of a nGPCR-x polypeptide. In one embodiment, the invention provides a method comprising the steps of contacting a nGPCR-x polypeptide expressed on the surface of a cell with a candidate binding partner compound and detecting binding of the candidate binding partner compound to the nGPCR-x polypeptide. In a preferred embodiment, the detection comprises detecting a calcium flux or other physiological event in the cell caused by the binding of the molecule.

30

Another aspect of the present invention is directed to methods of identifying compounds that bind to either nGPCR-x or nucleic acid molecules encoding nGPCR-x, comprising contacting nGPCR-x, or a nucleic acid molecule encoding the same, with a compound, and determining whether the compound binds nGPCR-x or a nucleic acid molecule encoding the same. Binding can be determined by binding assays which are well known to the skilled artisan, including, but not limited to, gel-shift assays, Western blots, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, ELISA, and the like, which are described in, for example, *Current Protocols in Molecular Biology*, 1999, John Wiley & Sons, NY, which is incorporated herein by reference in its entirety. The compounds to be screened include (which may include compounds which are suspected to bind nGPCR-x, or a nucleic acid molecule encoding the same), but are not limited to, extracellular, intracellular, biologic or chemical origin. The methods of the invention also embrace ligands, especially neuropeptides, that are attached to a label, such as a radiolabel (e.g.,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^3\text{H}$ ), a fluorescence label, a chemiluminescent label, an enzymic label and an immunogenic label. Modulators falling within the scope of the invention include, but are not limited to, non-peptide molecules such as non-peptide mimetics, non-peptide allosteric effectors, and peptides. The nGPCR-x polypeptide or polynucleotide employed in such a test may either be free in solution, attached to a solid support, borne on a cell surface or located intracellularly or associated with a portion of a cell. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

In another embodiment of the invention, high throughput screening for compounds having suitable binding affinity to nGPCR-x is employed. Briefly, large numbers of different test compounds are synthesized on a solid substrate. The peptide test compounds are contacted with nGPCR-x and washed. Bound nGPCR-x is then detected by methods well known in the art. Purified polypeptides of the invention can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the protein and immobilize it on the solid support.

Generally, an expressed nGPCR-x can be used for HTS binding assays in conjunction with its defined ligand, in this case the corresponding neuropeptide that activates it. The identified peptide is labeled with a suitable radioisotope, including, but not limited to,  $^{125}\text{I}$ ,  $^3\text{H}$ ,  $^{35}\text{S}$  or  $^{32}\text{P}$ , by methods that are well known to those skilled in the art. Alternatively, the peptides

may be labeled by well-known methods with a suitable fluorescent derivative (Baindur *et al.*, *Drug Dev. Res.*, **1994**, *33*, 373-398; Rogers, *Drug Discovery Today*, **1997**, *2*, 156-160).

Radioactive ligand specifically bound to the receptor in membrane preparations made from the cell line expressing the recombinant protein can be detected in HTS assays in one of several  
5 standard ways, including filtration of the receptor-ligand complex to separate bound ligand from unbound ligand (Williams, *Med. Res. Rev.*, **1991**, *11*, 147-184; Sweetnam *et al.*, *J. Natural Products*, **1993**, *56*, 441-455). Alternative methods include a scintillation proximity assay (SPA) or a FlashPlate format in which such separation is unnecessary (Nakayama, *Cur. Opinion Drug Disc. Dev.*, **1998**, *1*, 85-91 Bossé *et al.*, *J. Biomolecular Screening*, **1998**, *3*, 285-292.). Binding  
10 of fluorescent ligands can be detected in various ways, including fluorescence energy transfer (FRET), direct spectrophotofluorometric analysis of bound ligand, or fluorescence polarization (Rogers, *Drug Discovery Today*, **1997**, *2*, 156-160; Hill, *Cur. Opinion Drug Disc. Dev.*, **1998**, *1*, 92-97).

Other assays may be used to identify specific ligands of a nGPCR-x receptor, including  
15 assays that identify ligands of the target protein through measuring direct binding of test ligands to the target protein, as well as assays that identify ligands of target proteins through affinity ultrafiltration with ion spray mass spectroscopy/HPLC methods or other physical and analytical methods. Alternatively, such binding interactions are evaluated indirectly using the yeast two-hybrid system described in Fields *et al.*, *Nature*, 340:245-246 (1989), and Fields *et al.*, *Trends in*  
20 *Genetics*, 10:286-292 (1994), both of which are incorporated herein by reference in its entirety. The two-hybrid system is a genetic assay for detecting interactions between two proteins or polypeptides. It can be used to identify proteins that bind to a known protein of interest, or to delineate domains or residues critical for an interaction. Variations on this methodology have been developed to clone genes that encode DNA binding proteins, to identify peptides that bind  
25 to a protein, and to screen for drugs. The two-hybrid system exploits the ability of a pair of interacting proteins to bring a transcription activation domain into close proximity with a DNA binding domain that binds to an upstream activation sequence (UAS) of a reporter gene, and is generally performed in yeast. The assay requires the construction of two hybrid genes encoding (1) a DNA-binding domain that is fused to a first protein and (2) an activation domain fused to a  
30 second protein. The DNA-binding domain targets the first hybrid protein to the UAS of the reporter gene; however, because most proteins lack an activation domain, this DNA-binding hybrid protein does not activate transcription of the reporter gene. The second hybrid protein, which contains the activation domain, cannot by itself activate expression of the reporter gene because it does not bind the UAS. However, when both hybrid proteins are present, the

noncovalent interaction of the first and second proteins tethers the activation domain to the UAS, activating transcription of the reporter gene. For example, when the first protein is a GPCR gene product, or fragment thereof, that is known to interact with another protein or nucleic acid, this assay can be used to detect agents that interfere with the binding interaction.

- 5 Expression of the reporter gene is monitored as different test agents are added to the system. The presence of an inhibitory agent results in lack of a reporter signal.

The yeast two-hybrid assay can also be used to identify proteins that bind to the gene product. In an assay to identify proteins that bind to a nGPCR-x receptor, or fragment thereof, a fusion polynucleotide encoding both a nGPCR-x receptor (or fragment) and a UAS binding domain (*i.e.*, a first protein) may be used. In addition, a large number of hybrid genes each encoding a different second protein fused to an activation domain are produced and screened in the assay. Typically, the second protein is encoded by one or more members of a total cDNA or genomic DNA fusion library, with each second protein-coding region being fused to the activation domain. This system is applicable to a wide variety of proteins, and it is not even necessary to know the identity or function of the second binding protein. The system is highly sensitive and can detect interactions not revealed by other methods; even transient interactions may trigger transcription to produce a stable mRNA that can be repeatedly translated to yield the reporter protein.

Other assays may be used to search for agents that bind to the target protein. One such screening method to identify direct binding of test ligands to a target protein is described in U.S. Patent No. 5,585,277, incorporated herein by reference in its entirety. This method relies on the principle that proteins generally exist as a mixture of folded and unfolded states, and continually alternate between the two states. When a test ligand binds to the folded form of a target protein (*i.e.*, when the test ligand is a ligand of the target protein), the target protein molecule bound by the ligand remains in its folded state. Thus, the folded target protein is present to a greater extent in the presence of a test ligand which binds the target protein, than in the absence of a ligand. Binding of the ligand to the target protein can be determined by any method that distinguishes between the folded and unfolded states of the target protein. The function of the target protein need not be known in order for this assay to be performed. Virtually any agent can be assessed by this method as a test ligand, including, but not limited to, metals, polypeptides, proteins, lipids, polysaccharides, polynucleotides and small organic molecules.

Another method for identifying ligands of a target protein is described in Wieboldt *et al.*, *Anal. Chem.*, 69:1683-1691 (1997), incorporated herein by reference in its entirety. This technique screens combinatorial libraries of 20-30 agents at a time in solution phase for binding

to the target protein. Agents that bind to the target protein are separated from other library components by simple membrane washing. The specifically selected molecules that are retained on the filter are subsequently liberated from the target protein and analyzed by HPLC and pneumatically assisted electrospray (ion spray) ionization mass spectroscopy. This procedure  
5 selects library components with the greatest affinity for the target protein, and is particularly useful for small molecule libraries.

Determining whether a test compound binds to nGPCR-51 can also be accomplished by measuring the intrinsic fluorescence of nGPCR-51 and determining whether the intrinsic fluorescence is modulated in the presence of the test compound. Preferably, the intrinsic  
10 fluorescence of nGPCR-51 is measured as a function of the tryptophan residue(s) of nGPCR-51. Preferably, fluorescence of nGPCR-51 is measured and compared to the fluorescence intensity of nGPCR-51 in the presence of the test compound, wherein a decrease in fluorescence intensity indicates binding of the test compound to nGPCR-51. Preferred methodology is set forth in "Principles of Fluorescence Spectroscopy" by Joseph R. Lakowicz, New York, Plenum Press,  
15 1983 (ISBN 0306412853) and "Spectrophotometry And Spectrofluorometry" by C.L. Bashford and D.A. Harris Oxford, Washington DC, IRL Press, 1987 (ISBN 0947946691), each of which is incorporated herein by reference in its entirety.

Other embodiments of the invention comprise using competitive screening assays in which neutralizing antibodies capable of binding a polypeptide of the invention specifically  
20 compete with a test compound for binding to the polypeptide. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more antigenic determinants with nGPCR-x. Radiolabeled competitive binding studies are described in A.H. Lin *et al.* *Antimicrobial Agents and Chemotherapy*, 1997, vol. 41, no. 10. pp. 2127-2131, the disclosure of which is incorporated herein by reference in its entirety.

Another aspect of the present invention relates to methods of identifying a compound  
25 that binds to or modulates nGPCR-51. The methods comprise contacting a composition comprising nGPCR-51 and Peptide A with a test compound, or a plurality of test compounds, and determining whether the test compound competes with Peptide A for binding to nGPCR-51. A decrease in the amount of the complex between Peptide A, or a protein homologous thereto,  
30 and nGPCR-51 in the presence of a test compound or compounds confirms that the compound or compounds binds to nGPCR-51. In some embodiments, the affinity or displacement of Peptide A is measured, wherein a low affinity indicates that the test compound interacts with nGPCR-51. In these methods, the composition that comprises nGPCR-51 and Peptide A can be cells. Compounds identified as binding to nGPCR-51 are also expected to modulate nGPCR-51

activity. Binding of a test compound to nGPCR-51 can be determined by any of the binding assays described above.

As described above and in Example 5 below, the genes encoding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, and 70 have been detected in brain tissue indicating that these n-GPCR-x proteins are neuroreceptors. Accordingly, natural binding partners of these molecules include neurotransmitters.

#### Identification of modulating agents

The invention also provides methods for identifying a modulator of binding between a nGPCR-x and a nGPCR-x binding partner, comprising the steps of: (a) contacting a nGPCR-x binding partner and a composition comprising a nGPCR-x in the presence and in the absence of a putative modulator compound; (b) detecting binding between the binding partner and the nGPCR-x; and (c) identifying a putative modulator compound or a modulator compound in view of decreased or increased binding between the binding partner and the nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator. Following steps (a) and (b), compounds identified as modulating binding between nGPCR-x and an nGPCR-x binding partner can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate modulation of binding to nGPCR-x.

nGPCR-x binding partners that stimulate nGPCR-x activity are useful as agonists in disease states or conditions characterized by insufficient nGPCR-x signaling (*e.g.*, as a result of insufficient activity of a nGPCR-x ligand). nGPCR-x binding partners that block ligand-mediated nGPCR-x signaling are useful as nGPCR-x antagonists to treat disease states or conditions characterized by excessive nGPCR-x signaling. In addition nGPCR-x modulators in general, as well as nGPCR-x polynucleotides and polypeptides, are useful in diagnostic assays for such diseases or conditions.

In another aspect, the invention provides methods for treating a disease or abnormal condition by administering to a patient in need of such treatment a substance that modulates the activity or expression of a polypeptide having sequences selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120.

Agents that modulate (*i.e.*, increase, decrease, or block) nGPCR-x activity or expression may be identified by incubating a putative modulator with a cell containing a nGPCR-x polypeptide or polynucleotide and determining the effect of the putative modulator on nGPCR-x activity or expression. The selectivity of a compound that modulates the activity of nGPCR-x can be evaluated by comparing its effects on nGPCR-x to its effect on other GPCR compounds. Following identification of compounds that modulate nGPCR-x activity or expression, such



compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Selective modulators may include, for example, antibodies and other proteins, peptides, or organic molecules that specifically bind to a nGPCR-x polypeptide or a nGPCR-x-encoding nucleic acid. Modulators of nGPCR-x activity will be

5 therapeutically useful in treatment of diseases and physiological conditions in which normal or aberrant nGPCR-x activity is involved. nGPCR-x polynucleotides, polypeptides, and modulators may be used in the treatment of such diseases and conditions as infections, such as viral infections caused by HIV-1 or HIV-2; pain; cancers; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia,

10 hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); Parkinson's disease; and psychotic and neurological disorders, including anxiety, mental disorder, manic depression, schizophrenia, migraine, major depression, attention disorders including ADD and ADHD, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome, among others. nGPCR-x

15 polynucleotides and polypeptides, as well as nGPCR-x modulators, may also be used in diagnostic assays for such diseases or conditions.

Methods of the invention to identify modulators include variations on any of the methods described above to identify binding partner compounds, the variations including techniques wherein a binding partner compound has been identified and the binding assay is carried out in

20 the presence and absence of a candidate modulator. A modulator is identified in those instances where binding between the nGPCR-x polypeptide and the binding partner compound changes in the presence of the candidate modulator compared to binding in the absence of the candidate modulator compound. A modulator that increases binding between the nGPCR-x polypeptide and the binding partner compound is described as an enhancer or activator, and a modulator that

25 decreases binding between the nGPCR-x polypeptide and the binding partner compound is described as an inhibitor. Following identification of modulators, such compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

The invention also comprehends high-throughput screening (HTS) assays to identify

30 compounds that interact with or inhibit biological activity (*i.e.*, affect enzymatic activity, binding activity, *etc.*) of a nGPCR-x polypeptide. HTS assays permit screening of large numbers of compounds in an efficient manner. Cell-based HTS systems are contemplated to investigate nGPCR-x receptor-ligand interaction. HTS assays are designed to identify "hits" or "lead compounds" having the desired property, from which modifications can be designed to improve

the desired property. Chemical modification of the "hit" or "lead compound" is often based on an identifiable structure/activity relationship between the "hit" and the nGPCR-x polypeptide.

Another aspect of the present invention is directed to methods of identifying compounds which modulate (*i.e.*, increase or decrease) activity of nGPCR-x comprising contacting nGPCR-x with a compound, and determining whether the compound modifies activity of nGPCR-x. The activity in the presence of the test compared is measured to the activity in the absence of the test compound. Where the activity of the sample containing the test compound is higher than the activity in the sample lacking the test compound, the compound will have increased activity. Similarly, where the activity of the sample containing the test compound is lower than the activity in the sample lacking the test compound, the compound will have inhibited activity. Following identification of compounds that modulate an activity of nGPCR-x, such compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

The present invention is particularly useful for screening compounds by using nGPCR-x in any of a variety of drug screening techniques. The compounds to be screened include (which may include compounds which are suspected to modulate nGPCR-x activity), but are not limited to, extracellular, intracellular, biologic or chemical origin. The nGPCR-x polypeptide employed in such a test may be in any form, preferably, free in solution, attached to a solid support, borne on a cell surface or located intracellularly. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

The activity of nGPCR-x polypeptides of the invention can be determined by, for example, examining the ability to bind or be activated by chemically synthesized peptide ligands. Alternatively, the activity of nGPCR-x polypeptides can be assayed by examining their ability to bind calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and photons. Alternatively, the activity of the nGPCR-x polypeptides can be determined by examining the activity of effector molecules including, but not limited to, adenylate cyclase, phospholipases and ion channels. Thus, modulators of nGPCR-x polypeptide activity may alter a GPCR receptor function, such as a binding property of a receptor or an activity such as G protein-mediated signal transduction or membrane localization. In various embodiments of the method, the assay may take the form of an ion flux assay, a yeast growth assay, a non-hydrolyzable GTP assay such as a [<sup>35</sup>S]-GTP γS assay, a cAMP assay, an inositol triphosphate assay, a diacylglycerol assay, an Aequorin assay, a

Luciferase assay, a FLIPR assay for intracellular  $\text{Ca}^{2+}$  concentration, a mitogenesis assay, a MAP Kinase activity assay, an arachidonic acid release assay (*e.g.*, using [ $^3\text{H}$ ]-arachidonic acid), and an assay for extracellular acidification rates, as well as other binding or function-based assays of nGPCR-x activity that are generally known in the art. In several of these

embodiments, the invention comprehends the inclusion of any of the G proteins known in the art, such as  $\text{G}_{16}$ ,  $\text{G}_{15}$ , or chimeric  $\text{G}_{\text{qd5}}$ ,  $\text{G}_{\text{qs5}}$ ,  $\text{G}_{\text{qo5}}$ ,  $\text{G}_{\text{q25}}$ , and the like. nGPCR-x activity can be determined by methodologies that are used to assay for FaRP activity, which is well known to those skilled in the art. Biological activities of nGPCR-x receptors according to the invention include, but are not limited to, the binding of a natural or an unnatural ligand, as well as any one of the functional activities of GPCRs known in the art. Non-limiting examples of GPCR activities include transmembrane signaling of various forms, which may involve G protein association and/or the exertion of an influence over G protein binding of various guanidylate nucleotides; another exemplary activity of GPCRs is the binding of accessory proteins or polypeptides that differ from known G proteins.

The modulators of the invention exhibit a variety of chemical structures, which can be generally grouped into non-peptide mimetics of natural GPCR receptor ligands, peptide and non-peptide allosteric effectors of GPCR receptors, and peptides that may function as activators or inhibitors (competitive, uncompetitive and non-competitive) (*e.g.*, antibody products) of GPCR receptors. The invention does not restrict the sources for suitable modulators, which may be obtained from natural sources such as plant, animal or mineral extracts, or non-natural sources such as small molecule libraries, including the products of combinatorial chemical approaches to library construction, and peptide libraries. Examples of peptide modulators of GPCR receptors exhibit the following primary structures: GLGPRPLRFamide, GNSFLRFamide, GGPQGPLRFamide, GPSGPLRFamide, PDVDHVFLRFamide, and pyro-EDVDHVFLRFamide.

Other assays can be used to examine enzymatic activity including, but not limited to, photometric, radiometric, HPLC, electrochemical, and the like, which are described in, for example, *Enzyme Assays: A Practical Approach*, eds. R. Eisinger and M. J. Danson, 1992, Oxford University Press, which is incorporated herein by reference in its entirety.

The use of cDNAs encoding GPCRs in drug discovery programs is well-known; assays capable of testing thousands of unknown compounds per day in high-throughput screens (HTSs) are thoroughly documented. The literature is replete with examples of the use of radiolabelled ligands in HTS binding assays for drug discovery (see Williams, *Medicinal Research Reviews*, 1991, 11, 147-184.; Sweetnam, *et al.*, *J. Natural Products*, 1993, 56, 441-455 for review).

Recombinant receptors are preferred for binding assay HTS because they allow for better specificity (higher relative purity), provide the ability to generate large amounts of receptor material, and can be used in a broad variety of formats (see Hodgson, *Bio/Technology*, 1992, 10, 973-980; each of which is incorporated herein by reference in its entirety).

5 A variety of heterologous systems is available for functional expression of recombinant receptors that are well known to those skilled in the art. Such systems include bacteria (Strosberg, *et al.*, *Trends in Pharmacological Sciences*, 1992, 13, 95-98), yeast (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494), several kinds of insect cells (Vanden Broeck, *Int. Rev. Cytology*, 1996, 164, 189-268), amphibian cells (Jayawickreme *et al.*, *Current Opinion in Biotechnology*, 1997, 8, 629-634) and several mammalian cell lines (CHO, HEK293, COS, etc.; see Gerhardt, *et al.*, *Eur. J. Pharmacology*, 1997, 334, 1-23). These examples do not preclude the use of other possible cell expression systems, including cell lines obtained from nematodes (PCT application WO 98/37177).

15 In preferred embodiments of the invention, methods of screening for compounds that modulate nGPCR-x activity comprise contacting test compounds with nGPCR-x and assaying for the presence of a complex between the compound and nGPCR-x. In such assays, the ligand is typically labeled. After suitable incubation, free ligand is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular compound to bind to nGPCR-x.

20 It is well known that activation of heterologous receptors expressed in recombinant systems results in a variety of biological responses, which are mediated by G proteins expressed in the host cells. Occupation of a GPCR by an agonist results in exchange of bound GDP for GTP at a binding site on the  $G_\alpha$  subunit; one can use a radioactive, non-hydrolyzable derivative of GTP,  $GTP\gamma[^{35}S]$ , to measure binding of an agonist to the receptor (Sim *et al.*, *Neuroreport*, 1996, 7, 729-733). One can also use this binding to measure the ability of antagonists to bind to the receptor by decreasing binding of  $GTP\gamma[^{35}S]$  in the presence of a known agonist. One could therefore construct a HTS based on  $GTP\gamma[^{35}S]$  binding, though this is not the preferred method.

The G proteins required for functional expression of heterologous GPCRs can be native constituents of the host cell or can be introduced through well-known recombinant technology. 30 The G proteins can be intact or chimeric. Often, a nearly universally competent G protein (e.g.,  $G_{\alpha 16}$ ) is used to couple any given receptor to a detectable response pathway. G protein activation results in the stimulation or inhibition of other native proteins, events that can be linked to a measurable response.

Examples of such biological responses include, but are not limited to, the following: the ability to survive in the absence of a limiting nutrient in specifically engineered yeast cells (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494); changes in intracellular  $\text{Ca}^{2+}$  concentration as measured by fluorescent dyes (Murphy, *et al.*, *Cur. Opinion Drug Disc. Dev.*, 1998, 1, 192-199). Fluorescence changes can also be used to monitor ligand-induced changes in membrane potential or intracellular pH; an automated system suitable for HTS has been described for these purposes (Schroeder, *et al.*, *J. Biomolecular Screening*, 1996, 1, 75-80). Melanophores prepared from *Xenopus laevis* show a ligand-dependent change in pigment organization in response to heterologous GPCR activation; this response is adaptable to HTS formats (Jayawickreme *et al.*, *Cur. Opinion Biotechnology*, 1997, 8, 629-634). Assays are also available for the measurement of common second messengers, including cAMP, phosphoinositides and arachidonic acid, but these are not generally preferred for HTS.

Preferred methods of HTS employing these receptors include permanently transfected CHO cells, in which agonists and antagonists can be identified by the ability to specifically alter the binding of  $\text{GTP}\gamma[^{35}\text{S}]$  in membranes prepared from these cells. In another embodiment of the invention, permanently transfected CHO cells could be used for the preparation of membranes which contain significant amounts of the recombinant receptor proteins; these membrane preparations would then be used in receptor binding assays, employing the radiolabelled ligand specific for the particular receptor. Alternatively, a functional assay, such as fluorescent monitoring of ligand-induced changes in internal  $\text{Ca}^{2+}$  concentration or membrane potential in permanently transfected CHO cells containing each of these receptors individually or in combination would be preferred for HTS. Equally preferred would be an alternative type of mammalian cell, such as HEK293 or COS cells, in similar formats. More preferred would be permanently transfected insect cell lines, such as *Drosophila* S2 cells. Even more preferred would be recombinant yeast cells expressing the *Drosophila melanogaster* receptors in HTS formats well known to those skilled in the art (*e.g.*, Pausch, *Trends in Biotechnology*, 1997, 15, 487-494).

The invention contemplates a multitude of assays to screen and identify inhibitors of ligand binding to nGPCR-x receptors. In one example, the nGPCR-x receptor is immobilized and interaction with a binding partner is assessed in the presence and absence of a candidate modulator such as an inhibitor compound. In another example, interaction between the nGPCR-x receptor and its binding partner is assessed in a solution assay, both in the presence and absence of a candidate inhibitor compound. In either assay, an inhibitor is identified as a compound that decreases binding between the nGPCR-x receptor and its binding partner.

Following identification of compounds that inhibit ligand binding to nGPCR-x receptors, such compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Another contemplated assay involves a variation of the dihybrid assay wherein an inhibitor of protein/protein interactions is identified by detection  
5 of a positive signal in a transformed or transfected host cell, as described in PCT publication number WO 95/20652, published August 3, 1995.

Candidate modulators contemplated by the invention include compounds selected from libraries of either potential activators or potential inhibitors. There are a number of different libraries used for the identification of small molecule modulators, including: (1) chemical  
10 libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules. Chemical libraries consist of random chemical structures, some of which are analogs of known compounds or analogs of compounds that have been identified as "hits" or "leads" in other drug discovery screens, some of which are derived from natural products, and some of which arise from non-directed synthetic organic chemistry.  
15 Natural product libraries are collections of microorganisms, animals, plants, or marine organisms which are used to create mixtures for screening by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of plants or marine organisms. Natural product libraries include polyketides, non-ribosomal peptides, and variants (non-naturally occurring) thereof. For a review, see *Science* 282:63-68 (1998). Combinatorial  
20 libraries are composed of large numbers of peptides, oligonucleotides, or organic compounds as a mixture. These libraries are relatively easy to prepare by traditional automated synthesis methods, PCR, cloning, or proprietary synthetic methods. Of particular interest are non-peptide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of  
25 combinatorial chemistry and libraries created therefrom, see Myers, *Curr. Opin. Biotechnol.* 8:701-707 (1997). Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to modulate activity.

Still other candidate inhibitors contemplated by the invention can be designed and  
30 include soluble forms of binding partners, as well as such binding partners as chimeric, or fusion, proteins. A "binding partner" as used herein broadly encompasses non-peptide modulators; as well as such peptide modulators as neuropeptides other than natural ligands, antibodies, antibody fragments, and modified compounds comprising antibody domains that are immunospecific for the expression product of the identified nGPCR-x gene.

The polypeptides of the invention are employed as a research tool for identification, characterization and purification of interacting, regulatory proteins. Appropriate labels are incorporated into the polypeptides of the invention by various methods known in the art and the polypeptides are used to capture interacting molecules. For example, molecules are incubated with the labeled polypeptides, washed to remove unbound polypeptides, and the polypeptide complex is quantified. Data obtained using different concentrations of polypeptide are used to calculate values for the number, affinity, and association of polypeptide with the protein complex.

Labeled polypeptides are also useful as reagents for the purification of molecules with which the polypeptide interacts including, but not limited to, inhibitors. In one embodiment of affinity purification, a polypeptide is covalently coupled to a chromatography column. Cells and their membranes are extracted, and various cellular subcomponents are passed over the column. Molecules bind to the column by virtue of their affinity to the polypeptide. The polypeptide-complex is recovered from the column, dissociated and the recovered molecule is subjected to protein sequencing. This amino acid sequence is then used to identify the captured molecule or to design degenerate oligonucleotides for cloning the corresponding gene from an appropriate cDNA library.

Alternatively, compounds may be identified which exhibit similar properties to the ligand for the nGPCR-x of the invention, but which are smaller and exhibit a longer half time than the endogenous ligand in a human or animal body. When an organic compound is designed, a molecule according to the invention is used as a "lead" compound. The design of mimetics to known pharmaceutically active compounds is a well-known approach in the development of pharmaceuticals based on such "lead" compounds. Mimetic design, synthesis and testing are generally used to avoid randomly screening a large number of molecules for a target property. Furthermore, structural data deriving from the analysis of the deduced amino acid sequences encoded by the DNAs of the present invention are useful to design new drugs, more specific and therefore with a higher pharmacological potency.

Comparison of the protein sequence of the present invention with the sequences present in all the available databases showed a significant homology with the transmembrane portion of G protein coupled receptors. Accordingly, computer modeling can be used to develop a putative tertiary structure of the proteins of the invention based on the available information of the transmembrane domain of other proteins. Thus, novel ligands based on the predicted structure of nGPCR-x can be designed.

In a particular embodiment, the novel molecules identified by the screening methods according to the invention are low molecular weight organic molecules, in which case a composition or pharmaceutical composition can be prepared thereof for oral intake, such as in tablets. The compositions, or pharmaceutical compositions, comprising the nucleic acid  
5 molecules, vectors, polypeptides, antibodies and compounds identified by the screening methods described herein, can be prepared for any route of administration including, but not limited to, oral, intravenous, cutaneous, subcutaneous, nasal, intramuscular or intraperitoneal. The nature of the carrier or other ingredients will depend on the specific route of administration and particular embodiment of the invention to be administered. Examples of techniques and protocols that are  
10 useful in this context are, *inter alia*, found in Remington's Pharmaceutical Sciences, 16<sup>th</sup> edition, Osol, A (ed.), 1980, which is incorporated herein by reference in its entirety.

The dosage of these low molecular weight compounds will depend on the disease state or condition to be treated and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound. For treating human or animals,  
15 between approximately 0.5 mg/kg of body weight to 500 mg/kg of body weight of the compound can be administered. Therapy is typically administered at lower dosages and is continued until the desired therapeutic outcome is observed.

The present compounds and methods, including nucleic acid molecules, polypeptides, antibodies, compounds identified by the screening methods described herein, have a variety of  
20 pharmaceutical applications and may be used, for example, to treat or prevent unregulated cellular growth, such as cancer cell and tumor growth. In a particular embodiment, the present molecules are used in gene therapy. For a review of gene therapy procedures, see *e.g.* Anderson, *Science*, 1992, 256, 808-813, which is incorporated herein by reference in its entirety.

The present invention also encompasses a method of agonizing (stimulating) or  
25 antagonizing a nGPCR-x natural binding partner associated activity in a mammal comprising administering to said mammal an agonist or antagonist to one of the above disclosed polypeptides in an amount sufficient to effect said agonism or antagonism. One embodiment of the present invention, then, is a method of treating diseases in a mammal with an agonist or antagonist of the protein of the present invention comprises administering the agonist or  
30 antagonist to a mammal in an amount sufficient to agonize or antagonize nGPCR-x-associated functions.

In an effort to discover novel treatments for diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that modulate the function of G



protein coupled receptors. Some small organic molecules form a class of compounds that modulate the function of G protein coupled receptors.

Exemplary diseases and conditions amenable to treatment based on the present invention include, but are not limited to, thyroid disorders (*e.g.* thyreotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, Schizophrenia, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

Methods of determining the dosages of compounds to be administered to a patient and modes of administering compounds to an organism are disclosed in U.S. Application Serial No. 08/702,282, filed August 23, 1996 and International patent publication number WO 96/22976, published August 1 1996, both of which are incorporated herein by reference in their entirety, including any drawings, figures or tables. Those skilled in the art will appreciate that such descriptions are applicable to the present invention and can be easily adapted to it.

The proper dosage depends on various factors such as the type of disease being treated, the particular composition being used and the size and physiological condition of the patient, including such factors as, for example, weight, age, sex, disease state, *etc.* Therapeutically effective doses for the compounds described herein can be estimated initially from cell culture and animal models. For example, a dose can be formulated in animal models to achieve a circulating concentration range that initially takes into account the IC<sub>50</sub> as determined in cell culture assays. The animal model data can be used to more accurately determine useful doses in humans.

Plasma half-life and biodistribution of the drug and metabolites in the plasma, tumors and major organs can also be determined to facilitate the selection of drugs most appropriate to inhibit a disorder. Such measurements can be carried out. For example, HPLC analysis can be performed on the plasma of animals treated with the drug and the location of radiolabeled compounds can be determined using detection methods such as X-ray, CAT scan and MRI. Compounds that show potent inhibitory activity in the screening assays, but have poor pharmacokinetic characteristics, can be optimized by altering the chemical structure and retesting. In this regard, compounds displaying good pharmacokinetic characteristics can be used as a model.

Toxicity studies can also be carried out by measuring the blood cell composition. For example, toxicity studies can be carried out in a suitable animal model as follows: 1) the compound is administered to mice (an untreated control mouse should also be used); 2) blood samples are periodically obtained via the tail vein from one mouse in each treatment group; and 3) the samples are analyzed for red and white blood cell counts, blood cell composition and the percent of lymphocytes versus polymorphonuclear cells. A comparison of results for each dosing regime with the controls indicates if toxicity is present.

At the termination of each toxicity study, further studies can be carried out by sacrificing the animals (preferably, in accordance with the American Veterinary Medical Association guidelines Report of the American Veterinary Medical Assoc. Panel on Euthanasia, Journal of American Veterinary Medical Assoc., 202:229-249, 1993). Representative animals from each treatment group can then be examined by gross necropsy for immediate evidence of metastasis, unusual illness or toxicity. Gross abnormalities in tissue are noted and tissues are examined histologically. Compounds causing a reduction in body weight or blood components are less preferred, as are compounds having an adverse effect on major organs. In general, the greater the adverse effect the less preferred the compound.

For the treatment of many diseases, the expected daily dose of a hydrophobic pharmaceutical agent is between 1 to 500 mg/day, preferably 1 to 250 mg/day, and most preferably 1 to 50 mg/day. Drugs can be delivered less frequently provided plasma levels of the active moiety are sufficient to maintain therapeutic effectiveness. Plasma levels should reflect the potency of the drug. Generally, the more potent the compound the lower the plasma levels necessary to achieve efficacy.

nGPCR-42, 46, 48, 49, 51, 52, 61, 63, and 70 mRNA transcripts may found in many tissues, including, but not limited to, brain, peripheral blood lymphocytes, pancreas, ovary, uterus, testis, salivary gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues. Within the brain,

nGPCR-42, 46, 48, 49, 51, 52, 61, 63, and 70 mRNA transcripts may be found in many tissues, including, but not limited to, frontal lobe, hypothalamus, pons, cerebellum, caudate nucleus, and medulla. Tissues and brain regions where specific nGPCR mRNA transcripts are expressed are identified in the Examples, below.

5 Sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60 will, as detailed above, enable screening the endogenous neurotransmitters/hormones/ligands which activate, agonize, or antagonize nGPCR-x and for compounds with potential utility in treating disorders including, but not limited to, thyroid disorders (*e.g.* thyreotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases related to cell  
10 differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, Schizophrenia, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's  
15 Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies,  
20 atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

For example, nGPCR-x may be useful in the treatment of respiratory ailments such as  
25 asthma, where T cells are implicated by the disease. Contraction of airway smooth muscle is stimulated by thrombin. Cicala *et al* (1999) Br J Pharmacol 126:478-484. Additionally, in bronchiolitis obliterans, it has been noted that activation of thrombin receptors may be deleterious. Hauck *et al.* (1999) Am J Physiol 277:L22-L29. Furthermore, mast cells have also been shown to have thrombin receptors. Cirino *et al* (1996) J Exp Med 183:821-827. nGPCR-x  
30 may also be useful in remodeling of airway structures in chronic pulmonary inflammation via stimulation of fibroblast procollagen synthesis. See, *e.g.*, Chambers *et al.* (1998) Biochem J 333:121-127; Trejo *et al.* (1996) J Biol Chem 271:21536-21541.

In another example, increased release of sCD40L and expression of CD40L by T cells after activation of thrombin receptors suggests that nGPCR-x may be useful in the treatment of

unstable angina due to the role of T cells and inflammation. See Aukrust *et al.* (1999) Circulation 100:614-620.

A further example is the treatment of inflammatory diseases, such as psoriasis, inflammatory bowel disease, multiple sclerosis, rheumatoid arthritis, and thyroiditis. Due to the tissue expression profile of nGPCR-x, inhibition of thrombin receptors may be beneficial for these diseases. See, *e.g.*, Morris *et al.* (1996) Ann Rheum Dis 55:841-843. In addition to T cells, NK cells and monocytes are also critical cell types which contribute to the pathogenesis of these diseases. See, *e.g.*, Naldini & Carney (1996) Cell Immunol 172:35-42; Hoffman & Cooper (1995) Blood Cells Mol Dis 21:156-167; Colotta *et al.* (1994) Am J Pathol 144:975-985.

Expression of nGPCR-x in bone marrow and spleen may suggest that it may play a role in the proliferation of hematopoietic progenitor cells. See DiCuccio *et al.* (1996) Exp Hematol 24:914-918.

As another example, nGPCR-x may be useful in the treatment of acute and/or traumatic brain injury. Astrocytes have been demonstrated to express thrombin receptors. Activation of thrombin receptors may be involved in astrogliosis following brain injury. Therefore, inhibition of receptor activity may be beneficial for limiting neuroinflammation. Scar formation mediated by astrocytes may also be limited by inhibiting thrombin receptors. See, *e.g.*, Pindon *et al.* (1998) Eur J Biochem 255:766-774; Ubl & Reiser. (1997) Glia 21:361-369; Grabham & Cunningham (1995) J Neurochem 64:583-591.

nGPCR-x receptor activation may mediate neuronal and astrocyte apoptosis and prevention of neurite outgrowth. Inhibition would be beneficial in both chronic and acute brain injury. See, *e.g.*, Donovan *et al.* (1997) J Neurosci 17:5316-5326; Turgeon *et al.* (1998) J Neurosci 18:6882-6891; Smith-Swintosky *et al.* (1997) J Neurochem 69:1890-1896; Gill *et al.* (1998) Brain Res 797:321-327; Suidan *et al.* (1996) Semin Thromb Hemost 22:125-133.

The attached Sequence Listing contains the sequences of the polynucleotides and polypeptides of the invention and is incorporated herein by reference in its entirety.

As described above and in Example 5 below, the genes encoding nGPCR-42, 46, 48, 49, 51, 52, 61, 63, and 70 have been detected in brain tissue indicating that these n-GPCR-x proteins are neuroreceptors. The identification of modulators such as agonists and antagonists is therefore useful for the identification of compounds useful to treat neurological diseases and psychiatric disorders. Such neurological diseases and disorders, including but are not limited to, mental disorder, affective disorders, ADHD/ADD, and neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, schizophrenia, and senile dementia as well as depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like.

## Methods of Screening Human Subjects

Thus in yet another embodiment, the invention provides genetic screening procedures that entail analyzing a person's genome – in particular their alleles for GPCRs of the invention - to determine whether the individual possesses a genetic characteristic found in other individuals that are considered to be afflicted with, or at risk for, developing a mental disorder or disease of the brain that is suspected of having a hereditary component. For example, in one embodiment, the invention provides a method for determining a potential for developing a disorder affecting the brain in a human subject comprising the steps of analyzing the coding sequence of one or more GPCR genes from the human subject; and determining development potential for the disorder in said human subject from the analyzing step.

More particularly, the invention provides a method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of: (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain, wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120, or an allelic variant thereof, and wherein the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of allele in the nucleic acid correlates with an increased risk of developing the disorder. In preferred variations, the seven transmembrane receptor is nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 comprising an amino acid sequence set forth in SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120, or an allelic variant thereof, and the disease is mental disorder.

By "human subject" is meant any human being, human embryo, or human fetus. It will be apparent that methods of the present invention will be of particular interest to individuals that have themselves been diagnosed with a disorder affecting the brain or have relatives that have been diagnosed with a disorder affecting the brain.

By "screening for an increased risk" is meant determination of whether a genetic variation exists in the human subject that correlates with a greater likelihood of developing a disorder affecting the brain than exists for the human population as a whole, or for a relevant racial or ethnic human sub-population to which the individual belongs. Both positive and negative determinations (i.e., determinations that a genetic predisposition marker is present or is absent) are intended to fall within the scope of screening methods of the invention. In preferred

embodiments, the presence of a mutation altering the sequence or expression of at least one nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 seven transmembrane receptor allele in the nucleic acid is correlated with an increased risk of developing mental disorder, whereas the absence of such a mutation is reported as a negative determination.

5       The “assaying” step of the invention may involve any techniques available for analyzing nucleic acid to determine its characteristics, including but not limited to well-known techniques such as single-strand conformation polymorphism analysis (SSCP) (Orita *et al.*, *Proc Natl. Acad. Sci. USA*, 86: 2766-2770 (1989)); heteroduplex analysis (White *et al.*, *Genomics*, 12: 301-306 (1992)); denaturing gradient gel electrophoresis analysis (Fischer *et al.*, *Proc. Natl. Acad. Sci. USA*, 80: 1579-1583 (1983); and Riesner *et al.*, *Electrophoresis*, 10: 377-389 (1989)); DNA sequencing; RNase cleavage (Myers *et al.*, *Science*, 230: 1242-1246 (1985)); chemical cleavage of mismatch techniques (Rowley *et al.*, *Genomics*, 30: 574-582 (1995); and Roberts *et al.*, *Nucl. Acids Res.*, 25: 3377-3378 (1997)); restriction fragment length polymorphism analysis; single nucleotide primer extension analysis (Shumaker *et al.*, *Hum. Mutat.*, 7: 346-354 (1996); and 10       Pastinen *et al.*, *Genome Res.*, 7: 606-614 (1997)); 5’ nuclease assays (Pease *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:5022-5026 (1994)); DNA Microchip analysis (Ramsay, G., *Nature Biotechnology*, 16: 40-48 (1999); and Chee *et al.*, U.S. Patent No. 5,837,832); and ligase chain reaction (Whiteley *et al.*, U.S. Patent No. 5,521,065). (See generally, Schafer and Hawkins, *Nature Biotechnology*, 16: 33-39 (1998).) All of the foregoing documents are hereby 15       incorporated herein by reference in their entirety. 20

Thus, in one preferred embodiment involving screening nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 sequences, for example, the assaying step comprises at least one procedure selected from the group consisting of: (a) determining a nucleotide sequence of at least one codon of at least one nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 allele of the human subject; (b) 25       performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; (c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and (d) performing a restriction endonuclease digestion to determine whether nucleic 30       acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

In a highly preferred embodiment, the assaying involves sequencing of nucleic acid to determine nucleotide sequence thereof, using any available sequencing technique. (See, *e.g.*, Sanger *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 74: 5463-5467 (1977) (dideoxy chain termination

method); Mirzabekov, *TIBTECH*, 12: 27-32 (1994) (sequencing by hybridization); Drmanac *et al.*, *Nature Biotechnology*, 16: 54-58 (1998); U.S. Patent No. 5,202,231; and *Science*, 260: 1649-1652 (1993) (sequencing by hybridization); Kieleczawa *et al.*, *Science*, 258: 1787-1791 (1992) (sequencing by primer walking); (Douglas *et al.*, *Biotechniques*, 14: 824-828 (1993) (Direct sequencing of PCR products); and Akane *et al.*, *Biotechniques* 16: 238-241 (1994); Maxam and Gilbert, *Meth. Enzymol.*, 65: 499-560 (1977) (chemical termination sequencing), all incorporated herein by reference in its entirety). The analysis may entail sequencing of the entire nGPCR gene genomic DNA sequence, or portions thereof; or sequencing of the entire seven transmembrane receptor coding sequence or portions thereof. In some circumstances, the analysis may involve a determination of whether an individual possesses a particular allelic variant, in which case sequencing of only a small portion of nucleic acid -- enough to determine the sequence of a particular codon characterizing the allelic variant -- is sufficient. This approach is appropriate, for example, when assaying to determine whether one family member inherited the same allelic variant that has been previously characterized for another family member, or, more generally, whether a person's genome contains an allelic variant that has been previously characterized and correlated with a mental disorder having a heritable component.

In another highly preferred embodiment, the assaying step comprises performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences. In a preferred embodiment, the hybridization involves a determination of whether nucleic acid derived from the human subject will hybridize with one or more oligonucleotides, wherein the oligonucleotides have nucleotide sequences that correspond identically to a portion of the GPCR gene sequence taught herein, such as the nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 coding sequence set forth in SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120, or that correspond identically except for one mismatch. The hybridization conditions are selected to differentiate between perfect sequence complementarity and imperfect matches differing by one or more bases. Such hybridization experiments thereby can provide single nucleotide polymorphism sequence information about the nucleic acid from the human subject, by virtue of knowing the sequences of the oligonucleotides used in the experiments.

Several of the techniques outlined above involve an analysis wherein one performs a polynucleotide migration assay, *e.g.*, on a polyacrylamide electrophoresis gel (or in a capillary electrophoresis system), under denaturing or non-denaturing conditions. Nucleic acid derived from the human subject is subjected to gel electrophoresis, usually adjacent to (or co-loaded with) one or more reference nucleic acids, such as reference GPCR-encoding sequences having

a coding sequence identical to all or a portion of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 (or identical except for one known polymorphism). The nucleic acid from the human subject and the reference sequence(s) are subjected to similar chemical or enzymatic treatments and then electrophoresed under conditions whereby the polynucleotides will show a differential migration pattern, unless they contain identical sequences. (See generally Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, New York: John Wiley & Sons, Inc. (1987-1999); and Sambrook *et al.*, (eds.), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989), both incorporated herein by reference in their entirety).

10 In the context of assaying, the term "nucleic acid of a human subject" is intended to include nucleic acid obtained directly from the human subject (*e.g.*, DNA or RNA obtained from a biological sample such as a blood, tissue, or other cell or fluid sample); and also nucleic acid derived from nucleic acid obtained directly from the human subject. By way of non-limiting examples, well known procedures exist for creating cDNA that is complementary to RNA  
15 derived from a biological sample from a human subject, and for amplifying (*e.g.*, via polymerase chain reaction (PCR)) DNA or RNA derived from a biological sample obtained from a human subject. Any such derived polynucleotide which retains relevant nucleotide sequence information of the human subject's own DNA/RNA is intended to fall within the definition of "nucleic acid of a human subject" for the purposes of the present invention.

20 In the context of assaying, the term "mutation" includes addition, deletion, and/or substitution of one or more nucleotides in the GPCR gene sequence (*e.g.*, as compared to the seven transmembrane receptor-encoding sequences set forth of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 and other polymorphisms that occur in introns (where introns exist) and that are identifiable via sequencing, restriction fragment length polymorphism, or  
25 other techniques. The various activity examples provided herein permit determination of whether a mutation modulates activity of the relevant receptor in the presence or absence of various test substances.

In a related embodiment, the invention provides methods of screening a person's genotype with respect to GPCRs of the invention, and correlating such genotypes with  
30 diagnoses for disease or with predisposition for disease (for genetic counseling). For example, the invention provides a method of screening for an nGPCR-63 hereditary mental disorder genotype in a human patient, comprising the steps of: (a) providing a biological sample comprising nucleic acid from the patient, the nucleic acid including sequences corresponding to said patient's nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 alleles; (b) analyzing the nucleic acid



for the presence of a mutation or mutations; (c) determining an nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 genotype from the analyzing step; and (d) correlating the presence of a mutation in an nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 allele with a hereditary mental disorder genotype. In a preferred embodiment, the biological sample is a cell sample containing human  
5 cells that contain genomic DNA of the human subject. The analyzing can be performed analogously to the assaying described in preceding paragraphs. For example, the analyzing comprises sequencing a portion of the nucleic acid (e.g., DNA or RNA), the portion comprising at least one codon of the nGPCR-63 alleles.

Although more time consuming and expensive than methods involving nucleic acid  
10 analysis, the invention also may be practiced by assaying protein of a human subject to determine the presence or absence of an amino acid sequence variation in GPCR protein from the human subject. Such protein analyses may be performed, e.g., by fragmenting GPCR protein via chemical or enzymatic methods and sequencing the resultant peptides; or by Western analyses using an antibody having specificity for a particular allelic variant of the GPCR.

15 The invention also provides materials that are useful for performing methods of the invention. For example, the present invention provides oligonucleotides useful as probes in the many analyzing techniques described above. In general, such oligonucleotide probes comprise 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides that have a  
20 sequence that is identical, or exactly complementary, to a portion of a human GPCR gene sequence taught herein (or allelic variant thereof), or that is identical or exactly complementary except for one nucleotide substitution. In a preferred embodiment, the oligonucleotides have a sequence that corresponds in the foregoing manner to a human GPCR coding sequence taught herein, and in particular, the coding sequences set forth in SEQ ID Numbers 61, 62, 68, 91, 94,  
25 96, 97, 99, 100, and 111-120. In one variation, an oligonucleotide probe of the invention is purified and isolated. In another variation, the oligonucleotide probe is labeled, e.g., with a radioisotope, chromophore, or fluorophore. In yet another variation, the probe is covalently attached to a solid support. (See generally Ausubel *et al.* and Sambrook *et al.*, *supra*.)

In a related embodiment, the invention provides kits comprising reagents that are useful  
30 for practicing methods of the invention. For example, the invention provides a kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association: (a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 seven transmembrane receptor gene, the oligonucleotide comprising 6-50 nucleotides that have a sequence that is identical or exactly

complementary to a portion of a human nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 gene sequence or nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and (b) a media packaged with the oligonucleotide

5 containing information identifying polymorphisms identifiable with the probe that correlate with mental disorder or a genetic predisposition therefor. Exemplary information-containing media include printed paper package inserts or packaging labels; and magnetic and optical storage media that are readable by computers or machines used by practitioners who perform genetic screening and counseling services. The practitioner uses the information provided in the media  
10 to correlate the results of the analysis with the oligonucleotide with a diagnosis. In a preferred variation, the oligonucleotide is labeled.

In still another embodiment, the invention provides methods of identifying those allelic variants of GPCRs of the invention that correlate with mental disorders. For example, the invention provides a method of identifying a seven transmembrane allelic variant that correlates  
15 with a mental disorder, comprising steps of: (a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny; (b) analyzing the nucleic acid for the presence of a mutation or mutations in at least one seven transmembrane receptor that is expressed in the brain, wherein the at least one seven transmembrane receptor comprises an amino acid sequence selected from  
20 the group consisting of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding the at least one seven transmembrane receptor; (c) determining a genotype for the patient for the at least one seven transmembrane receptor from said analyzing step; and (d) identifying an allelic variant that correlates with the mental disorder from the determining step.  
25 To expedite this process, it may be desirable to perform linkage studies in the patients (and possibly their families) to correlate chromosomal markers with disease states. The chromosomal localization data provided herein facilitates identifying an involved GPCR with a chromosomal marker.

The foregoing method can be performed to correlate GPCRs of the invention to a  
30 number of disorders having hereditary components that are causative or that predispose persons to the disorder. For example, in one preferred variation, the disorder is a mental disorder, and the at least one seven transmembrane receptor comprises nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 having an amino acid sequence set forth in SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 or an allelic variant thereof.

Also contemplated as part of the invention are polynucleotides that comprise the allelic variant sequences identified by such methods, and polypeptides encoded by the allelic variant sequences, and oligonucleotide and oligopeptide fragments thereof that embody the mutations that have been identified. Such materials are useful in *in vitro* cell-free and cell-based assays for identifying lead compounds and therapeutics for treatment of the disorders. For example, the variants are used in activity assays, binding assays, and assays to screen for activity modulators described herein. In one preferred embodiment, the invention provides a purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 receptor allelic variant identified according to the methods described above; and an oligonucleotide that comprises the sequences that differentiate the allelic variant from the nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 sequences set forth in SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120. The invention also provides a vector comprising the polynucleotide (preferably an expression vector); and a host cell transformed or transfected with the polynucleotide or vector. The invention also provides an isolated cell line that is expressing the allelic variant GPCR polypeptide; purified cell membranes from such cells; purified polypeptide; and synthetic peptides that embody the allelic variation amino acid sequence. In one particular embodiment, the invention provides a purified polynucleotide comprising a nucleotide sequence encoding a nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 seven transmembrane receptor protein of a human that is affected with a mental disorder; wherein said polynucleotide hybridizes to the complement of SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 under the following hybridization conditions: (a) hybridization for 16 hours at 42C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60C in a wash solution comprising 0.1x SSC and 1% SDS; and wherein the polynucleotide encodes a nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 amino acid sequence that differs from SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120 by at least one residue.

An exemplary assay for using the allelic variants is a method for identifying a modulator of nGPCR-x biological activity, comprising the steps of: (a) contacting a cell expressing the allelic variant in the presence and in the absence of a putative modulator compound; (b) measuring nGPCR-x biological activity in the cell; and (c) identifying a putative modulator compound in view of decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator.

Additional features of the invention will be apparent from the following Examples. Examples 1-6, 12, 14, and 15 are actual, while the remaining Examples are prophetic.

Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, including the detailed description, and all such features are intended as aspects of the invention. Likewise, features of the invention described herein can be re-combined into additional embodiments that also are intended as aspects of the invention, irrespective of whether the combination of features is specifically mentioned above as an aspect or embodiment of the invention. Also, only such limitations which are described herein as critical to the invention should be viewed as such; variations of the invention lacking limitations which have not been described herein as critical are intended as aspects of the invention.

## EXAMPLES

### Example 1: Identification of nGPCR-x

#### A. Database search

The Celera database was searched using known GPCR receptors as query sequences to find patterns suggestive of novel G protein-coupled receptors. Positive hits were further analyzed with the GCG program BLAST to determine which ones were the most likely candidates to encode G protein-coupled receptors, using the standard (default) alignment produced by BLAST as a guide.

Briefly, the BLAST algorithm, which stands for Basic Local Alignment Search Tool is suitable for determining sequence similarity (Altschul *et al.*, J. Mol. Biol., 1990, 215, 403-410, which is incorporated herein by reference in its entirety). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information through the world wide web of the Internet (ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension for the word hits in each direction are halted when: 1) the cumulative alignment score falls off by the quantity X from its maximum achieved value; 2) the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or 3) the end of either sequence is reached. The Blast algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The Blast program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff *et al.*, Proc. Natl. Acad. Sci. USA, 1992,

89, 10915-10919, which is incorporated herein by reference in its entirety) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm (Karlin *et al.*, Proc. Natl. Acad. Sci. USA, 1993, 90, 5873-5787, which is incorporated herein by reference in its entirety) and Gapped BLAST perform a statistical analysis of the similarity between two sequences. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a GPCR gene or cDNA if the smallest sum probability in comparison of the test nucleic acid to a GPCR nucleic acid is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Homology searches are performed with the program BLAST version 2.08. A collection of about 200 to about 350 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

Multiple query sequences may have a significant alignment to the same genomic region, although each alignment may not cover exactly the same DNA region. A procedure is used to determine the region of maximum common overlap between the alignments from several query sequences. This region is called the consensus DNA region. The procedure for determining this consensus involves the automatic parsing of the BLAST output files using the program MSPcrunch to produce a tabular report. From this tabular report the start and end of each alignment in the genomic DNA is extracted. This information is used by a PERL script to derive the maximum common overlap. These regions are reported in the form of a unique sequence identifier, a start and the end position in the sequence. The sequences defined by these regions were extracted from the original genomic sequence file using the program fetchdb.

The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCRs. These sequences were then submitted for further sequence analysis.

Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs.

*nGPCR-70*

Homology searches were performed with the program BLAST version 2.08. A collection of 286 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.1 were collected from each BLAST search. The amino acid sequences were edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program "GelStart". After assembly with GelStart, a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-70. Further sequence analysis involved the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Transmembrane regions within the amino acid sequences were visually inspected with an outliner (MaxThink) and the Celera sequences with promising transmembrane regions were kept. In a last step, the blast results were compared against SWPLUS and the transmembrane regions of the sequences and promising Celera sequences for full length cloning were selected. The sequence of nGPCR-70s is shown above in Table 5.

#### *nGPCR-63*

Homology searches were performed with the program BLAST version 2.08. A collection of 286 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.1 were collected from each BLAST search. The amino acid sequences were edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program "GelStart". After assembly with "GelStart" a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-63. Further sequence analysis involved the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Transmembrane regions within the amino acid sequences were visually inspected with an outliner (MaxThink) and the Celera sequences with promising transmembrane regions were kept. In a last step, the blast results were compared against SWPLUS and the

transmembrane regions of the sequences and promising Celera sequences for full length cloning were selected. The sequence of nGPCR-63 is shown above in Table 5.

#### *nGPCR-42*

Homology searches were performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-42. These sequences were then submitted for further sequence analysis. Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Only sequences that contained transmembrane regions in a pattern found in GPCRs were retained. These nGPCR-42s are shown above in Table 5.

#### *nGPCR-46*

Homology searches are performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-46. These sequences were then submitted for further sequence analysis. Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Only sequences that contained transmembrane regions in a pattern found in GPCRs were retained. These nGPCR-46s are shown above in Table 5.

#### *nGPCR-48*

Homology searches are performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-48. These sequences were then submitted for further sequence analysis. Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmtrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Only sequences that contained transmembrane regions in a pattern found in GPCRs were retained. These nGPCR-48s are shown above in Table 5.

#### nGPCR-49

Homology searches were performed with the program BLAST version 2.08. A collection of 286 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.1 were collected from each BLAST search. The amino acid sequences were edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program "GelStart". After assembly with GelStart a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-49. Further sequence analysis involved the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmtrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Transmembrane regions within the amino acid sequences were visually inspected with an outliner (MaxThink) and the Celera sequences with promising transmembrane regions were kept. In a last step, the blast results were compared against SWPLUS and the transmembrane regions of the sequences and promising Celera sequences for full length cloning were selected. The sequence of nGPCR-49 is shown above in Table 5.

#### nGPCR-61



Homology searches were performed with the program BLAST version 2.08. A collection of 286 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.1 were collected from each BLAST search. The amino acid sequences were edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-61. Further sequence analysis involved the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Transmembrane regions within the amino acid sequences were visually inspected with an outliner (MaxThink) and the Celera sequences with promising transmembrane regions were kept. In a last step, the blast results were compared against SWPLUS and the transmembrane regions of the sequences and promising Celera sequences for full length cloning were selected. The sequence of nGPCR-61 is shown above in Table 5.

#### *nGPCR-51*

Homology searches were performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences were edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions were assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-51. These sequences were then submitted for further sequence analysis. Further sequence analysis involved the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Only sequences that contained transmembrane regions in a pattern found in GPCRs were retained. The sequence of nGPCR-51 is shown above in Table 5.

#### *nGPCR-52*

Homology searches are performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the  
5 sequence that produce non-significant alignments with proteins that are not related to GPCRs.

The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCR-52. These sequences were then submitted for further sequence analysis. Further sequence analysis involves the removal of sequences previously isolated and  
10 removal of sequences that are related to olfactory GPCRs. The transmembrane regions for the sequences that remained were determined using a FORTRAN computer program called "tmtrest.all" (Parodi *et al.*, Comput. Appl. Biosci. 5:527-535(1994)). Only sequences that contained transmembrane regions in a pattern found in GPCRs were retained. These nGPCR-52s are shown above in Table 5.

15 nGPCR-x cDNAs were sequenced directly using an ABI377 fluorescence-based sequencer (Perkin-Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM™ Ready Dye-Deoxy Terminator kit with Taq FS™ polymerase. Each ABI cycle sequencing reaction contained about 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98C for 1 minute, followed by 50 cycles using the following  
20 parameters: 98C for 30 seconds, annealing at 50C for 30 seconds, and extension at 60C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using Centriflex™ gel filtration cartridges (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B  
25 tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 5 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90C for three minutes and loaded into the gel sample wells for sequence analysis using the ABI377 sequencer. Sequence analysis was performed by importing  
30 ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp were obtained. Potential sequencing errors were minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers annealing at different locations until all sequencing ambiguities were removed.

The following Table 5 contains the sequences of the polynucleotides and polypeptides of the invention. Start and stop codons within the polynucleotide sequence are identified by boldface type. The transmembrane domains within the polypeptide sequence are identified by underlining.

5

Table 5

The following DNA sequence nGPCR-Seq61 <SEQ ID NO. 1> was identified in *H. sapiens*:

CCTATGCTACAATCGGCAGATGGGAACTAGGAGCCATGATCTCTCAGATTGCAGGTCTCATTG  
GAACCACATTTATTGGATTTTCCTTTTATAGTAGTACTAACATCATACTACTCTTTTGTAAAGCCA  
TCTGAGAAAAATAAGAACCTGTACGTCCATTATGGAGAAAGATTGACTTACAGTTCTGTGAA  
AAGACATCTTTTGGTCATCCAGATTCTACTAATAGTTTGCTTCCTTCCTTATAGTATTTTAAAC  
CCATTTTTTATGTTCTACACCAAAGAGATAACTGTCTAGCAATTGAATTATTTAATAGAAACAA  
AAAACATTCTCACCTGTCTTGCTTCGGCCAGAAGTAGCACAGACCCCATTAATTTCTTTTATT  
AGACAAAACATTCAAGAAGACACTATATAATCTCTTTACAAAGTCTAATTCAGCACATATGCA  
ATCATATGGTTGACTTTTGAATGGAAAACCCACAATATTAAGAAAAGCATTTCATGTGACTTT  
ATTAGGGACACTAAACTACATCATTAACATGTCACAG

The following amino acid sequence <SEQ ID NO. 61> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 1:

YATIGRWE LGAMISQIAG LIGTTFIGFS FLVVLTSYYS FVSHLRKIRT CTSIMEKDLT  
YSSVKRHLV IQILLVCF LYSIFKPIFY VLHQRDNCQQ LNYLIETKNI LTCLASARSS  
TDPHFLLD KTFKKTLYNL FTKSNSAHMQ SYG\*LLNGKP HNIKKSIHVT LLGTLNYIIN MSQ

The following DNA sequence nGPCR-Seq63 <SEQ ID NO. 2> was identified in *H. sapiens*:

CAGTGAGCCG AGATGGTGCC ATTGCACTCT AGCCTGGGGC AACAGAGCCG ACTCCATCT  
CCAAAAA AAAA AAGGCCATTC TGAGGATCAA GGCACCACTA GCAACAGGGA GCCCATGGG  
TCTCAGACCC TCTCCCCACA TCTCCTGGTCCCTGCCCCCA CCTGGCGTAC AGGGACCAGC  
CCCACGGAAG GCTCTTGAGG CCAGGTAACC ATGGGGAGGG GAGGAATGGG GACACCTTCC  
TCCTGAGTGT CTTAGGGAAG AGAAGCTTAG GTCAGGTGGC TGAGGGTGGG AATGAGAGAG  
GGGTCTCCTC CTGGAGGGTC TCACCATTC CTTGGTCACC CACCCAACTC TCATCTCCCC  
TGATGTGGGG AGGAGCAGGG GGCATGGATT CCTGAGCCCC AGACTCAACT GTTGTGGTTT  
ACAGGGGCAT CAGGAGAGAG AGCGAGCAGA ACACACTCCT GCAGCATCCC CTGGCCCCC  
GCCCATGAT GGAGCCCAGA GAAGCTGGAC AGCACGTGGG GGCCGCCAAC GGCGCCAGG  
AGGATGTGGC CTTCAACCTC ATCATCCTGT CCCTCACC GA GGGGCTCGGC CTCGGTGGGC  
TGCTGGGGAA TGGGGCAGTC CTCTGGCTGC TCAGCTCCAA TGTCTACAGA AACCCTTCG  
CCATCTACCT CCTGGACGTG GCCTGCGCGG ATCTCATCTT CTTGGCTGC CACATGGTGG  
CCATCGTCCC CGACTTGCTG CAAGGCCGGC TGGACTTCCC GGGCTTCGTG CAGACCAGCC  
TGGCAACGCT GCGCTTCTC TGCTACATCG TGGGCCTGAG TCTCCTGGCG GCCGTCAGCG  
TGGAGCAGTG CCTGGCCGCC CTCTTCCAG CCTGGTACTC GTGCCGCCG CCACGCCACC  
TGACCACCTG TGTGTGCGCC CTCACCTGGG CCTCTGCCT GCTGCTGCAC CTGCTGCTCA  
GCGGCGCCTG CACCCAGTTC TTCGGGGAGC CCAGCCGCCA CTTGTGCCG ACGCTGTGGC  
TGGTGGCAGC GGTGCTGCTG GCTCTGCTGT GTTGACCAT GTGTGGGGCC AGCCTTATGC  
TGCTGTGCG GGTGGAGCGA GGCCCCCAGC GGCCCCCACC CCGGGGCTT CCTGGGCTCA  
TCTCTCTCAC CGTCTCCTC TCTCTTCT GCGGCTGCC CTTCCGCATC TACTGGCTGT  
CCCGGAACCT GCTCTGGTAC ATCCCCCACT ACTTCTACCA CTTACGCTC CTCATGGCCG  
CCGTGCACTG CGCGGCCAAG CCCGTCGTCT ACTTCTGCCT GGGCAGTGCC CAGGGCCGCA  
GGCTGCCCT CCGGCTGGTC CTCCAGCGAG CGCTGGGAGA CGAGGCTGAG CTGGGGGCCG  
TCAGGGAGAC CTCCCGCCGG GGCCTGGTGG ACATAGCAGC CTGAGCCCTG GGGCCCCCGA  
CCCCAGCTGC AGCCCCCGTG AGGCAAGAGG GTGACGTGGG GAAGGTGGTG GGGTCAGAGG  
CTGGGGCCAG CCGGACCTGG AGGAGGCCTT GGTGGGTGAC CCGGTCATGT GCTGTCAAAG  
TTGTGACCCT TGGTCTGGAG CATGAGGCTC CCCTGGGAGG CAGCTGGAAA GG

The following amino acid sequence <SEQ ID NO. 62> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 2:

VSRDGA IAL\*PGATEP DSISKKKRPF \*GSRHH\*QQG APWVSDPLPT SPGPCPHLAY  
 RDQPHGRLLR PGNHGEGRNG DTFLLSVLGK RSLGQVAEGG NERGVSSWRV SPFPWSPTQL  
 SSPLMWGGAG GMD\*APDST VVVYRGIRRE SEQNTLLQHP LAPRPMMEPR EAGQHVGAAN  
 GAQEDVAFNL ILSLTEGLG LGLLGNGAV LWLLSSNVYR NPFAYLLDV ACADLIFLGC  
 HMVAIVPDL QGRLDFFPGFV QTSLATLRF CYIVGLSLLA AVSVEQCLAA LFPAYWSCRR  
 PRHLTTCVCA LTWALCLLLH LLLSGACTQF FGEPsrHLCR TLWLVA AVL ALLCCTMCGA  
 SLMLLLRVER GPQRPPRGF PGLILLTVLL FLFCGLPFI YWLSRNLLWY IPHYFYHFSF  
 LMAAVHCAAK PVVYFCLGSA QGRRPLRLV LQRALGDEAE LGAVRETSRR GLVDIAA\*AL  
 GPPTPAAAPV RQEGDVGKVV GSEAGASRTW RRPWWVTRSC AVKVVTGLGLE HEAPLGGSWK

The following DNA sequence nGPCR-Seq64 <SEQ ID NO. 3> was identified in *H. sapiens*:

AAAATTGCTC TTCCTCCTGA GCTTGACAC AATGATTGAG TTCAAGATGA AGAAGATGGA  
 GCAGGGCACC AGGTAGACGG TGAAGCAGTG GATCCAGATG AGGACGTGAT GCACAGAGGT  
 GCTGATGTAG TCTTCAGTCC AGATGTTGGG CCACCAGTAA TAGGGGATGC TGGTCAGGAA  
 GCAGGTGATG TAAACACTTA CAATGACTTT CCGGGTGC GG GCTGGGTATG AGACCGTGTG  
 GTACTTGAGC GGGTGGCAGA CAGCGATATA CCTGTCAATG GTTAACGGTA CAGTAATCCA  
 TATGGAGGTG TGGATGGATG AGAATTCCAG CACTTCTATG ATCTTGTCGG GGACCTGAGG  
 CATCTGCATG TTCAAGATGA AATCTTCCAA CAGGAAGTCC ACAAACACTA TGAAAAAGAG  
 GACCAAGATG TCGGCAGCAG CGAGTGCCAA GAGATAGTTG TAGGAGGACT TCTGTCTTCT  
 TGCCACCAGC TGGGAGAGGA TGATCACTGT CAAGATATTT GCTGTGGAGA GAAGAAAAAC  
 TGGTTTAGCT CTGAAGCAAA GATGACTTCG TTGGCTCCTA TGGGGGCCCT AGGCATATGT  
 TTATTTTGCA CTCCCATGGA AGTGAAAATG ATTGAATCAA TGCTTTTGAG GGACAACCTC  
 AGCATTACAA ATAGCACCTC ATACAATTAG TGGATACTAT TTTAAAGTTA TGCTTATATT  
 CTAACACAAC CATGAGAGGT GGTGCCTCCA TTCTCCTCAT CTTAGAAGTG AAACCTGGGGC  
 TCTGAGAGCC TCACACAGCC ATGAGAGGTG GTGCCGCCAT TCTCCTCACA CAACCATGAG  
 TGGTGTGCC GCCATTCTCC TCACACAACA ATGAGAGGTG GTGCCACCAT TCTTCTCACA  
 CAACCGTGAG AGGCGATGCT GCCATTATCC

The following amino acid sequence <SEQ ID NO. 63> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 3:

G\* WQHRLSRLCE KNGGTTSHCC VRRMAATPLM VV\*GEWRHHL SWLCEALRAP  
 VSLLR\*GEWR HHLSWLC\*NI SITLK\*YPLI V\*GAICNAEV VPQKH\*FNHF HFHGSAK\*TY  
 A\*GPHRSQRS HLCFRAKPVF LLSTANILTV ILSQLVARR QKSSYNYLLA LAAADILVLF  
 FIVFVDFLE DFILNMQMPQ VPDKIIEVLE FSSIHTSIWI TVPLTIDRYI AVCHPLKYHT  
 VSYPARTRKV IVSVYITCFL TSIPYYWWPN IWTEYISTS VHHVLIWIHC FTVYLVPCSI  
 FFILNSIIVY KLRRKSNFDN GSIASHGCVR

The following DNA sequence nGPCR-Seq65 <SEQ ID NO. 4> was identified in *H. sapiens*:

GGCCCGCTCCCACGCTGTGTAGTGTACTTTTCATTTTCAATAAATCACTTCATTCTTCTCT  
 TGCTTTGTTTGTGCGTTTTGTCCAATTCITTTGTTCAAGACGCCAAGAACCTGGACAGCCT  
 CCACCATTAAAGAATACAAGAGCAGTTTCTGTACATGTACATATGGGGGTGGGTGGATCT  
 CGCTCAGCCTTTCCAGGACACAGCGGTTTGATGGAATGCTTTTCTGAAACTCGTGGCAGA  
 ATGAGTACGGAAGAGGGCTGATAGCCGAATTCAGCCAGTGGAGCCAGAAGCGTGTCTCAT  
 AGAGAAAGTGCTGGACATGTGACCCGTGGCAGGCTCCGATGATCAGAGCAGTGTAG  
 GGAGCCCAGCACAGGACAAATGCACACCATGACGTCAAGCGACGTGGACGCTCATTTG  
 TCCCGGGAGACGCCGTGCTGGGGAGGCTCGGCTGTCTGCTGCCATTGCAGGCGTTCTCGC  
 TGCCCTGTAGCGGGTCTTCTGCGTAGGATTATCAAGTCCACCTGCAGAGGTGA

The following amino acid sequence <SEQ ID NO. 64> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 4:

HLCRWT\*\*SY AEDPLQSEN ACNGSRQPSL PSTASPGTNE RPRRLTSWC HLSCAGLPTL  
 LVIIGAACHG SHVQHFLYET RFWLHWLNSA ISPLPYSFCH EFQKSIPSNR CVLERLSEIH  
 PPPYVHVTET ALVFLMVEAV QVLGVLNKL DKTHKQSKG MK\*FIENEST LHSVAG

The following DNA sequence nGPCR-Seq66 <SEQ ID NO. 5> was identified in *H. sapiens*:

TACTTTTACCCAAAACATTATGTTTCATTATTAGAGTTTCCTAGAAAAATACCTAAGGAG  
 TTAATGGTTTCATATCTTTTGCTCTATTATGAAGAAGAAAAATGTTTTACAAATATTTTC  
 ATTATTGGAGCATTTTTTGTGTTAGTGAAATTATCAAAACTAGGATTGATTCTATTCT  
 GTTTACTTTTGTATAATCTTTATCCTTTTCTCTTAATTTCTGTATTTTGGATGCCTAAC  
 TTAGAATACATTACCAAAGTTACCTTTTCATTTAGTCTCTCAATACAAGATGATTTAAAA  
 CATTTATGGTTACCTTTTTTAATTTTTTGTCTATGCAAATTTATAAAAGGGCAAAGTCTT  
 TGTGCTCTAATAATACCTGCTTTCTCATGTTTTACATGTTCTACGATTTATTTTGTTTTT  
 ATAATGTAATTTTCGTTTACCTAATTGTGCACATAGTGAATAATAGATTATAATGAAGAA  
 AACTTGGATTAAAAATCTATTGTTAAAAAGGTTTTTCAGGCAATAATAAATCATTGGATTT  
 TTCTGATGTATTTTAAAAAGATATGTTTATTTTGTAGCAACTCGTGTGC

The following amino acid sequence <SEQ ID NO. 65> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 5:

YFYPKLLCSL LEFPRKIPKE LMVSYLLLYY EEEKCFNIF IIGAFFVVSE IIKTRIDFYS  
 VYFCYNLYPF LLISVFWMPN LEYITKVTF SLSIQDDLK HLWLPFLIFL LCKFIKGQSL  
 CALIPAFSC FTCSTIYFVF IM\*FSFT\*LC T\*\*IIDYNEE NLD\*NLLKR FFRQ\*\*IIG F\*CILKRYVY  
 F\*ATRV

The following DNA sequence nGPCR-Seq67 <SEQ ID NO. 6> was identified in *H. sapiens*:

GGCTACTTGT TATGGAAAAG TTATGATTGG TGTAGAAGTAC TACAGGTTGC TTCTGGCACT  
 TGAAGAAAGC CTTGAGCGTG GCGGCTGCCA TGGTGCGGAA CCGCTTGCTG ATGAAGCAGT  
 AGAGGAAGAA GTTGATGGCT GTGTTGAGAA GGGCTAGCAT GTTGGCAATG TCGGACATGA  
 TGTGTACCAG CCAGCGGTTT TGGATGGGCG CCCCATAGAG GTGGTAAAGA ATCATGATGA  
 TCGCGGGGGG CCAAAGTGTG GCAAAGATGG AGGTAATGGT GAACAAGATG GCGGTGGTCT  
 TCCCCGTGGA GTAGCCACGG AGACGAAAAT TGCTCTTCCT CCTGAGCTTG TACACAATGA  
 TTGAGTTCAA GATGAAGAAC ATGGAGCAGG GCACCAAGTA GACGGTGAAG CAGTGGATCC  
 AGATGAGGAC GTGATGCACA GAGGTGCTGA TGTAATCTTC AGTCCAGATG TTGGGCCACC  
 AGTAATAGGG GATGCTGGTC AGGAAGCAGG TGATGTAAAC ACTTACAATG ACTTTCCGGG  
 TCGCGGCTGG GTATGAGACC GTGTGGTACT TGAGCGGGTG GCAGACAGCG ATATACCTGT  
 CAATGGTTAA CGGTACAGTA ATCCATATGG AGGTGTGGAT GGATGAGAAT TCCAGCACTT  
 CTATGATCTT GTCGGGGACC TGAGGCATCT GCATGTTCAA GATGAAATCT TCCAACAGGA  
 AGTCCACAAA CACTATGAAA AAGAGGACCA AGATGTCGGC AGCAGCGAGT GCCAAGAGAT  
 AGTTGTAGGA GGACTTCTGT CTTCTTGCCA CCAGCTGGGA GAGGATGATC ACTGTCAAGA  
 TATTTGCTGT GGAGAGAAGA AAAC

The following amino acid sequence <SEQ ID NO. 66> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 6:

FLL STANILTVII LSQLVARRQK SSYNYLLALA AADILVLFVI VFVDFLLEDF ILNMQMPQVP  
DKIIEVLEFS SIHTSIWITV PLTIDRYIAV CHPLKYHTVS YPARTRKVIV SVYITCFILTS  
IPYYWWPNIW TEDYISTSVH HVLWIHCFT VYLVPSCMFF ILNSIIYVKL RRKSNFRLRG  
YSTGKTTAIL FTITSIFATL WAPRIMILY HLYGAPIQNR WL VHIMSDIA NMLALLNTAI  
NFFLYCFISK RFRTMAAATL KAFFKCQKQP VQFYTNHNFS ITSS

The following DNA sequence nGPCR-Seq68 <SEQ ID NO. 7> was identified in *H. sapiens*:

GACACCTGCC AACATGTTCA TTATCAACCT CGCGGTCAGC GACTTCCTCA TGTCCTTCAC  
 CCAGGCCCTT GTCTTCTTCA CCAGTAGCCT CTATAAGCAG TGGCTCTTTG GGGAGACAGG  
 TAGATGCTGG GGCTCCCTTT TGCTGGAGGG AGGAGGAGGG TTTTGACCTG GGGATGCCCT  
 CAATGGAGGG TGGCCCAAAG GAGGTGATTT GCTGCTTCTG GGCAGAGAGT GGGTAGCTGC  
 CCTCAGTCCTGTGAGTAAGC AAGAAGGGAA GATGCAGTGT TGGTCCTAAG GCCTCTGCCA  
 GCCTTGGCCTA GATGTGGCAG GTGGAGGGGG TGGAGTGCGC TCAGTCCTGC TCTTCTGTG  
 AGGTGAAGGC CAGAGCAGAG TCTACCCTGT CCCAGACCC TCCTCCCCAG GACTCAGAGC  
 AGGGGCTGTG CCCACAGGCT GCGAGTTCTA TGCCTTCTGT GGAGCTCTCT TTGGCATTTT  
 CTCCATGATC ACCCTGACGG CCATCGCCCT GGACCGCTAC CTGGTAATCA CACGCCCGCT  
 GGCCACCTTT GGTGTGGCGT CCAAGAGGCG TCGGCATTT GTCTGCTGG GCGTTTGGCT  
 CTATGCCCTG GCCTGGAGTC TGCCACCCTT CTTCGGCTGG AGTAAGTGGG CTGCTGGAAC

TGGAAGGGGG GCAGATGGGC TGGGAGGGGC ACATTCAAGG GTAAGTAGGT GACTTGGGT  
CAGCCAGCTG GCGGGAGCAG GGTGCCAGG AGCTACCTGA GCCTCAGGTG AGATGGACAT  
TCAGGGGGAC ATGACTGGCA GCAAGGGAAA CTGACACTGC CCCA

The following amino acid sequence <SEQ ID NO. 67> is a predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 7:

TPANMF IINLAVSDFL MSFTQAPVFF TSSLYKQWLF GETGRCWGS LLEGGGGF\*P  
GDALNGGWPK GGDLLLLGRE WVAALSPVSK QEGKMQCWS\* GLCQPWPDVA GGGGGVRSVL  
LFL\*GEGQSR VYPVPRPSSP GLRAGAVPTG CEFYAFCGAL FGISSMITLT AIALDRYLVI  
TRPLATFGVA SKRRAAFVLL GVWLYALAWS LPPFFGWSKW AAGTGRGADG LGGAHSRVS  
R WTWVSQLAGA GCPGAT\*ASG EMDIQGDMTG SKGN\*HCPHL

The following DNA sequence nGPCR-Seq70 <SEQ ID NO. 8> was identified in *H. sapiens*:

TTCTCCCCTTGACGGGTGACTAACTCTGCCTGCGTGTTTCTTTTGTACCAGCATAGGCA  
CTGAGTGCGGTCTGTGCACCCCTTTGCCACCCACCGGTGCCGGCACTGAGCCTGCAACCT  
GTCTACGCCCTCTGGCTGTTGCCATGACGTCCACCTGCACCAACAGCACGCGGAGAGT  
AACAGCAGCCACACGTGCATGCCCTCTCCAAAATGCCCATCAGCCTGGCCCACGGCATC  
ATCCGCTCAACCGTGCTGGTTATCTTCCTCGCCGCTCTTTCTGTCGGCAACATAGTGCTG  
GCGCTAGTGTTGCAGCGCAAGCCGCGAGCTGCTGCAGGTGACCAACCGTTTATCTTTAAC  
CTCCTCGTCACCGACCTGCTGCAGATTTCGCTCGTGGCCCCCTGGGTGGTGGCCACCTCT  
GTGCCTCTCTTCTGGCCCCCTCAACAGCCACTTCTGCACGGCCCTGGTTAGCCTCACCCAC  
CTGTTTCGCTTCGCCAGCGTCAACACCATTTGCTGTTGGTGTGAGTGGATCGCTACTTGTCC  
ATCATCCACCTCTCTTCTACCCGTCCAAGATGACCCAGCGCCGCGGGTACCTGCTCCTC  
TATGGCACCTGGATTGTGG

The following amino acid sequence <SEQ ID NO. 68> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 8:

FSP\*RVTNSA CVFLLSPA\*A LSAVCAPLCH PPVPALSLQP VSRPLAVAMT STCTNSTRES  
NSSHTCMPLS KMPISLAHGI IRSTVLVIFL AASFVGNIVL ALVLQRPQL LQVTNRFIEN  
LLVTDLLQIS LVAPWVWATS VPLFWPLNSH FCTALVSLTH LFAFASVNTI VVVSVDRLYS  
IHLPLFYPSK MTQRRGYLLL YGTWIV

The following DNA sequence nGPCR-Seq71 <SEQ ID NO. 9> was identified in *H. sapiens*:

AGGACAACGATGGTCACTGATTTGGTGACCTTCGACAGTCTCCGGGCGCTGGCTCCGGTC  
GGGCGTCTCCGGCTACCGCGGCCCTCCTTTGGTCCCCGCGCGCGCGGTCCGGCGATG  
AAGCGCACCAGCAGCAGGTAGCACAAGATAATGATGCCAGCGGCAGCACGAAGCCCAGC  
AGCACCTTCTGCGAGTTGGTAGAGGCCAGCCAGAACTGCCTGTCGCGGCCAGCAACTTG  
TCCGGGAAACGCACCAGGCACAGCTCCTCGCCCATCACCTTGACCGTGGTGGAGAAAATG  
GCACTGGGCAGCGAGGCCAGCGCGGCCAAAGCCCAGATCCACACACACAGCGCCTTGGCC  
GAGAAGCAGCAGCTGTCCCCAGGCTCCGGCCGCGAGCAGTCGCCCCGGCGGTGTCTCGG  
GTCCCGGTGGCTCTTCAGAGCCGAGGCCACCGAATGGTAGCGCGTCACTCATGGCAGT  
GAGGAAGAACACGCTGGCGTACATGTTTATGGGACGTCACCATGGACACGATCTTACACA  
TGGCCTTGCC

The following amino acid sequence <SEQ ID NO. 69> is a predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 9:

QGHV\*D RVHGDVP\*TC TPACSSSLP\* V\*RATIRWPR L\*RATGTRGH GRGDCCGRSL  
GDSCCFSKA LCVWTWALAA LASLPSAIFS TTVKVMGEEL CLVRFPDKLL GRDRQFWLGL  
YHSQKVLLGF VLPLGIILC YLLLVRFIAD RRAAGTKGGA AVAGGRPTGA SARRLSKVTK  
SVTIV

The following DNA sequence nGPCR-Seq69 <SEQ ID NO. 10> was identified in *H. sapiens*:

CTGGAAAGGG TGCTGGTGGC GCGCCGCGAC GCCGCGGCGC GACTGCCTGC CTGGTACTCA  
ACCTCTTCTG CGCGGACCTG CTCTTCATCA GCGCTATCCC TCTGGTGCTG GCCGTGCGCT  
GGACTGAGGC CTGGCTGCTG GGCCCCGTTG CCGCCACCT GCTCTTCTAC GTGATGACCC

TGAGCGGCAG CGTCACCATC CTCACGCTGG CCGCGGTCAG CCTGGAGCGC ATGGTGTGCA  
TCGTGCACCT GCAGCGCGGC GTGCGGGGTC CTGGGCGGCG GGC GCGGGCA GTGCTGCTGG  
CGCTCATCTG GGGCTATTCG GCGGTCGCC CTCTGCCTCT CTGCGTCTT TCCGAGTCG  
TCCCGCAACG GCTCCCCGGC GCCGACCAGG TGAGCGCCCC TCTGTGTGTG CCGGGCAGGT  
GTCCTGCGCA GGCTGGGAAG CGGGGCCCCG ACGGAAGCTG GGATGAGGAT GATCAAGAAC  
AACAATAGCC ATTTATTGCA CTTAATCGTT GTGCCAAATC TTGTGCCCAT GGCTGTGAAG  
TTTAATCTCT TAAATCTCAC TACAACGCTG TGCACACGCC CTCCTAAATG ATGTAAGTGG  
AGTCCCCCAA ATTCTTGCAA AATGCAATGA CTGTTGCGAG GTTAATTAAC GAGTAGTTTA  
GGAGCGAGAC GGAACTTTGG GGGTGCAGGG TGGCCAAACA CTTTGTATTG AATCATGATT  
CCTCGCCAGG TGCTACAATA CTGTTATTAT CACACCCATT TCACAGATGA GAACCAGAGG  
CACACCGAAG TGTATAATAA CTTGCCCGA GTATTTTATC CGTAATTCGA GGAGGAGATG  
GGCTCCTTCC AGAAGTTTAC CCGTAATTCA AGGAGGAGTT GGGCTCCTGT CCAGGGTTGG  
GTTATGGTCC TGCTTTGAAA GCGCGCGGAC AGGCATGTGA GACCCGGGGA CCCAGATGC  
AATGCTGTCT TTAGGGGACT TGTGACAGAA TTCCCTTCCG GGGTCTTCAG TTTTTCAGC  
TGCAAAACGG AAGGATTACA CTAGACCTC GAGGTGTCCT GGGCGCCTGA AATGTGCAGA  
TTACAGAGGC TGGACCGACG AGCT

The following amino acid sequence <SEQ ID NO. 70> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 10:

KKGAG GAPRRRGATA CLVLNLF CAD LLFISAIP LV LAVRWTEAWL LGPVACHLLF  
YVMTLSGSVT ILTLAAVSLE RMVCIVHLQR GVRGPGRRAR AVLLALI WGY SAVAALPLCV  
FFRVVPQRLP GADQVSAPLC VPGRCPAQAG KRPGDGSWDE DDQEQQ\*PFI ALNRCAKSCA  
HGCEV\*SLKS HYNAVHTPS\* MM\*VESPKFL QNAMTVARLI NE\*FRSETEL WGCRAVAKHFV  
LNHDSSPGAT ILLSHPFHR \*EPEAHRV\* \*LAQSILSVI RGGDGLLEPV YP\*FKEELGS  
CPGLGYGPAL KARGQACETR GPQMCCCL\*G TCDRIPFRGL QFFQLQNGRI TDLRGLVGA  
\*NVQITEAGP TS

The following DNA sequence nGPCR-Seq2011 <SEQ ID NO. 11> was identified in *H. sapiens*:

TTAATCCCTGGAAGTCCACGAACAATGAATCCATTTTCATGCATCTTGTGGAACACCTCT  
GCCGAACTTTTAAACAAATCCTGGAATAAAGAGTTTGCTTATCAAACAGGGCTGGTTGGCAAC  
GATACAGTCATCCTCCCTTCCATGATTGGGATTATCTGTTCAACAGGGCTGGTTGGCAAC  
ATCCTCATTTGATTCACTATAATAAGGTAAGGAATGGCTCCTTTTTTTTTTTTTCCTTC  
CATACTTTAGGAACTACAGTCAAAGCTCCCTAAATGAGTCCTTTCCCTGTAGCATTTT  
GCTTAATGAAATGCAATTTTGAAAATATTTGCTTAAGATAATTAATGAAGATTCTACAGA  
TATTTTCGTCATGCATTAGGTAACATCTCAGTTGCAAAATCTCAACATGCTAAGACCTAGG  
CCAATGCTTACTGCTGGGTCAGTGAGTTTTTAGGGAATGACTCTCACTCTCAGTCTTAG  
CTGCATATTAGAATCATCTGGGGAGCTTTAAAAACTCCTGATATGCAGTTTACCCCAGA  
CCCATTA ACTCAGAATCTCTAAGGGTAGGGCCCGGTAAGATTAAAACTG

The following amino acid sequence <SEQ ID NO. 71> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 11:

LIPGSPRTMN PFHASCWNTS AELLNKSWNK EFAYQTASVV DTVILPSMIG IICSTGLVGN  
ILIVFTIR\* GMAPFFFFSF HTLGNYSQSS LNESFPL\*HF A\*\*NAILEIF A\*DN\*\*RFYR  
YFRHALGNIS VANLNMLRPR PMLTAGSVSF \*GNDSHSQS\* LHIRI WGAL KTPDMQFHPR  
PINSESLRVG PG\*DLKL

The following DNA sequence nGPCR-Seq2012 <SEQ ID NO. 12> was identified in *H. sapiens*:

GGGTGACAGGA ACTCTGGAAGGTGATAGTTTTCCAGGTGAAGAAGGGACAGTAGGTTTCC  
CTGCTTTACTGCTATCTCCCTATGGCCTTCGGTCAGGCCTGAACTGTGATGAGAGTCTCT  
GCCCCGTGCTCTTTGGGATTCTCAAATATCCTCCCTTCATTGTTCCCTGAATTATCTAGA  
TCAATACAAACCAATCCCTCCTCTGCTTAAACTTTTAATTGCTTACAGGGGAAAGTGCA  
GACTCCTTAGCTGGCGCACAAGCCTTTTGACCTGGCACACATAGAGCTTTCTAGGGCTA  
TCTCCTGCCATTCTCTAGCCTGTCTCCTTCTGCTCTGGGTCTCCAGAACTGCCCATGAAT  
TCCCTCACATGCCTTTCT

The following amino acid sequence <SEQ ID NO. 72> is the predicted amino acid sequence derived from

the DNA sequence of SEQ ID NO. 12:

VTGTLEGDS FPGEEGTVGF PALLSPYGL RSGLNCDL CPCSLGFLKY PPFIVP\*II\*  
INTNQSLCL KLLIAYRGKC RLLSWRTSL HLAHIELSRA ISCHSLACL LLWVSRTAHE  
FPHMPF

The following DNA sequence nGPCR-Seq2013 <SEQ ID NO. 13> was identified in *H. sapiens*:

CCTATGCTTTCTATAGCTCATGGACCAAGCCATTTCTCCAACATCATAATTTTCTCTTA  
TCATCCCTTTCTTGCAATTATCTCCAGATGAAAAAGTTCAAATGCCTCCTCCCTATTT  
TCAGCTGAGAAGGTGAGATGCCCGTGCTGGATTTCCAGGCTCCTGCCTGCTTGGCAGAC  
CACTGTGACCGTGACGACTAAGTCCACTCTTCCCAGCCAGCATCCCCTGCTCTCCCCAG  
GCCAACTTCTTTGTGTTCTATTCTATCTCCTTGCCGGAAGTACTCAGACCCCTCT  
GTCTTTCTGCCTCTTGCCCAATCTTCCAGGCTCTAGTTTGTGCTGGCTTTCTGCCTCCAAAA  
ATGATTTTAAGCATTTGTGAGTCTTTCTCTCCACAGAACTCCAGACTCTTTGAAAATGCC  
GTAGTTCGTAAATTACTTCCAATATTTAAGAAGTGCTTATCATGCTTCCCTGCGGGTTCA  
TGCTCTTTGGTATTGATTTCT

The following amino acid sequence <SEQ ID NO. 73> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 13:

YAFLLMDQAISPTS\*FFLIIPFLALFSR\*KKFKCLLPIFS\*EGEMPVPGFPGSCLLGRPL\*PCTT  
KSTLPSQHPLLSPGQLLCVLFIPISELLRPLCLASCPIFQALVCWLSASKNDFKHL\*  
VFLSTELQTL\*KCRSS\*ITSNI\*EVLIMLP CGFMLFGIDF

The following DNA sequence nGPCR-Seq2014 <SEQ ID NO. 14> was identified in *H. sapiens*:

CAAATGAGGTGGATGCACTTAGAAGGAAGGGTAGAACAAATATTCGTTGATGATATACAA  
TGTGTCAGATACAGAGATACAGACTAGGGCTGTGGGAATCTCAGATTTTAATTTGTCA  
ACAGTGTGTTTTGATTTTTTTGTTTTGTATATTTGCCTCCCCAATTAATTTACAGGCTT  
AATCATCTTTACAAGACATTATTTTAAAGAGAAAGCGAACTTACTGAAGTTTTATGCTTC  
CCTGATTGTGATGAGCTGGTTGATTCTAGCTCTAGTTTCAATGTTCTGGAAAATACTGAA  
GTACTTCCACCTGGTGGCACTTAGTGAACATTGCAGAACCGAGTAATAGGTTATTCCGTT  
GGGTTTCTCGAACAAATCTGAGTTATAGCTAGGAAGTCTCAACTAACAATTTATGAGAAC  
CTTCTGCTACACATGTGAATTTACATATTTATTCCTTTGCAGTTGAAGGATGGGATGTAT  
GCAAAGGAGAATGAAGTCTACTACAGTAATGGAAGAAGTGATAGAAGATGAAACTCCAA  
AATGCCTATCTGCTACTTGAGTGGAAGTCACTAGAATATATCAAATGCAGAGAGAAACAT  
GTGACTAGGCTCTAGTCAAACAATTGTCATCAAATACTTGCTGAATATATAACACATTTT  
AGGGAGCTGTG

The following amino acid sequence <SEQ ID NO. 74> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 14:

K\*GGCT\*KEG\*NKYSMLIYNVSDTEIHLGLWESQILICQQCVLIFLCIFASPIFNLGLIIFTRHY  
FKEKANLLKFYASLIVMSWLILALVSMFWKILKYFHLVALSEHCRTE\*\*V IPLGFSNKSE  
L\*LGTLN\*QFMRFTCYTCEF TYLFLCS\*RM GCMQRRMNST TVMERSDRR\*NSKMPICYLS  
GTH\*NISNAERNM\*LGSSQTIVIKYLLNI\*HILGSC

The following DNA sequence nGPCR-Seq2015 <SEQ ID NO. 15> was identified in *H. sapiens*:

AAAAAATACATTCCTGTATGGAAGACTAAATATTAAGAAAAAACAAGTTATCCATATG  
TTGTTTCATAGCTTTAAGTATCCTAGTCTAAAATCCAAAGGGACTTTTTTGGGGACTCAA  
AAAAAATTTCTAAAGTTGATCCAGAAGAGTAAAGAGACTAATATAGTCAAAATAATAT  
TTGAAAAGGTTAAATAATTATGGGAAAACATGCTTAGCAAAAATATAAACAACAAAAC  
TCAAACAGACAAAACCCCTATGAAAAACCAAAACCTATTTAAAAACTAAATTCTAAATG  
AGAACTAACATACAAATCAAGAATCAGCATAGATAGCTAAGAAGCAAATTATAGCATATA  
AACTAATTCAATTTATTATAAAG

The following amino acid sequence <SEQ ID NO. 75> is the predicted amino acid sequence derived from



the DNA sequence of SEQ ID NO. 15:

FI IN\*ISLYAII CFLAIYADS\* FVC\*FSFRI\* FLNRFLVFHR VFVCLKFCLF IFLLSMFSHN  
YLTFNSNILT ILVSLLFWIN FRIFFFESPK KSLWILD\*DT \*SYETTYG\*L FFSLIFSLPY RNVFF

The following DNA sequence nGPCR-Seq2016 <SEQ ID NO. 16> was identified in *H. sapiens*:

CTAGGCTGCTCCTGTGTGGTTATAATGAATCTCATCAGCTACAATCATTTCCCAAAAAGG  
ACAGAAAGTGAATCTGTCTTATAAAGATGGTTTTCAAAGCCAACTACAGATATCATAGAA  
AAAAAAAAAAAAACAGTGGCAGATAAGACCTTTCATCTTTTCTTATCCCATGGGCTTTT  
CTGCCTCCAGATTTCTTGCACATCAAGGGGGTCTGTGCAAATCAGTGGGCCTGGCCCT  
CTGCCCACCTGGGCTAAGCTGTGGTTGCCATGGCTACGAAGACAAGAATGACCCTTGGT  
TTTATAGAGCCCTGGGTCCTTGCTCACAGGCCTTTTCAGTTGATATTTCTTTCATCTTC  
TTCAGAACCCCATGTGGCAGGTGCAACAAGGAGTTCACCAATGCTCCGGCCAAAGTGAGT  
AATGAGGGACACATGCACGACCAGGCAGAGCAGCCCTGAAGAAGTGGTTTGTGAGGGTGT  
GGAGGAAGGGCTTGCTTCCACACTCTGCACTTCTGGGTCCTAAGGCACTAATCACACCTG  
GCTCAGGCATCATTCTCAGCCTCATGCTACTTTTGTCCAGGGACAGAACAAAGAGGTGCCA  
GCACTCAAGGGAATCCTGCACCACAACAGGCAGGGGACCATCATGACTGCTGAAATAGGC  
AGCAGGAGTCCGAACTATGGCATAAAAATGTCACAGCAACACCACAGGGGGGAACCCCTG  
GGGGGAAGTAGGGGCATGGTGA

The following amino acid sequence <SEQ ID NO. 76> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 16:

RLLLCG YNESHQLQSF PKKDRK\*ICL IKMVFKANYR YHRKKKKNSG R\*DLSSFSYP  
MGFSASRIPC TSRGFCANQW AWPSAHPGLS CGCHGYEDKN DPWFYRALGP  
LLTGFLS\*YF FHLLQNPMWQ VQOGVHQCSG QSE\*\*GTHAR PGRAALKKWF  
VRVWRKGLLP HSALLGPKAL ITPGSGHLS LMLLLSRDRT RGASTQGNPA  
PQQAGDHHDC \*NRQSESELW HKNVTATPQG GTPGGK\*GHG

The following DNA sequence nGPCR-Seq2017 <SEQ ID NO. 17> was identified in *H. sapiens*:

ATTATCTCCATTATGCTCTATCAGTTTCTTTATAGAATATGATTACTACTTAGTGATAAA  
GCTTCCTTTGCTAAGATTTTCAGCCTACGAACCATGATCCAAACCTACTTCTAAAACATA  
TAAACATGCTTTACAAGTATCCTATATATGGAAGAGTCCTTGGAATTATTTGGGTAATTA  
ACCTGGTATTACATGTGTTATTTCTGATCTACTATTGCAAATTGACA

The following amino acid sequence <SEQ ID NO. 77> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 17:

SIC NSRSGNNTCN TRLITQIIPR FPYIGYL\*S MFICFRSRVW IMVRRLKS\*Q RKL YH\*VVII  
FYKETDRA\*W FPYIGYLAS MFICF

The following DNA sequence nGPCR-Seq2018 <SEQ ID NO. 18> was identified in *H. sapiens*:

TCTTGATTTACACAAAAACCATAACTAACTAAAATAGAAAACCTTTTAATTATCTGTATT  
GTTGTATAAAATGTTTTATATATTAGGAATCTAAAAATTTGTTTTTGGCTTTTATCCTCT  
GTCATAGGAAGAAGACATCATGTCTCTGTTTACTGTAACCATTAAATAAAGGCTAATAACA  
GACAGTACATGATGATACTTTTAACTAGGGCAAACAAAAGTAATATTTTAAACAATGAGGT  
TTGGTCTTTGCTATCTATACCTCATGTCTAATTTCCCTACAATGTAAATGTCATTCTCTC  
CTCTCTACCCATTTGTAAGGGTCTCAGTTTCTGCTCTTGTCATGACTTATTTTAAAGGGT  
CACAATAAGGCCAGGTAATTCATATTTTAAAAAATTCATTTAGAATAATTACATCTAAAA  
ATTCACAAGAAAGACAATTTCAATATAAAATAATAAATTACTAATATTGGAATTTCAAGC  
ATTAGTCATGGCAAAAAAGAGATAATTTGTAGCAGAATATTTTAATGGCAACTTTCTTAT  
TCTATCACTTATTTGTGTTCTATTTGTTATGACCAAAGAAATTACTCTATATCCACTACAA  
TTCATAAAACAGGCATGGAAGAAGTCTTTTTTTTCTTGGTGCTCATGTCTAAGAAGATGA  
ACCTCAGAAGTATGTCATTTTCAATACTATGTTCTGAACAGACAGCACACATTATTTTT  
GAATGGACACCAAATCTCAAACATATATAGA



GTGGTATAATCTGCCAAACACAAGCCCCGTTTTACTGTTGCTAATGCACAAAGAGAGGG  
CTGGCAGACATGGCCTGCATGCTCTGTACACCTGCCCATCTGCACCACACTCCCCCATGG  
TCTTGCTGTTGTTGCTGTGTTGGGCTCATCCCGCCTCTCACA

The following amino acid sequence <SEQ ID NO. 81> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 21:

CERRDEPNTA TTARPWGSVV QMGRCTEHAG HVCQPSLCAL ATVKRGLCLG RLYHWSCIPN  
VSLFTRLYKH AQ\*YLCIFTV TSFSYIHKVP IYIPYMYTIN TQMHTHKHTC TELLDQSLLI  
LCIFTV TSFSYIHKVP IYIP R\*STEVQSSP SSDLMII\*LE HLIFLYFIMY LIIT\*LFSEC QLIRRMTCLL  
SCSS\*YPP\*I GFISPLTRCG DWAG

The following DNA sequence nGPCR-Seq2002 <SEQ ID NO. 22> was identified in *H. sapiens*:

AGGCAGAGATTATGTGAGACTTTGCATGATAGGGGTCTGTGCATTCTTGCGCATTCTTCTTA  
GCTCTGTGTGAGCAATATTTAAGTAGTCATTACATAAAATCTATTGAACCTATACAAAGA  
AGTGAACACACTGTGATGAGTTGGCTCCTACCAAGTAAGGCCTGGGGGTTGGTCCTTTAT  
GGTTCCCCAGCACACATCTTAAATAGTTCCCTTTGTGGATAAACTCTCTTTGAATTATTC  
TATTTTGAGTGTGCCATCTGATTCTTGTTAGGACTCTGACTTTAAAAGGTAGAGCATTG  
CTCTCAT

The following amino acid sequence <SEQ ID NO. 82> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 22:

RQRLCETLHD RGLCILAHFL ALCEQYLSSH YKSIEPIQR SEHTVMSWLL  
PSKA WGLVLY GSPAHLK\*F LCG\*TLFELF YFECAL\*FLL GL\*L\*KGRAL LS

The following DNA sequence nGPCR-Seq2003 <SEQ ID NO. 23> was identified in *H. sapiens*:

CAAATGGATTAATTTACATATGGATGAACATGATCTGTTGCTCTCCAGGTCCCAAAGGAT  
ACATAAGAAGAAAAATTTAGTGATGTTACTGGATGATGTCCTTTAAGACAACACAATACA  
ATATCTGAGTATGTACCCTTACGACATAGAGAAGGGATTTTCAAAATATTTTAACTTAAA  
TAGATTCATAAAAGAAATCACCTTCCAACCACTGTTTCTTGTCTCTGGTCAATTAGGGT  
CATAATATTGTTTTCATTGTATTACAAAAGGTAAGAATGTACACTGTATAAATGAATAAA  
TAATATAGATTACTAGATAAGCAGATAAAATAAATACAAAAGCACAAAATACAAAAGCAA  
ATGACCACTCAACTCACTTCCACCCACTTCAAGGAAAACACAGTTCCTTTATCATAGTTA  
CTATTAGAAGTCTTTCCTGTCTTCATTTTCTTACAAGCATGCAGAAATATATATGTGGAC  
ACATTTGTGCCAGAATTCCTTTTTTCTGACACTCACTTTTTCTCCTACTCCACAATA  
TGTCAGGACAATTTTCTACAAGATATAGCAAATGGAGTAACATAGAAATAGAGCAAAACAT  
GAAAACCTCAAACCTCATTAGTGATGATGTTTT

The following amino acid sequence <SEQ ID NO. 83> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 23:

KWINLHMDE HDLLLSRSQR IHKKKNLVML LDDVF\*DNTI QYLSMYPYDI EKGFSKYFNL  
NRFTKRNLHP TTVPCLSIR VILFSLYK R\*ECTLYK\*I NNIDY\*ISR\* INTKAQKYKS  
K\*PLNSLPPT SRKTQFLYHS YY\*KSFSSS SYKHAIEYMW THLCQNSFFS DTHFFPPTPQ  
YVRTIFYKI\* QME\*HRIEQN MKTSNSLVDD V

The following DNA sequence nGPCR-Seq2004 <SEQ ID NO. 24> was identified in *H. sapiens*:

GGGAAGGCTCTTCTAAGAAACCACGCCACACACAAATTAGTAAATTCGACAAAACAGGA  
AACAAACAAAAATTTTGTGTTAAAAGTAACACTAAATCAAAATGAAAAGATAAAAGGCAA  
ACACAAAATTGACAAGGATGCTTGCAACTTGTGTAAGTAAAGGGCTATTTTCTGAA  
TATATTTCTCTATTCTATAATTACTTTCTTTAATATATTTTAAAAAATTCTGTAAAGTAAG  
TATGAAATAGACAAATGATAAGAACAGTTATCAGAAAATGAAATACAAATGATTCAAACA  
TGAAAAGATGCTCGAGCTCAGTCTATAATAATAGACCCTTAAAAGTATAATGAAATATCT  
TTTTTTAAACCCCTATCTGATTAGCAAAGACCAGCAAGTTGGAAAAACAGAGGGCTTTTT  
CAAAGATCAAGCTTGTGAGCCAAACTAAAGGATTCTGTGCTTCAAGAATTACACTGTTT

AGAGTTTGGACTTTTTGAAAAAATGTACATGTCTATGAAATAAATTGGCCTTTTAAAAA  
 AGAGTTTGCAGCAATGCTGAAAGTAGTGGCCAAAAGCATGATCAAGAGATAGGATATTCA  
 TCTAATCTCAAACCATCTCCCTACAAGCTATTTATCAATTACAAAGTGAAAAATACTAAC  
 TGCACAGTGGTGAA

The following amino acid sequence <SEQ ID NO. 84> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 24:

SPLC S\*YFSLCN\*\* IACREMV\*D\* MNILSLDHAF GHYFQHCKL FFKRPIYFID MYIFFKKS  
 \*TV\*FLKAQN PLVWLTSLIF EKALCFNLL VFANQIGV\*K KIFHYTFKGL LL\*TELEHLF  
 MFESFVFHFL ITVLIICLFH TYLQKFLKYI KESNYRIEKY IQENSPLSVT QVASILVNFV  
 FAFYLFILI\* CYF\*HKIFCC FLFCRIY\*FV CGRGFLEEPS

The following DNA sequence nGPCR-Seq2005 <SEQ ID NO. 25> was identified in *H. sapiens*:

ACAAAAGTCACTCCTCTGCTCAAAATCTTTGCACGGTTTCCTGCTGTACTTGGGAATACAA  
 CCCCAGTCCTTTCCCTAGGGCACTAGCCTTGATTGGCCTTGATCACTGCTCCTGCCAC  
 CTCATACCACTTGTCATTTGGCCATTATAGTCCAACAACATCCACTCATCCGCTTTCTCG  
 CAGGCCCTGGAATATTCTAAGCACTTTCCCATCTCAGGGTGTGTGAATGCTGTTTCCTC  
 TACCTGCTCTCATCCTCCCACTCTTATCAACTAAGTTCTCACATACACATGTTTGATCT  
 CAAGCTGTGTGTGTTTACTTGTTGTTATTTGAAGCACTATGCTGAGCCATAGTGGAATC  
 TGAGGTAAAAGGAAAAATAAGTAATATTGACTCTGAGTTTATGTAATATTTGATTTTTT  
 CCCATGAAGGAATTTTGCATTAAATTTTATTTTAAAAATATTGTATTAAAAATTTTATTTA  
 TCTTGATTACTGAGTATTTTGGTGTCCCTTAAATTTCTACCAAAGGACAAGTGCCTCG  
 CTCACCTCACCTTTTTCCAGTACTGGTCATTGGTCTGTCCACCAAACCTGTAAGTAAAC

The following amino acid sequence <SEQ ID NO. 85> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 25:

QKSLCSKSL HGFLLYLEYN PKSFPRALAL YWP\*SLLLPT SYHLSFGHYS PTTSTHPLSR  
 RPWNILSTFP SQGVVNAVSS TCSHPPTLYQ LSSHIMFDL KLCVFTCLLF EALC\*AIVES  
 EVKGKISNID SEFM\*YFDFP P\*RNFAIFI LKILY\*NFIY LDY\*VFWCPL KILTKGQVPR  
 SPHPFSSTGH WSVPPNCK\*

The following DNA sequence nGPCR-Seq2006 <SEQ ID NO. 26> was identified in *H. sapiens*:

CCGAGGGCTTTCCCTGTGAGAATCAGTTTCAAATATGAGGTTGATTTTATTTTCCCGGT  
 TGCAATTAACTGAAAAAATATTTCAAAGCTGTATAATTAAAGCCGGCATTTTATTAGG  
 GTTTAAATAGTCGCTATGAAATTTGCTCCTGCCATTGGTAATTACAACGTACATCTTAA  
 TATCCCCCTGCTTGCCAGCCCCGAGATAACAGCGCTGTGTGTGCTGCATTTGGCTTTGCA  
 GAGAAGGTAAATCCTTATCTTATCTCCATCTGCACACAACAACAGCCTCTCTTCCAAT  
 TCAACCCTAATCTTCTTAGTAAATAATACTGATATTTTTCACCATAACAAAATGGAGCTG  
 GTGCCAGATTCTCACTGATAACACCTCTCGTTTACACACCATTTCTCTGTCAAAATCGGCC  
 CGATTTTTCTTAATCTCTTTAACTTATCAAAAAAATACTTTTTACATTTGGCCATAGTA  
 AGTTCTTTTCTACCAAATAATAATTATAAAATAAATCCCTTTGCCTACAGCTGACTCAT  
 TGAAAAGATACTCAGCCTCCTCATTTTCTTCATGAATGGAAATTGTTCTGCTCTTCAATT  
 TTCTCCCAAGTTTGGGGGTCCACACCAAGGGAATTGTTCCAGGCAGGCCCTGAGCCAT  
 CTTTGATCTTGCAGGGGGTCACTTGGTCCAGGATGTGCATGCTTCTC

The following amino acid sequence <SEQ ID NO. 86> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 26:

EKH AHPGPR\*PPA RYKDGSGPAW EQFPWCGPPN LGGKLKSRTI SIHEENEEAE YLFNESA  
 VGK GIYL\*YYYLV EKNLLWPNVK SIFFDKLKRL RKIGPILTEN GV\*TRGWISE NLAPAPFCY  
 G EKYQYYLLRR LGLNWKRGLW LCADGDKIRI LPSLQSQMQH TQRCYLGAQK QGDIKMYVVI  
 TNGRSKFHSD YFKP\*\*NAGL \*LYSFEIFFS VNCNRGK\*NQ PHI\*N\*FSQG KPS

The following DNA sequence nGPCR-Seq2007 <SEQ ID NO. 27> was identified in *H. sapiens*:

AGTATAAAATCTGGGAGAAAAATCCATCAAATTATAAAATAGAACTAATAATGCTTTCAT

GGGTAAGCAATGAAACAAACCAAGGTAAACCAGCGTTTTAAAGATTGATATTTCTTTAA  
GAAGAGAATGGAAAATAAATTAATAATATACATTTTATATTGATTATCATGAAGTTTTTCT  
AGTTTCTTACCTGAAGTTTCTGGAGGAACTTTAACAGCACTCATTTGACATTAGGAAGA  
AGCCGAATGTAAATAAATCCTAGGTGGATAAGCATAATATAATGAGTAATATGATCTTAA  
ATGGAAAAAAGATTATGTAAAGTTGTTTATACTCTTAACATTTTCCACCCACTACACTGT  
GAAGCAGATTAACTTTTGTCAAAGGTGTAACCTTCT

The following amino acid sequence <SEQ ID NO. 87> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 27:

EGY TFDKRLICFT V\*WVENVKSI NNFT\*SFHFL RSYYSLYYAY PPRIYLHSAS S\*CQMSAVKV  
PPGNFR\*ETR KTS\*\*SI\*NV YFNLSILFL KKYQSLKRWTFGLFHCLPM KALLVLFYNL  
MDFSPRFYT

The following DNA sequence nGPCR-Seq2008 <SEQ ID NO. 28> was identified in *H. sapiens*:

CTTTGAAGGATGAAGAGAAGTTACCAAGTGAACGTGGAGGAAAAGCAGTCCAGGCAGAA  
GGAACAGCTGCTGAAAAGGTGAGGTGCCCAAACAGAAAAGGCATGTGAGGGGAATTCATGG  
GCAGTTCTGGAGAACCAGAGCAGAAGGAGACAGGCTAGAGAATGGCAGGAGATAGCCCTA  
GAAAGCTCTATGTGTGCCAGGTGCAAAAGGCTTGTGCGCCAGCTAAGGAGTCTGCACTTT  
CCCCTGTAAGCAATTAAGTTTAAAGCAGAGGAGGGATTGGTTTGTATTGATCTAGATA  
ATTCAGGGAACAATGAAGGGAGGATATTTGAGGAATCCCAAAGAGCAGGGGCAGAGACTC  
TCATCACAGTTCAGGCCTGACCGAAGGCCATAGGGAGATAGCAGTAAAGCAGGGAAACCT  
ACTGTCCCTTCTTACCTGGAAAACCTATCACCTTCCAGAGTTCCTGTCACCCCAGTGAA  
GGCAACCACTGTCTCTTCTGTTACCTAACACGTGGTACAAACAGCTACCATAGCATG

The following amino acid sequence <SEQ ID NO. 88> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 28:

MLW\*LF VPRVR\*QKRQ WLPSLGVGTGT LEGDSFPGEE GTVGFPAALL SPYGLRSLN  
CDESLCPCSL GFLKYPPFIV P\*II\*INTNQ SLLCLKLLIA YRGKCRLLSW RTSLLHLAHI  
ELSRASCHS LACLILLWFS RTAHEFPHPMP FWFGHLTFSA AVPSAWTAFPRSLGELLFI LQ

The following DNA sequence nGPCR-Seq2009 <SEQ ID NO. 29> was identified in *H. sapiens*:

TATAAGGTAGTTGTGTAGCTTTAAGACTAAGACCTCTACACCTGGAAATTGGTATACTTT  
TTCAATAAAGCATCAGATAGTAAAGTTTGTGGGCCATATGTCTCTGCTGCAACTACTCA  
GCTCTGCCATTGTGCAAAAGCAGCCACAGATAATATGTAAATGAATGAACCTGGCTGTTT  
TCACAAAAACAGGCGGCTGTTAGGATTTGGCTTGCTGACTCTTAACCTGGATTCTCACTT  
CTGAAAGCATGGTCTAGGAGACCAGCAGCATCAGCATTACCTGTGAACCTGTTAGAGATA  
CAGAATGTTGAGCCCCTTCTCATACTGATTGCTCTAGAGTCTAAACCATGTTTCTCAAAC  
TTCAGCATACATAGGACTACCTGGGGAGCTTGTGAAAGTACAGACTCTGAGATAGCAGG  
GCTGAACACCATACTAGCCTTGGGGAGGCCAATGTTATGACCTGGTCAGTGTTCCTTGGGG  
AGTCCGCAATCTATTAGGAGAGACTGCTATATAAACAATAATTTCAATACAGACTAGTA  
AATGCTATAACGGAGGTATCTTCAGCAGAGGAGCACCA

The following amino acid sequence <SEQ ID NO. 89> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 29:

YKVVV\*L\*D\* DLYTWKL VYF FNKASDSKVL WAICLCCNYS ALPLCKSSHR \*YVNE\*TWLF  
SQKQAAVRIW LADS\*LGFS LKAWSRRPAA SALPVNLEI QNVEPLLLI ALESKPCFSN  
FSIHRTHLGI L\*KYRL\*DSR AEHHTSLGEA ML\*PGQCSLG SPQSIRRDY INK\*FQYRLV  
NAITEVSSAE EH

The following DNA sequence nGPCR-Seq2010 <SEQ ID NO. 30> was identified in *H. sapiens*:

ATTTTCATGTTGATGGCAATCATCCAAGAGAGAGCAGGCCATGTCACGCAATAATAGAAC  
AAGGAGAAATGCACACTTTTCTTTTCCATGTCTCTGCTTGCTTGCATTACATTTGCTAATACC  
ACTTTAGCCAGATAAATTAACACAGCGCAGCACATAGTCAAAAAAATAGAAGACCAGAAG

TTACAGAACAACCTGACATGGTTTAGAAAAGCAATTAAGACCATCAGTGAAATCAGTCTAC  
 CAAAAAGGTAATTCATTTGTTTACAAACGCTGTTGAAAAGACTTATCTTCCAAGACAGGA  
 AATGGTTCTCCACTGAAGGGTGAAGACATTTCAATTTTCAGTCATTTGGGGAAGAGTTGG  
 ATCTCCAACGAGTAACTTTTCATGCAAGGACAAGAAATTTAGTAGTGAAATAGAGGTTATTC  
 GTTTTTTTACCATAAAATAATTAATAATCTTGGAGGCAGTTTCCTCATAGCAGTTATTAT  
 GGCAGTTGTGTTTCATTTACAGGAAAACCTGAGAAAACCTAAGATGTTTTTGGGAAAAAAA  
 GTATTTTGAAAGCTTGCGAGTGTTAACTTCCACAATAGATATACTCTTC

The following amino acid sequence <SEQ ID NO. 90> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 30:

R VYLLWKLTLA SFQNTFFSQK HLRVSQFSCK \*TQLP\*\*LL\* GNCLQDY\*LF YGKKTNNLYF  
 TTKFLSLHES YSLEIQLFPK \*LKIEMSSPF SGEPPFVLED KSFQQRCEQM NYLLW\*TDFT  
 DGLNCFKPC QLFCNFWSSI FLTMCCAVLI YLAKVVLANV MQAETWKKKS VHFSLFYYCV  
 TWPALSWMIA INMK

The following DNA sequence nPCR-Seq42 <SEQ ID NO. 31> was identified in *H. sapiens*:

CGCACAGCGC GCAGGTCCTC ACCAGAGCTC TGGTGGCCAC CTCTGTCCCG CCATGCTGCT  
 CACCGACAGT GGCCAGGGCC CACAGCACCA AGAGGCTTGG GCCACAAAGT  
 AAAGGGTTCG GCAGCCTCGC CGGCCGCCAT GTGGAGCTGC AGCTGGTTCA ACGGCACAGG  
 GCTGGTGGAG GAGCTGCCTG CCTGCCAGGA CCTGCAGCTG GGGCTGTCAC TGTTGTGCGT  
 GCTGGGCCTG GTGGTGGGCG TGCCAGTGGG CCTGTGCTAC AACGCCCTGC TGGTGTGCGC  
 CAACCTACAC AGCAAGGCCA GCATGACCAT GCCGGACGTG TACTTTGTCA ACATGGCAGT  
 GGCAGGCCTG GTGCTCAGCG CCCTGGCCCC TGTGCACCTG CTCGGCCCCC CGAGCTCCCG  
 GTGGGCGCTG TGGAGTGTGG GCGGCGAAGT CCACGTGGCA CTGCAGATCC CTTCAATGT  
 GTCCTCACTG GTGGCCATGT ACTCCACCGC CCTGCTGAGC CTCGACCACT ACATCGAGCG  
 TGCACTGCCG CGGACCTACATGGCCAGCGT GTACAACACG CGGCACGTGT GCGGCTTCGT  
 GTGGGGTGGC GCGCTGCTGA CCAGCTTCTC CTCGCTGCTC TTCTACATCT GCAGCCATGT  
 GTCCACCCGC GCGCTAGAGT GCGCCAAGAT GCAGAACGCA GAAGCTGCCG ACGCCACGCT  
 GGTGTTTCATC GGCTACGTGG TGCCAGCACT GGCCACCCTC TACGCGCTGG TGCTACTCTC  
 CCGCGTCCGC AGGGAGGACA CGCCCCTGGA CCGGGACACG GGCCGGCTGG AGCCCTCGGC  
 ACACAGGCTG CTGGTGGCCA CCGTGTGCAC GCAGTTTGGG CTCTGGACGC CACACTATCT  
 GATCCTGCTG GGGCACACGG GCATCATCTC GCGAGGGAAG CCCGTGGACG CACACTACCT  
 GGGGCTACTG CACTTTGTGA AGGATTTCTC CAAACTCCTG GCCTTCTCCA GCAGCTTTGT  
 GACACCACTT CTCTACCGCT ACATGAACCA GAGCTTCCCC AGCAAGCTCC AACGGCTGAT  
 GAAAAAGCTG CCCTGCGGGG ACCGGCACTG CTCCCCGGAC CACATGGGGG TGCAGCAGGT  
 GCTGGCGTAG GCGGCCAGC CCTCCTGGGG AGACGTGACT CTGGTGGACG CAGAGCACTT  
 AGTTACCCTG GACGCTCCCC ACATCCTTCC AGAAGGAGAC GAGCTGCTGG AAGAGAAGCA  
 GGAGGGGTGT TTTTCTTGAA GTTTCCTTTT TCCACAAAT GCCACTCTTG GGCCAAGGCT  
 GTGGTCCCCG TGGCTGGCAT CTGGCTTGAG TCTCCCCGAG GCCTGTGCGT CTCCCAAACA  
 CGCAGCTCAA GGTCCACATC CGCAAAAGCC TCCTCGCCTT CAGCCTCTC AGCATTCACT  
 TTGTCAATGA AGTGATGAAA GCTTAGAGCC AGTATTTATA CTTTGTGGTT AAAATACTTG  
 ATTCCCCCTT GTTTGTTTTA CAAAAACAGA TGTTTCCTAG AAAAATGACA AATAGTAAAA  
 TGAACAAAAC CCTACGAAAG AATGGCAACA GCCAGGGTGG CCGGGCCCTG CCAAGTGGGCG  
 GCGTGTGCTA GCAAGGCCTG CCGGGTGTGC CGCAGTCACC ACAGGGTTCT GAGAACATTT  
 CACAGAAGTG CCTGAGACGC GGAGACATGG CTGGTGTAA ATGGAGCTAT TCAATAGCAG  
 TGACGCGCTC TCCTCAGCCA

The following amino acid sequence <SEQ ID NO. 91> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 31:

MWSC SWFNGTGLVE ELPACQDLQL GLSLLSLLGL VVGVPVGLCY NALLVLANLH  
 SKASMTMPDV YFVNMAVAGL VLSALAPVHL LGPPSSRWAL WSVGGEVHVA LQIPFNVSSL  
 VAMYSTALLS LDHYIERALP RTYMASVYNT RHVCGFVWGG ALLTSFSSLL FYICSHVSTR  
 ALECAKMQNA EAADATLVFI GYVVPALATL YALVLLSRVR REDTPLDRDT GRLEPSAHL  
 LVATVCTQEG LWTPHYLILL GHTGIISRGK PVDAYHLGLL HFVKDFSKLL AFSSSFVTP  
 LYRYMNQSF SKLQRLMKKL PCGDRHCSPD HMGVQVQVLA

The following DNA sequence nPCR-Seq44 <SEQ ID NO. 32> was identified in *H. sapiens*:

CCACAAGACAGTGCCATTTAGTGGCAGCCCTAGGATAAAGATGATACTGTAGGCCAGGGA  
 GAGGTAGACTTGCTTGACTTCTCTGAGAACTGGCAGAGACCTTGTTCTGTGATGTATT  
 CATGTCCACCTTCTCCATGTCCCGGGAGGCTCCCTCCAGGAGCAGAGCTCCACGACGGCT  
 CCCGCTTCTGCTTCCCCTGGAAGGAAGCAAAATGGACAGCATAACATCATGACCTCAGCA  
 TGGCATCAGCCTCTGCCCCAACCCGGCATTCAACTTTGTCCCCCAAAGTCTAGTCAGAGG  
 CCTTGTTTAAAAGAGACCAACCCAGGAAAATAGAGTATTCAACAAGGGCCAAATAATCT  
 TCCAAAGCTGGGCGTCTAGACTTACCAGGTCCAAAGGGTTGGCCTTACATGGACTTGTAT  
 GCTGTCTACTCTAGGCTGGGTTGTGATAGAAATGGTGGTGGGATGACACTGTCATTTAAA  
 CAGGGTACTCAGAACCTCCAGCTCTGTGTCTCCTCCTGCTAGAATCCATTGCATAGAGTC  
 AGAAAATCTCC

The following amino acid sequence <SEQ ID NO. 92> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 32:

TRLWGTKLNA GLGQRLMPC\* GHDVMLSILL PSRGSRSRSGR RGALLLEGAS  
 RDMEKVDMNT SQEQGLCQFS EKYKQVYLSL AYSIIFILGL PLNGTVLW

The following DNA sequence nGPCR-Seq45 <SEQ ID NO. 33> was identified in *H. sapiens*:

CATCCTAAGCTCGGGCACACAAAGTCCCAGAGGCACACAGATTCAGTGGCTTGGGGATAG  
 CATATTCCAACCTTGGGCCCCCTAACTCAAAATCCCCAACACCCCTCAGTCACCTGTGGC  
 TCCAGTCCTGTTTTGGTTTTCCACAGATCACAAAGGCATCAAGGAAGAGGCTCACGGTAA  
 GCCTGGCCTACTCGGAGAGCCACCAGATCCGCGTGTCCCAGCAGGACTTCCGGCTCTTCC  
 GCACCCTCTTCTTCTAATGGTCTCCTTCTTCATCATGTGGAGCCCCATCATCATTACCA  
 TCCTCCTAATCCTGATCCAGAACTTCAAGCAAGACCTGGTCATCTGGCCGTCCCTCTCT  
 TCTGGGTGGTGGGCTTCACATTTGCTAATTCAGCCCTAAACCCCATCCTCTACAACATGACACT  
 GT

The following amino acid sequence <SEQ ID NO. 93> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 33:

S\*ARAHK VPEAHRFSGL GIAYSNSWAP NSKSPTTSLH LWLQSCFGFP QITKASRKRL  
 TVSLAYSESH QIRVSQQDFR LFRTLFLMV SFFIMWSPII ITILLILQN FKQDLVTWPS  
LFFWVVGFTF ANSALNPILY NMTL

The following DNA sequence nGPCR-Seq46 <SEQ ID NO. 34> was identified in *H. sapiens*:

GCTGGGAAGTGCTCCTGGTGCAGCCGGCGGTAGATGGCGGAGGGCACGGTGAGCAGCAAG  
 GCCAGTGTCAGGCTGCCCCACAGGCCACCTGCACCCCGCACGCCCGCTGAACCGTATAC  
 CACCAGGCAGGCCCCGAGAGGCCAGGAAGCAGAGGTTCGGCACTGAGAGCTGCCAGGAGCAGG  
 ACGCTGGCATACATGGTCAGCAGGATGATGGAGGGCAGCGCCCGACAGCCCACTGCACCA  
 TACGGTCAGTGGCCTCCACGGGCAATGGGCACTGCCAGGATGGGCAGAGACAAACAGCAC  
 AGCAAATCCGCCACGGCCAGGTGGAGCAACCAGGTGGCACCCACCCTCCGGCGGGGCCACC  
 TTCCCAGCCACCCAGGCCACCATGGCATTGCCCCGGCACCCCCACCAGGAAGATGGCGGCA  
 TACAGTGGGAGCGGGGCCACGCGCAGCGGGTCGATGGCCAGGCAGGCGCCATCCAGGCAG  
 TCCACAGGGCGGTCCGAGAGGTCGCTGTAATCCCCATACTCGTAGCTGACAGAATCG

The following amino acid sequence <SEQ ID NO. 94> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 34:

DSVSYE YGDYSDLSDR PVDCLDGACL AIDPLRVAPL PLYAAIFLVG VPGNAMVAWV  
AGKVARRRVG ATWLLHLAVA DLLCCLSLPI LAVPIARGGH \*PYGAVGCRA LPSIILLTMY  
ASVLLLAALS ADLCFLALGPAW WYTVQRAC GVQVACGAAW TLALLLTVPS AIYRRLHQEH FP

The following DNA sequence nGPCR-Seq47 <SEQ ID NO. 35> was identified in *H. sapiens*:

GGCTCCACTGGCCTTTCGGCAAGGGGCTCTGCAGGCTGACGGCGTTTGTGCTCTACACCG  
 ACACCTACGGGGGGTCTACCTCATGGCCTGTGTGAGCGTGGACCATTACCCAGCTGTGGT  
 CTGTGCCCACTGGGGCCCGTGCTCCGCACGGCTGGCCGCGCCAGGCTGGTCTGCGTGCC

CATCTGGACCTTGGTGCTGCTGCAGACGATGCCCTTGCTCTTGATGCCCATGACCAAGCC  
 GCTGGTGGGCAAGCTGGCCTGCATGGAGTACAGCAGCATGGAGTCAGTCCTCGGGGCTGC  
 CCTCATGGTCTTGGTGGCCTTTGCCATTGGCTTCTGTGGGGCAGTGGGGATCATCTGT  
 CCTGCTATATGAAGATCACCTGGAAGCTGTGCAGCACAGCTGGGAGAACCCAGTGACCAG  
 CGGGAAAGGACACCAACCGCGGGGACCCCGGGAGGACCCAGTGACCAGCAGGAAAGGAC  
 GCCACCGGCGGGGCAGCCCAGGAGGACCCAGTGACCAGCGGGAAAGGACACCACCG

The following amino acid sequence <SEQ ID NO. 95> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 35:

AP LAFRQALQA DGVICALHRHLR GVYLMACVS VDHYPVVCA HWGPCLRTAG  
 RARLVCAIW TLVLLQTMPL LLMPMTKPLV GKLACMEYSS MESVLGAAPH GPGGLCHWLL  
 WASGDHPVLLYEDHLEAVQH SWENPVTSGK GHHRRGSPGG PSDQQERTPP AGQPRRTQ\*P  
 AGKDTT

The following DNA sequence nGPCR-Seq48 <SEQ ID NO. 36> was identified in *H. sapiens*:

GAGGCAGGCT GCAGGGAAGT AAGGAGGAGG CATGGCACCT TCTCATCGGG CATCACAGGT  
 GGGGTTTTGC CCCACCCCTG AACGCCCTCT GTGGCGCCTT CCACCCACCT GTAGGCCAG  
 AAGGATGTCG GTCTGCTACC GTCCCCCAGG GAACGAGACA CTGCTGAGCT GGAAGACTTC  
 GCGGGCCACA GGCACAGCCT TCTGCTGCT GGCGGCGCTG CTGGGGCTGC CTGGCAACGG  
 CTTCTGTTG TGGAGCTTGG CGGGCTGGCG GCCTGCACGG GGGCGACCGC TGGCGGCCAC  
 GCTTGTGCTG CACCTGGCGC TGGCCGACGG CGCGGTGCTG CTGCTACGC CGCTCTTGT  
 GGCTTCTCTG ACCCGGCAGG CCTGGCCGCT GGGCCAGGCG GGCTGCAAGG CGGTGTACTA  
 CGTGTGCGCG CTCAGCATGT ACGCCAGCGT GCTGCTCACC GGCCTGCTCA GCCTGCAGCG  
 CTGCCTCGCA GTCACCCGCC CCTTCTGGC GCCTCGGCTG CGCAGCCCGG CCCTGGCCCG  
 CCGCCTGCTG CTGGCGGTCT GGCTGGCCGC CTTGTGCTC GCCGTCCCGG CCGCCGTCTA  
 CCGCCACCTG TGGAGGGACC GCGTATGCCA GTGTGCCAC CCGTCGCCGG TCCAGCCCGC  
 CGCCACCTG AGCCTGGAGA CTCTGACCGC TTTCGTGCTT CCTTCGGGC TGATGCTCGG  
 CTGCTACAGC GTGACGCTGG CACGGCTGCG GGGCGCCCGC TGGGGCTCCG GCGGGCACGG  
 GGCGCGGGTG GGCCGGCTGG TGAGCGCCAT CGTGCTTGCC TTCGGCTTGC TCTGGGCCCC  
 CTACCACGCA GTCAACCTTC TGCAGGCGGT CGCAGCGCTG GCTCCACCGG AAGGGGCCTT  
 GGCGAAGCTG GGCGGAGCCG GCCAGGCGGC GCGAGCGGGA ACTACGGCCT TGGCCTTCTT  
 CAGTTCTAGC GTCAACCCGG TGCTCTACGT CTTACCGCT GGAGATCTGC TGCCCCGGGC  
 AGGTCCCCGT TTCCTCACGC GGCTCTTCGA AGGCTCTGGG GAGGCCCGAG GGGGCGGCCG  
 CTCTAGGGAA GGGACCATGG AGTCCGAAC TACCCCTCAG CTGAAAGTGG TGGGGCAGGG  
 CCGCGGCAAT GGAGACCCGG GGGGTGGGAT GGAGAAGGAC GGTCCGGAAT GGACCTTG  
 ACAGCAGACC CTACAACCTG CTGCCCTTCC CTGTCCCTTT CCACCCCCACCCACCCTCC  
 AGAGGTCAGT GTTCTGGGAC ATTTGGGGAC CCTTCTTTGA CTAGAGTTTG GATCTGGCTG  
 GGTAGGATTA CTATACACTT GGGGCAGGCC CAGGCTCCTC CAACTGAGG GATTATGAGG  
 GTGGTGATGG TCCCTGTAA GGAATATTGT GTGCTTGCAA GTTGGCATGT ACCCATGTGC  
 CAGCATTGCT

The following amino acid sequence <SEQ ID NO. 96> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 36:

MS VCYPGNET LLSWKTSRAT GTAFLLLAAL LGLPGNGFVV WSLAGWRPAR GRPLAATLVL  
 HLALADGAVL LLTPLFVAF TRQAWPLGQA GCKAVYYVCA LSMYASVLLT GLLSLQRC LA  
 VTRPFLAPRL RSPALARLL LAVWLAALL AVPAAVYRHL WRDRVCQLCH PSPVHAAHL  
 SLETLTAFVL PFGLMLGCYS VTLARLRGAR WSGRHRGARV GRLVSAIVLA FGLLWAPYHA  
 VNLLQAVAAL APPEGALAKL GGAGQAARAG TTALAFFSSS VNPVLYVFTA GDLLPRAGPR  
 FLTRLFEGSG EARGGGRSRE GTMELRTTPQ LKVVGGQGRGNDPGGGMEKD GPEWDL

The following DNA sequence nGPCR-Seq49 <SEQ ID NO. 37> was identified in *H. sapiens*:

CCAGCCGTCCAGGCGACGCGGGCCAGCACCAGGAACCAGGTGACGCTGGCCACGTGGTAG  
 CGCGGGCAGACCAGGGGGCAGCGCTGTTGCGCAGCAGGCCGAGGCGTTGGCGGGCAGC  
 GGCACGGCCGGCAGGGTCCGCGGTCCACACCAGGGCGCAGAGGACGGCCGAGGCGTGTCTG  
 GGCCGGCAGCCCTGGTAGCAGGCGGGGAAGAGGTCCGAGAGGCAGCGCTCCACGCTGAAG  
 GCCGCCAGCAGCCAGAGCCCCACCGCGAACACAGGAAGGTGAGCACGAAGTAGAGTGTG



TCCTGGGCGCCAGGGCAGCCTGAGCCACGGAGAAGCCACACGGCAGGAGAGGAACAGG  
AAGTCGGCGGCGGCCAGGTGCAGCAGGTAGATGGAGAAGGGGCCCTTCTTGATGCGGAAG  
CCGAGGTTCCAGAGCACCAGCCCGTTACCTACCGGTCCCCGAGGCCACGATCAGCGTC  
AGGTAGAAGACCACACTGTCGAAGGTTCTCCAGAGGCCGAACAGCCCAAACATCCTGGCC  
GGCTCAGAGGGTGTGGCGAGGGACCTGCAAGATGGAGACAGACATGGTCCTCAGGAGTC  
TTTTGTGCCCCCGCTGGGGGCACAGGGAGTGGCAT

The following amino acid sequence <SEQ ID NO. 97> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 37:

MPLPVPP AGAQKTPEDH VCLHLAGPSP APSEPARMFG LFGLWRTFDS VVFYLTLLIVG  
LGFPVGNGLV L WNLGFRICK GPFSTYLLHL AAADFLFLSC RVGFSVAQAA LGAQDTLYFV  
LTFLWFAVGL WLLAAFSV ER CLSDLFPACY QGCRPRHASA VLCALVWTPT LPAVPLPA NA  
CGLLRNSACP LVCPRYHVAS VTWFLVLARV AWTA

The following DNA sequence nGPCR-Seq50 <SEQ ID NO. 38> was identified in *H. sapiens*:

ATGTCTACAGAAACCCCTTCGCCATCTACCTCCTGGTACGTGGCCTGCAGCAGGATCTCA  
TCTTCCTTGGCTGCCACATGGTGGCCATCGTCCCCGACTTGCTGCAAGGCCGGCTGGACT  
TCCCGGGCTTCGTGCAGACGAGCCTGGCAACGCTGCGCTTCTTCTGCTACATCGTGGGCC  
TGAGTCTCCTGGCGGCCGTCAGCGTGGAGCAGTGCCTGGCCGCCCTCTTCCAGCCTGGT  
ACTCGTGCCGCCGCCACGCCACCTGACCACCTGTGTGTGCGCCCTCACCTGGGCCCTTT  
GCCTGCTGCTGCACCTGCTGCTCAGCAGCGCCTGACCCAGTTCTTCGGGGAGCCCAGCC  
GCCACTTGTGCCGGACGCTGTGGCTGGTGGCAGCGGTGCTGCTGGCTCTGCTGTGTTGCA  
CCATGTGTGGGGCCAGCCTTATGCTGCTGCTGCGGGTGGAGCGAGGCCCCCAGCGGCCCC  
CACCCCGGGGCTTCCCTGGGCTCATCCTTCTCACCCTCCTCCTCTTCTCTGCGGC  
CTGCCCTTCGGCATCTACTGGCTG

The following amino acid sequence <SEQ ID NO. 98> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 38:

VYRNP FAIYLLVRGL QQDLIFLGCH MVAIVPDLQ GRLDFFGFVQ TSLATLRFEC  
YIVGLSLLAA VSVEQCLAAL FPAWYSCRPR RHLTTCVCAL TWALCLLLHL LLSSACTQFF  
GEPSRHLCRT LWLVAAVLLA LLCCTMCGAS LMLLLRVERG PQRPPRGGFP GLILPHRPPL  
PLLRPALRHL LA

The following DNA sequence nGPCR-Seq51 <SEQ ID NO. 39> was identified in *H. sapiens*:

AAGGTTCTAGGAACACATTGCCCCATAATTTCTATGGTTGTTTATACCCTGTCTATCTGA  
ATGTTTCCTCTGCAGCTGCAATCCCAAGTGTACCAAAACAGAGAAGTGATGAAGTTGACAA  
AGATGGTGCTGGTGCTGGTGGTAGTCTTTATCCTGAGTGCTGCCCTTATCATGTGATAC  
AACTGGTGAACCTACAGATGGAACAGCCACACTGGCCTTCTATGTGGGTTATTACCTCT  
CCATCTGTCTCAGCTATGCCAGCAGCAGCATTAACCCCTTTCTCTACATCCTGCTGAGTG  
GAAATTTCCAGAAACGTCTGCCTCAAATCCAAAGAAGAGCGACTGAGAAGGAAATCAACA  
ATATGGGAAACACTCTGAAATCACACTTTTAGGAAAGTACATGGATCACCATGAGTCTAG  
ACATGATTGTCTATCTTACTGGTATTATTAGAAAGGGCAGGTGTACCGATATGTTTATGC  
CCATTCTTCTGTGTAAGTGTGACTCTTAGCAGCATGGAAGAGAAGTGTAAACCATGCAAA  
TACAAATGAGCTTAATATGCTAACTTTAGCAAGATGTAAATGTTGATCTATATTGTGGGT  
AGGGAATGGGATAGTCTGAGATACCCAGGCTTCATGATGGTGTATATTATTCAGCATAT  
TATAAACTAGTCACTAATG

The following amino acid sequence <SEQ ID NO. 99> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 39:

GSRNTLPH NFYGCLYPVY LNVSSAAAP VYQNREVMKL TKMVLVLVVV FILSAAPYHV  
IQLVNLQMEQ PTLAFYVGYL LSICLSYASS SINPFLYILL SGNFQKRLPQ IQRATEKEI  
NNMGNTLKSH F\*ESTWITMS LDMIVYLTGI IRKGRCTDMF MPILLVYL\*L LAAWKRSVTM  
QIQ\*A\*YANF SKM\*NVDLYC G\*GMG\*SEIP RLHDGVYYFS IL\*TS\*

The following DNA sequence nGPCR-Seq52 <SEQ ID NO. 40> was identified in *H. sapiens*:

CAATTTTCTATTGCCTCTCTGGCCTGTGCTGACTTCTTGGTAGGTGTGACTGTGATGCTT  
TTCAGCATGGTCAGGACGGTGGAGAGCTGCTGGTATTTTGGAGCCAAATTTTGTACTCTT  
CACAGTTGCTGTGATGTGGCATTGTTACTCTTCTGTCTCCACTTGTGCTTCATCTGC  
ATCGACAGGTACATTGTGGTTACTGATCCCCTGGTCTATGCTACCAAGTTCACCGTGTCT  
GTGTGCGGAATTTGCATCAGCGTGTCTGGATTCTGCCTCTCACGTACAGCGGTGCTGTG  
TTCTACACAGGTGTCAATGATGATGGGCTGGAGGAATTAGTAAGTGCTCTCAACTGCGTA  
GGTGGCTGTCAAATTATTGTAAGTCAAGGCTGGGTGTTGATAGATTTTCTGTTATTCTTC  
ATACCTACCCTTGTATGATAATTCTTTACAGTAAGATTTTCTTATAGCTAAACAACAA  
GCTATAAAAATTGAAACTACTAGTAGCAAAGTAGAATCATCCTCAGAGAGTTATAAAATCAG  
AG

The following amino acid sequence <SEQ ID NO. 100> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 40:

QFSIASLACA DFLVGVTVML FSMVRTVESC WYFGAKFCTL HSCCDVAFYCY SSVLHLCFIC  
IDRYIVTDP LVYATKFTVS VSGICISVSW ILPLTY SGAV FYTGVNDDGL EELVSALNCV  
GGCQII VSQG WVLIDFLLFF IPTLVMILY SKIFLIAKQ AIKIETTSSK VESSSESZYKI R

The following DNA sequence nGPCR-Seq2021 <SEQ ID NO. 41> was identified in *H. sapiens*:

ATTATTCATTCCCTTTAGACATTAAACATTTCATTGAGCACCTGCTGTATGCAAAGCACTGG  
GCACCCACACTAAGGATGAAAACCAAGGATAAGTAAGACAGCATGTAGATTCTAGCTGCC  
TGGTTGAGAGGTTACAGTCAGAAAGTCCTTATAGCCATTGAGTACATACAGATAGGGAT  
AGGAGAGGAAATGGGGTGAGCACAGGGAAGGGATAAGTATAGGGTCAGGGCTACCACCCT  
TCTTTTCCCCATGACCCCATGGGGGCAAATATGTCCTGTCTCTCCTGGGTGGGTGATGT  
TCACTGCCCCCTTTCTCCATTCCAATTGGAACCTTCTAGATTGAGCCCGAAGCTAGACTTGC  
AGATCACAATTTTAAGAAAGTTGGATGTTCTGCAACAGGTTACACAGGAGCTATCATTGA  
GTCATCTTTTCGTTCAACCAATCATTTATTCATTTCATTCAACAAGTATTTCCCAAG  
CACCAATTCCATCCTCAACAAAAGACTCCTAAATCACAGGCCAGATGAAAGACGTCACTA  
TTCATTGGAAAAATTGAGCTGATTATGATCCTGGTGGAGTGACTCAGCGTCC

The following amino acid sequence <SEQ ID NO. 101> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 41:

YSFL \*TLNIH\*APA VCKALGTHTK DENQG\*VRQH VDSSCLVERL QSESPYSHSV  
TYR\*G\*ERKW GEHREGISIG SGLPPFFSP\* PHGGKYVLSL PGWVMFTAPF  
SPIGTSRLS PKLDLQITL RKLDVLQQVT QELSLSHLFV HPIIYSFIHS  
TSISQAPIPS STKDS\*ITGQ MKDVTIHLEK FS\*L\*SWSEC TQR

The following DNA sequence nGPCR-Seq2022 <SEQ ID NO. 42> was identified in *H. sapiens*:

ATGGAATTAGAAAAATGAGAAAGTAATCAAAGATTGTAAAAAATACTGAGATAGAATTAA  
AAGGAATAAAAAATTTGTAAATCAAATATGTAATTCTTTTAAAGTTGACACATTGTAATT  
TTATATATTTGTGGAGTATGATTTGGTACATTAATTTGTATATGTGTTGTATGCTGATCA  
AATCAGGGTATGTACTAGGCCATTTTGCATTGGTATAAAGAAATACCTGAAGCTGGGTA  
ATTTAAAGAAAAGAAGTTTATTTGACTCATGGTCTATAGGCTGTGAACAACATCTGCTT  
GGCTTCTGGTGGGAGCCTCGGAAAGCTTTCAATAGTGGCAGAAGGGGAAGGGGAAGCTGG  
AGTATCACATGGTGTGAAAGGGAGCAAGAGAGAGAGAGAGAGACAGGAGGTCCTAGACTT  
TAAACAACCAGATCTTGTGTGAACTATTTTCATGTGACTAAGAACTCATTTCATACCAAG  
CGGAAGGTGCCAAGTCATTTCATGAGGGATCCACCCTCATAATACTACCTTCCACCAAG  
TCTCACTTACAACATTGGGAATGACATTTTCAGCATAAGATTTGGAGAGGACAAATGTCCA  
AACTATATCAGGGTAGTTAGCTCATTTCATCACCTCATGTCTTTATCATTTT

The following amino acid sequence <SEQ ID NO. 102> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 42:

NDKDMR\*\* MS\*LP\*YSLD ICPLQILC\*N VIPNVVSETW WKVYYEYEGGS LMNDLAPSAW  
\*\*MSS\*SHEI VHTRSGCLKV \*DLLSLSL APFHTM\*YSS FPFPFCHY\*K LSEAPTRSQA

DVVHSL\*NHE SNKLLFFKLP SFRYFFIPMQ KWPSTYPDLI SIQHIYKLMY  
QILHKYIKL QCVNLKRITY LIYKFFIPFN SISVFFTIFD YFLIFLIP

The following DNA sequence nGPCR-Seq72 <SEQ ID NO. 43> was identified in *H. sapiens*:

ATGTAAAGGGCCTTTTGAAGAAAAGTGTAACTCCACTGTGGGGCCATTAAAGAAGTCT  
TGAATACATAGATTAAATCCCTTGGTTTGAGATAATAAACTCTCAATATTTACCATGCT  
TTTGTCATTTAAATACAAATTGTGGACATTTTACCAGGGCAACAGATGAGTTACAACCA  
TCCTTTTATCAGTTGCACAGAATCTTATTGAATGGGTATATCCTACTTATGTCAGTTCTT  
AGTAGTGAATATTAGGTTGTTTCTAATTTTTCCATATTAGAAACAATGTTAAAAGAACTT  
GCTCTTATTTGTGTTTCTTATGTCCTACACTAGTATTCAATACCTTTCTAGGAGTGGAAT  
TCCTGGCTCAAACAACTGCCTTCCAAGAAGATTACTAATATACATTCTACTAATGCT  
GTTTTAGAATGCCTTTTTCTCCATATTCCTATGAACACTGAGTGTTCTCTTTTAACT  
TTTTTCTCAATCTACTAGACTCCTTCTAAAATATATACTTTTCAAAACGCCTGCTTAC  
CCTTCAATATATTGTCTTTTAAAAGATTATTTGTAGGTACTGTTTATGTGTTAGGTTGGT  
GGAAAATTAATTGTGATTTTGCAAAAAGTGAATTACTTTGCACCAACCTAATATTATACAT  
GTAGA

The following amino acid sequence <SEQ ID NO. 103> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 43:

M\*RAFWKKTV NSTVGPLKKS \*IHRNLPLV\* DNKTLNIYHA FVI\*IQIVDI FTRATDELQP  
SFYQLHRILL NGYILLMSVL SSEY\*VVSNF SILETMLKEL ALICVSYVYT SIQYLSRSGN  
SWLKQTAFAQE DLLIYPTNA VLECLFLHIP MNTECSLFNT FFPQSTRLLL KYILFKTPAY  
PSIYCLKDY L\*VLFCM\*VG GKLIVILQKL QLLHQPNII HV

The following DNA sequence nGPCR-Seq2024 <SEQ ID NO. 44> was identified in *H. sapiens*:

TCTCTTTTGGTTCCTTTTTTCTCATTACCTACCTCTTCTCCTTTGCTGGTCCTTCACA  
TCTTCCTGGCCTCTTCTTTGTCTGCATTTACTTGTGTGGTCTTCCGTCTTTAACACATCT  
ACCCATCAGCTACTCCCAAATGTATATATCTATTCCCGGACCTTTCTCTGAACTCCAGA  
TTTGTATTTCCTCACTACCTACTCAATAGTACCTTCTTGGATATTTATTAATAATTGAATC  
ACGACATGCCTAAAATTGAACTTCCTATCTCTGAAGCAAAGCCCTATCCTTCTATTGTTT  
TTCCCATCCCAATAAATTGCCACTCATTCTTCCAGATGCTTGGGCAACATTTTTGCAGTC  
ATCTTTGATTTCTTCTTTCTTTGATATCCACATACATTACACAGCATATGCTGCCCTG  
TCTACATTTTAAACATAACCAGCATCCAACATTTCTACAATTGCTACTAGCCTTGATAG  
CCCTATCCTCTCTCCTCTGGATTACTGTATTAGCCTCTGAACTGGGATCCCGCTTCAACC  
CTTTCCAACCTCCTCCTCCCAAAGCCAACCTCTGTGAACAGCAGCTAAAGTTGTTCTTTCA  
AAAA

The following amino acid sequence <SEQ ID NO. 104> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 44:

LFWFPFLI TYLFSFAGPS HLPGLFFVCI YLCGLPSLTH LPISYSQMYI SIPGPFL\*TP  
DLYFQLPTQ\* YLLGYLLKFE SRHA\*N\*TSY L\*SKALSFYC FSHPNKLPLI LPDAWATFLQ  
SSLISFFL\*Y PTYIHHMLP CLHFKHNQHP TFLQLLLALY SPILSPLDYC ISL\*TGIFLQ  
PFPTSSSQSQ LCEQQLKLFF QK

The following DNA sequence nGPCR-Seq2025 <SEQ ID NO. 45> was identified in *H. sapiens*:

CAACCCTGGTAAAGCATCGCACCTTGGGTTATGTACCTCAGGGTTATTTGACGCACTGGG  
CTAAAATGTTGAAGGACATCCTGTTTCCAGGTGGGGACTGGAACAGAGCCTGGACTGTTT  
TAGCCAATGGCTCCTTACCTCAGGATGTTGCATTTCCAGCACATTCTGGTTGGTGTGAG  
AACTACAAACAAGAAAGTGGGAAGAACTGTTCTGCACCATTTATGTAACTTCTAGGAAA  
GCAAACTAATGTATTGTGACAGAAGGAAGATGAGTGATTGCTTAAACACAAGGGAGGAAT  
GCTACACAGGGAAGGTCTGGAGAGTTGGATTACAAAGAGAGACAAGGACACTTTTGGGAG  
AGATGGATATGTTTATTATCTTGCTTATGGTGATAGTTTCATAGGTCCATAAATCCCAA  
AGCATCTCATTGTACACTAACTATGTATGATTTATTTATTAATTATACCTCAATAAAG  
TTGTTTTTTAAAAAAGTTACCACTTAATCCTGTAAATAGACCAGGAAGGCAATTAATTAA

TATTTTCTAGTTTACTTTTGAGAACTAAAGCTTAGTA

The following amino acid sequence <SEQ ID NO. 105> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 45:

NPGKASHL GLCTSGLFDA LG\*NVEGHPV SRWGLEQSLD CFSQWLLTSG CCIPSTFWLV  
LRTTNKKVGR TVLHHLCKLL GKQTNVL\*QK EDE\*LLKHKG GMLHREGLES WTKRDKDTF  
GRDGYVYYLA YGDSFIGP\*I PKASHCTLTM YDLFILIPQ \*SCFLKKLPL NPVNRPGRQL  
INIF\*FTFEK LKLS

The following DNA sequence nGPCR-Seq2026 <SEQ ID NO. 46> was identified in *H. sapiens*:

AGTTTATGGACCAGCCTTCCCTGTGAAATTTGACTTTTCCCTCTTTGCTGAATTGGTCAG  
GTTAACAATGGTTACCCCTGGATTACAGGAAGGGCATGTGCTAAAAGCCTCTTTGGAGAC  
CCACATGGCCCTCAGATGAGCAATTGTTGAGATTCTTTTCTTTTCTTTTCCATGGGAA  
TAAGCTTTCCCTCTCTCCAAAGTACATGTTTTAGGCTTTTTTATTTTCTTGCTACTCCCAA  
GGACCTGGTGATATTTTCTTTACCATGCATTAAACAGAATCTGTGAGTCTTTTCTGGAA  
AAAAAAAAGGCAGGAGGGAACATACTAGTTAAAAAGTTTCTGGGTACACTACCAAGATGT  
ACCTATTTATTGATATACAAATGGCATAAGTTATTGAATGCTTGCTATAGGCATTCTCTA  
AGAACTTTGTAAGAATTGACTTACATGAGCTACTTCATAGCAGTTCGATGATATACATGT  
TGTTATTATCACCACTTTACAGATAAGGAAATAGAGACAGACATACTGAATGACATGCTC  
AACGCCACTCCACTAGCAAGTGGCAGAACCCAAGC

The following amino acid sequence <SEQ ID NO. 106> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 46:

LGFCHL LVEWR\*ACHS VCLSLFPYL\* SGDNNNMYII ELL\*SSCKS ILTKFLENAY SKHSITYAIC  
ISINRYILVV YPETF\*LVCS LLPFFFPEKT HRFCLMHGKE KYHQVLGSSK KIKKPKTCTL  
ERGLIPMEK KKKRNLNNCS SEGHVGLQRF\*HMPFL\*SR GNHC\*PDQFS KEGKVKFHRE  
GWSIN

The following DNA sequence nGPCR-Seq2027 <SEQ ID NO. 47> was identified in *H. sapiens*:

CGTGGCAGGTTGCACCAACTTTTAAACCCACAAGCAATTATGAGAGGTTCAATTTCCCTTC  
ACTCACAACAAAAGTGCATCTTATCACGTTTAAAGAAATCTTTGCCATTTTGCTGGGATT  
AACTTGCTCTCATTATTCTATATTTGCATTTCTAAGCTATTAAGCTTGAATAATTCCA  
TATGGGCCATTTGTATTTGCAAAACAAACACTACCCCTTCAATGATTTGCCTGGCTTCT  
GCCTTCACTAGTATTTATTGCCTCTTTAAACACTAAGTTAATAGTTTTATTGTTCCTT  
TGTCAGCCTCCTCTGAAAATGATGACACTTTTCAAACAGTACCACATCTTTCTACTCATT  
TCAGTTCAATGCTCATATAGGGCATTGTGCATACTGACTGTCCAAGCTGTGCACTAAGCA  
GGTGAATCTTCCCTGCCCTTGGATCGAGTAATAACAGAACTTAAAGAAGGCTCCTGGGTA  
GGAAAATCATGGGTACTGGTAACAGAGAGATATGGATTCAAATCCTGGCTTCTCTACTTA  
CTCACTCAA

The following amino acid sequence <SEQ ID NO. 107> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 47:

VAGCTNF \*THKQL\*EVH FPSLTTKLHL ITF\*EIFAIL LGLTCLSLFY ICISKLLSLN NFHMGHLYLQ  
NKHYPFNDA WLLPSLVFLA SLKH\*VNSFI CSFVSL\*K\* \*HFSNSTTSF YSFQFNAHIG  
HCAY\*LSKLC TKQVNLPCPW IE\*\*QNLKKA PG\*ENHGYW\* QRDMDSNPGF STYSL

The following DNA sequence nGPCR-Seq2028 <SEQ ID NO. 48> was identified in *H. sapiens*:

ACACCAGGAACACACTTGGATGCCCATCGGCCCCAGGTTTCTTCTGCTGGTCTGTCTGG  
CCATGTGTTTGGGTTTGAAAGTGTCAGATTGCCTGGTTCTCCTGGATCTTCACGCAAAA  
GAAACGAGCACATGATGGTGACCTGGAACCTCCTCGCTGGCGTCACTGCATTTTGGCA  
AGCCTGTAACTGTCCTGTCTGCATTCTGGGCTCCAAGGCTCAGTCTCTGATCTTTCTGT  
ACCTGTCTTCCCCTAGGCTGCCTTCTCTTTTCTTATTACAGTAAAGTTCTGTATGT  
ACTGCAGCATATTCATTTATTGGGAATTGAATATATTTCTTCTATGCCAGGATTTAAAA

TTAGGATTGTTAATATTGTGGTTTGTGCTCTGGTTACAGAATTTCTAAGGTTTGGGTGTT  
 CTATCCCAGCACCATATTTTCTCAAAGCCTTGCTCTCTGCAGTGGGTGATTTTGCACAA  
 TCAAACCTTCTCAGGTACTTTCTTCTGAGCTCCAGGTCCCCATATCCAACATCTACTCAAC  
 ATCTTATTCTGAGATGTTACCTCAAACATGTGAGAATCAACATGTCAACATGTCAATAC  
 CTCTTGCAAGGATTTCTAACTCCACAGATGGTATCCGTCCAATTGTGCAAGCCAAGAGCA  
 AGGCGCCAGCTGGGACGTTTCCCATAACCAATCTATCATCATGTCCCATTAATTTTACC  
 AGTAGT

The following amino acid sequence <SEQ ID NO. 108> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 48:

TRNTLGC PSAPGFFCWS CLAMCLGLKV SRLPGSPGSS RKRNEHMMVT WNSPRWRHCI  
 FAKPVTLSA FWAPRLSPLI FPDLSF\*AA FLFLITVKF CMYCSIFHLL GIEYISSMPG  
 FKIRIVNIVV CALVTEFLRF GCSIPAPYFL KALLSAVGDF AQCKLLRYFL LSSRSPYPTS  
 TQHLILRCSP QTCENQHVNM SIPLAGFPNS TDGIRPIVQA KSKAPAGTFP IPNLSSCPIK FYQ\*

The following DNA sequence nGPCR-Seq2029 <SEQ ID NO. 49> was identified in *H. sapiens*:

CACTTGAGAGGTGATCTCATCTATAAAGAGAGTATATTTAAGGAGCTAATTTACACAGGT  
 AGATGACAAAACCTTACACCAACTCTTCAAGCACAGGAAGCACCTTACACGAAACACTTT  
 GGTGAATGCTCATTTCTACTCAATTTTAACTAAATTCAACTGAAAATACCTCATTTTAA  
 AGAGGATATTGAGTTGACTCACTGGGGAAAAACAATTGGATAAACTGGTTGTTTGCCTT  
 ACCGAGATAAAATCACTGATAAGTTGAATGCATTTTTGTAATGTGTCAAGACAGATGGCT  
 TTAATAGTCACACATATTAAGTATGGGATATGTGGCGAGGAATAGAGGTTTTATATACA  
 GAAGTAAGAACTGTAGAGGGGGGAAACCATGAAAGATAGAAAAAGATGAGAAGAAAGAGG  
 ACAGAGATGATACACTCAAGAAAAGCTGCCTATTTTCAAGTAGAATATATTGTAAGCCCC  
 AGTATAGTTGAATGCTTACAAGAGGAGAAGCTTTTAGCAACTCCTTTGTTTCTTAATGTA  
 TATATTATATCTGAACAGAATGCTTCTTTCCTGTGAGCTAGACAGTAGGAAGTAATCTAT  
 CTTCAACCTTGTTTGGTGTGTAAATAAACTTT

The following amino acid sequence <SEQ ID NO. 109> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 49:

KFYLHTKQ G\*R\*ITSYCL AHRKEAFCSD IYTLRNKGV AKSFSSCKHS TILGLTIYST  
 LK\*AAFLECI ISVLFLLIFF YLSWFPPSTV LTSVYKNLYS SPHIPYLICV TIKAICLDL  
QKCIQLISDF ISVRANNQFI QLFFPSESTE YPL\*NEVFSV EFSLKLSRNE HSPKCFV\*GA  
 SCA\*RVGVRF CHLPV\*ISSL NILSL\*MRSP LK

The following DNA sequence nGPCR-Seq2030 <SEQ ID NO. 50> was identified in *H. sapiens*:

GTGATACGGTTCGGCTGTGTCCCCATCCAAATCTAATCTTGAATTACAGCTCCCATAATT  
 TCCATGTGTTGTGGAAGGGGCTGGTGGGAGATAACCGAATCATGGGGGCGGCTCCCCC  
 ATACTGTTCTCTTGGTAGTGAATAAGTCTCATAAGATCTGATCGTTTTATAAGGGGAAAC  
 CCCTTTTGCTTGGTTCTTATTCTCTCTTTGCTGGCTGCCATGTAAGATGCTTCTTTGCTC  
 TTCCTTTGTGTTCTGCCATGACTGTGGGGCCTCTCCAGCCATGTGGAAATGTGAGGAAAT  
 TAAACCTCTTTCTTTATAAATTGCCAGTCTTGGGTATGTCTTTATCAGCAGTGTGAAA  
 TGGACTAATACACCAGAAGTTCAAGACTAGCTTGGGCAACACAGTAAGACCCTGTCTGTA  
 TAAAGAAAGAATAAGAAAAAGAAAAATATCAGTGGCATAATATTAGACAGAATACTATAA  
 AACTGATTAAAGGCACATTAATATAGCCAGAATAATTATTCTGATTATATCATTTTGAAAG  
 ACCAGATAATCAATATAACTGATTTTTATCCCGAGTAGCTCAGACTTGCAAGCATGCACT  
 ACCATACTTGGATAATTT

The following amino acid sequence <SEQ ID NO. 110> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 50:

\*YGS AVSP\$KSNLE LQLP\*FPCVV EGAWWEITES WGRLPYCSL GSE\*VS\*DLI VL\*GETPFAW  
FLFSLCWLPC KMLLCSSFVF CHDCGASPAM WKCEEIKPLS FINCPVLGMS LSAV\*NGLIH  
QKFKTSLGNT VRPLYKERI RKRKISVA\*Y \*TEYYKTD\*G TLI\*PE\*LF\* LYHFERPDNQ  
 YN\*FLSRVAQ TCKHALPYLD N

The following DNA sequence nGPCR-Seq63 <SEQ ID NO. 51> was identified in *H. sapiens*:

CAGTGAGCCG AGATGGTGCC ATTGCACTCT AGCCTGGGGC AACAGAGCCA GACTCCATCT  
 CCAAAAAAAAA AAGGCCATTC TGAGGATCAA GGCACCACTA GCAACAGGGA GCCCATATGGG  
 TCTCAGACCC TCTCCCCACA TCTCTGGTC CCTGCCCCA CCTGGCGTAC AGGGACCAGC  
 CCCACGGAAG GCTCTTGAGG CCAGGTAACC ATGGGGAGGG GAGGAATGGG GACACCTTCC  
 TCCTGAGTGT CTTAGGGAAG AGAAGCTTAG GTCAGGTGGC TGAGGGTGGA AATGAGAGAG  
 GGGTCTCTC CTGGAGGGTC TCACCATTCC CTTGGTCACC CACCCAATC TCATCTCCCC  
 TGATGTGGGG AGGAGCAGGG GGCATGGATT CCTGAGCCCC AGACTCAACT GTTGTGGTTT  
 ACAGGGGCAT CAGGAGAGAG AGCGAGCAGA ACACACTCCT GCAGCATCCC CTGGCCCCC  
 GCCCCATGAT GGAGCCCAGA GAAGCTGGAC AGCACGTGGG GGCCGCCAAC GGCGCCCAGG  
 AGGATGTGGC CTTCAACCTC ATCATCTGT CCCTACCGA GGGGCTCGGC CTCGGTGGGC  
 TGCTGGGGAA TGGGGCAGTC CTCTGGCTGC TCAGCTCAA TGTCTACAGA AACCCCTTCG  
 CCATCTACCT CCTGGACGTG GCCTGCGCG ATCTCATCTT CCTTGGCTGC CACATGGTGG  
 CCATCGTCCC CGACTTGCTG CAAGGCCGGC TGGACTTCCC GGGCTTCGTG CAGACCAGCC  
 TGGCAACGCT GCGTTCTTC TGCTACATCG TGGGCTGAG TCTCTGGCG GCCGTCAGCG  
 TGGAGCAGTG CCTGGCCGCC CTCTTCCAG CCTGGTACTC GTGCCGCCG CCACGCCACC  
 TGACCACCTG TGTGTGCGCC CTCACCTGGG CCCTCTGCCT GCTGCTGCAC CTGCTGCTCA  
 GCGGCGCCTG CACCCAGTTC TTCGGGGAGC CCAGCCGCCA CTTGTGCCGG ACGCTGTGGC  
 TGGTGGCAGC GGTGCTGCTG GCTCTGCTGT GTTGACCAT GTGTGGGGCC AGCCTTATGC  
 TGCTGCTGCG GGTGGAGCGA GGCCCCCAGC GGCCCCACC CCGGGGCTTC CCTGGGCTCA  
 TCCTCTCAC CGTCTCTC TTCCTCTTCT GCGGCTGCC CTTCCGGCATC TACTGGCTGT  
 CCCGGAACCT GCTCTGGTAC ATCCCCACT ACTTCTACCA CTTAGCTTC CTCATGGCCG  
 CCGTGCATG CGCGGCCAAG CCCGTCGTCT ACTTCTGCCT GGGCAGTGCC CAGGGCCGCA  
 GGCTGCCCC CTGGCTGGTC CTCCAGCGAG CGCTGGGAGA CGAGGCTGAG CTGGGGGCCG  
 TCAGGGAGAC TCCCCGCCG GGCCTGGTGG ACATAGCAGC CTGAGCCCTG GGGCCCCGA  
 CCCCAGCTGC AGCCCCGTG AGGCAAGAGG GTGACGTGGG GAAGGTGGTG GGTGAGAGG  
 CTGGGGCCAG CCGGACCTGG AGGAGGCCTT GGTGGGTGAC CCGGTCATGT GCTGTCAAAG  
 TTGTGACCCT TGGTCTGGAG CATGAGGCTC CCCTGGGAGG CAGCTGGAAA GG

The following amino acid sequence <SEQ ID NO. 111> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 51:

MMEPREAGQHVGAANGAQEDVAFNLHLSTGLGLGGLLNGAVLWLLS  
 SNVYRNPFAIYLLDVACADLIFLGCHMVAIVPDLLQGRLDFFGQVQTSLA  
 TLRFCYIVGLSLLAAVSVEQCLAAFLPAWYSCRRPRHLTTCVCALTWALC  
 LLLHLLSGACTQFFGEPSRHLCRTLWLVAAVLLALLCCTMCGASLMLLL  
 RVERGPQRPPPRGFPLILLTVLLFLFCGLPFGIYWLSRNLLWYIPHYFY  
 HFSFLMAAVHCAAQPVVYFCLGSAQGRRLPLRLVLQRALGDEAELGAVRE  
 TSRRGLVDIAA

The following DNA sequence nGPCR-Seq70 <SEQ ID NO. 52> was identified in *H. sapiens*:

ATGACGTCCACCTGCACCAACAGCACGCGGAGAGTAACAGCAGCCACACGTGCATGCC  
 CTCTCCAAAATGCCATCAGCCTGGCCACGGCATCATCCGCTCAACCGTGCTGGTTATC  
 TTCCTCGCCGCTCTTTCGTCGGCAACATAGTGCTGGCGTAGTGTTCAGCGCAAGCCG  
 CAGCTGCTGCAGGTGACCAACCGTTTTATCTTTAACTCCTCGTACCGACCTGCTGCAG  
 ATTTGCTCGTGGCCCCCTGGGTGGTGGCCACCTCTGTGCCTCTTCTGGCCCCCTAAC  
 AGCCACTTCTGCACGGCCCTGGTTAGCCTCACCCACCTGTTGCGCTTCGCCAGCGTCAAC  
 ACCATTGTCGTGGTGTGAGTGGATCGCTACTTGTCCATCATCCACCCTCTCTCCTACCCG  
 TCCAAGATGACCCAGCGCCGCGGTTACCTGCTCCTCTATGGCACCTGGATTGTGGCCATC  
 CTGCAGAGCACTCCTCACTCTACGGCTGGGGCCAGGCTGCCTTTGATGAGCGCAATGCT  
 CTCTGCTCCATGATCTGGGGGGCCAGCCCCAGCTACACTATTCTCAGCGTGGTGTCTTC  
 ATCGTCAATCCACTGATTGTGATGATTGCTGCTACTCCGTGGTGTCTGTGACGCCCG  
 AGGCAGCATGCTCTGCTGTACAATGTCAAGAGACACAGCTTGAAGTGCGAGTCAAGGAC  
 TGTGTGGAQAATGAGGATGAAGAGGGAGCAGAGAAGAAGGAGGAGTTCAGGATGAGAGT  
 GAGTTTCGCCGCCAGCATGAAGGTGAGGTCAAGGCCAAGGAGGGCAGAATGGAAGCCAAG  
 GACGGCAGCCTGAAGGCCAAGGAAGGAAGCACGGGGACCAAGTGAAGAGTAGTGTAGAGGCC  
 AGGGGCAGCGAGGAGGTGAGAGAGAGCAGCACGGTGGCCAGCGACGGCAGCATGGAGGGT  
 AAGGAAGGCAGCACCAAAGTTGAGGAGAACAGCATGAAGGCAGACAAGGGTGCACAGAG

GTCAACCAGTGCAGCATTGACTTGGGTGAAGATGACATGGAGTTTGGTGAAGACGACATC  
 AATTCAGTGAGGATGACGTCGAGGCAGTGAACATCCCGGAGAGCCTCCCACCCAGTCGT  
 CGTAACAGCAACAGCAACCCCTCCTCTGCCCAGGTGCTACCAGTGCAAAGCTGCTAAAGTG  
 ATCTTCATCATCATTTTCTCCTATGTGCTATCCCTGGGGCCCTACTGCTTTTAGCAGTC  
 CTGGCCGTGTGGGTGGATGTGCAAACCCAGGTACCCAGTGGGTGATCACCATAATCATC  
 TGGCTTTTCTTCTGCACTGCTGCATCCACCCCTATGTCTATGGGTACATGCACAAGACC  
 ATTAAGAAGGAAATCCAGGACATGCTGAAGAAGTTCTTCTGCAAGGAAAAGCCCCGAAA  
 GAAGATAGCCACCCAGACCTGCCCCGAACAGAGGGTGGGACTGAAGGCAAGATTGTCCCT  
 TCCTACGATTCTGCTACTTTTCCTTGA

The following amino acid sequence <SEQ ID NO. 112> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 52:

MTSTCTNSTRESNSSHTCMPLSKMPISLAHGIIRSTVLVIFLAASFVGNIVLALVLQRKP  
 QLLQVTNRIFNLLVTDLLQISLVAPWVATSVPLFWPLNSHFCTALVSLTHLFAFASVN  
 TIVVVSVDRLSYPSKMTQRRGYLLLYGTWIVAILQSTPPL YGWGQA AFDERNA  
 LCSMIWGA SPSTILSVVSFIVPLIVMIACYSVVFCAARRQHALLYNVKRHSLEVRVKD  
 CVENEDEEGAEKKEEFQDESEFRQHEGEVKAKEGRMEAKDGLKAKEGSTGTSESSVEA  
 RGSEEVRESSTVASDGSMEGKEGSTKVEENSMKADKGRTEVNQCSIDLGEDDMEFGEDDI  
 NFSEDDVEAVNIPESLPPSRNSNSNPPLPRCYQCKAAKVIFHFSYVLSLGPYCFLAV  
 LAVWVDVETQVPQWVITIIWLFLLQCCIHYPVYGYMHKTIKKEIQDMLKKFFCKEKPPK  
 EDSHPDLPGTEGGTEGKIVPSYDSATFP

The following DNA sequence nGPCR-Seq63 <SEQ ID NO. 53> was identified in *H. sapiens*:

ATGAT GGAGCCCAGA GAAGCTGGAC AGCACGTGGG GGCCGCCAAC AGCGCCCAGG  
 AGGATGTGGC CTTCAACCTC ATCATCCTGT CCCTCACCGA GGGGCTCGGC CTCGGTGGGC  
 TGCTGGGGAA TGGGGCAGTC CTCTGGCTGC TCAGCTCAA TGTCTACAGA AACCCCTTCG  
 CCATCTACCT CCTGGACGTG GCCTGCGCGG ATCTCATCTT CCTTGGCTGC CACATGGTGG  
 CCATCGTCCC CGACTTGCTG CAAGGCCGGC TGGACTTCCC GGGCTTCGTG CAGACCAGCC  
 TGGCAACGCT GCGCTTCTC TGCTACATCG TGGGCCTGAG TCTCCTGGCG GCCGTCAGCG  
 TGGAGCAGTG CTTGGCCGCC CTCTTCCAG CCTGGTACTC GTGCCGCCGC CCACGCCACC  
 TGACCACCTG TGTGTGCGCC CTCACCTGGG CCCTCTGCCT GCTGCTGCAC CTGCTGTCA  
 GCGGCGCCTG CACCCAGTTC TTCGGGGAGC CCAGCCGCCA CTTGTGCCCG ACGCTGTGGC  
 TGGTGGCAGC GGTGCTGCTG GCTCTGCTGT GTTGACCAT GTGTGGGGCC AGCCTTATGC  
 TGCTGTGCG GGTGGAGCGA GGCCCCCAGC GGCCCCCACC CCGGGGCTTC CCTGGGCTCA  
 TCTCTCTCAC CGTCTCTCTC TTCCTCTTCT GCGGCCTGCC CTTCCGGCATC TACTGGCTGT  
 CCCGGAACCT GCTCTGGTAC ATCCCCCACT ACTTCTACCA CTTAGCTTC CTCATGGCCG  
 CCGTGCACTG CGCGGCCAAG CCCGCTGTCT ACTTCTGCCT GGGCAGTGCC CAGGGCCGCA  
 GGCTGCCCT CCGGCTGGTC CTCCAGCGAG CGCTGGGAGA CGAGGCTGAG CTGGGGGCCG  
 TCAGGGAGAC CTCCCGCCGG GGCCTGGTGG ACATAGCAGC CTGA

The following amino acid sequence <SEQ ID NO. 113> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 53:

MMEPREAGQHVGAANSAQEDVAFNLILSLTEGLGLGGLLGN GAVLWLLSSNVYR  
 NPFAIYLLDVACADLIFLGCHMVAIVPDLQGRLD FPGFVQTSLATLRF CYIVGL  
 SLLAAVSVEQCLAA LFP AWYSCRPRHLTTCVCALTWALCLLHLLSGACTQFF  
 GEPSRHL CRTLWLVA AVLALLCCTMCGASLMLLR VERGPQRPPPRGFPGLILL  
 TVLLFLFCGLPFGIY WLSRNLLWYIPHYFYHFSFLMAA VHC AKPVVYFCLGSAQ  
 GRRPLRLVLQRALGDEAELGAVRETSRRGLVDIAA

The following DNA sequence nGPCR-Seq42 <SEQ ID NO. 54> was identified in *H. sapiens*:

GGAGCCTCGC CGGCCGCCAT GTGGAGCTGC AGCTGGTTCA ACGGCACAGG GCTGGTGGAG  
 GAGCTGCCCTG CCTGCCAGGA CCTGCAGCTG GGGCTGTAC TGTGTGCTGCT GCTGGGCCTG  
 GTGGTGGGCG TGCCAGTGGG CTTGTGCTAC AACGCCCTGC TGGTGTGGC CAACCTACAC  
 AGCAAGGCCA GCATGACCAT GCCGGACGTG TACTTTGTCA ACATGGCAGT GGCAGGCCTG  
 GTGCTCAGCG CCTGGCCCC TGTGCACCTG CTCGGCCCCC CGAGCTCCCG GTGGGCGCTG  
 TGGAGTGTGG GCGGCGAAGT CCACGTGGCA CTGCAGATCC CCTTCAATGT GTCCTCACTG

GTGGCCATGT ACTCCACCGC CCTGCTGAGC CTCGACCACT ACATCGAGCG TGCACTGCCG  
 CGGACCTACA TGGCCAGCGT GTACAACACG CGGCACGTGT GCGGCTTCGT GTGGGGTGGC  
 GCGCTGCTGA CCAGCTTCTC CTCGCTGCTC TTCTACATCT GCAGCCATGT GTCCACCCGC  
 GCGTAGAGT GCGCCAAGAT GCAGAACGCA GAAGCTGCCG ACGCCACGCT GGTGTTTCATC  
 GGCTACGTGG TGCCAGCACT GGCCACCCTC TACGCGCTGG TGCTACTCTC CCGCGTCCGC  
 AGGGAGGACA CGCCCCTGGA CCGGGACACG GGCCGGCTGG AGCCCTCGGC ACACAGGCTG  
 CTGGTGGCCA CCGTGTGCAC GCAGTTTGGG CTCTGGACGC CACACTATCT GATCCTGCTG  
 GGGCACACGG TCATCATCTC GCGAGGGAAG CCGTGGACG CACACTACCT GGGGCTACTG  
 CACTTTGTGA AGGATTTCTC CAAACTCCTG GCCTTCTCCA GCAGCTTTGT GACACCATT  
 CTCTACCGCT ACATGAACCA GAGCTTCCCC AGCAAGCTCC AACGGCTGAT GAAAAAGCTG  
 CCCTGCGGGG ACCGGCACTG CTCCCCGGAC CACATGGGGG TGCAGCAGGT GCTGGCGTAG  
 GCGGCCAGC CCTCCTGGGG AGACGTGACT CTGGTGGACG CAGAGCACTT AGTTACCCTG  
 GACGCTCCCC ACATCCTTCC AGAAGGAGAC GAGCTGCTGG AAGAGAAGCA GGAGGGGTGT  
 TTTTCTTGAA GTTTCCTTTT TCCACAAAT GCCACTCTTG GGCCAAGGCT GTGGTCCCCG  
 TGGCTGGCAT CTGGCTTGAG TCTCCCCGAG GCCTGTGCGT CTCCCAAACA CGCAGCTCAA  
 GGTCCACATC CGCAAAAGCC TCCTCGCCTT CAGCCTCCTC AGCATTCACT TGTCAATGA  
 AGTGATGAAA GCTTAGAGCC AGTATTTATA CTTTGTGGTT AAAATACTTG ATTCCCCCTT  
 GTTTGTTTTA CAAAAACAGA TGTTTCCTAG AAAAATGACA AATAGTAAAA TGAACAAAC  
 CCTACGAAAG AATGGCAACA GCCAGGGTGG CCGGGCCCTG CCAGTGGGCG GCGTGTGCTA  
 GCAAGGCCTG CCGGGTGTGC CGCAGTCACC ACAGGGTTCT GAGAACATTT CACAGAAGTG  
 CCTGAGACGC GGAGACATGG CTGGTGTAA ATGGAGCTAT TCAATAGCAG TGACGCGCTC  
 TCCTCAGCCA CCAAATGTCC CTGACACCCT CCCAGCCCC CACAGATAAC ATCAGCTGAG  
 GTTTTTTTCA GTATGAACCT GTCCTAAATC AATTCCTCAA AGTGTGCACA AAATAAAGA  
 ATATAAATAA ACAAAGAAA GGCAAAAAA AAAAAAA

The following amino acid sequence <SEQ ID NO. 114> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 54:

MWSCSWFNGTGLVEELPACQDLQLGLSLLGLLVGVVGLCYNALLVL  
 ANLHKSASMTMPDVYFVNMAVAGLVLSALAPVHLLGPPSSRWALWSVGGE  
 VHVALQIPFNVSSLVAMYSTALLSLDHYIERALPRTYMASVYNTRHVCGF  
 VWGGALLTSFSSLLFYICSHVSTRALECAKMQNAEADATLVFIGYVVP  
 LATLYALVLLSRVRREDTPLDRDTGRLEPSAHRLLVATVCTQFGLWTPHY  
 LILLGHTVVISRGKPVDAHYLGLLHFVKDFSKLLAFSSSFVTPLLYRYMN  
 QSFPSKLQRLMKKLPCGDRHCSPDHMGVQQLA

The following DNA sequence nGPCR-Seq46 <SEQ ID NO. 55> was identified in *H. sapiens*:

ATGGGGAA CGATTCTGTC AGCTACGAGT ATGGGGATTA CAGCGACCTC TCGGACCGCC  
 CTGTGGACTG CCTGGATGGC GCCTGCCTGG CCATCGACCC GCTGCGCGTG GCCCGCTCC  
 CACTGTATGC CGCCATCTTC CTGGTGGGGG TGCCGGGCAA TGCCATGGTG GCCTGGGTGG  
 CTGGGAAGGT GGCCCGCCGG AGGGTGGGTG CCACCTGGTT GCTCCACCTG GCGTGGCGG  
 ATTTGCTGTG CTGTTTGTCT CTGCCATCC TGGCAGTGCC CATTGCCCGT GGAGGCCACT  
 GGCCGTATGG TGCAGTGGGC TGTCGGGCGC TGCCCTCCAT CATCCTGCTG ACCATGTATG  
 CCAGCGTCCT GCTCCTGGCA GCTCTCAGTG CCGACCTCTG CTCCTGGCT CTCGGGCCTG  
 CCTGGTGGT TACGGTTCAG CGGGCGTGCG GGGTGCAGGT GGCCTGTGGG GCAGCCTGGA  
 CACTGGCCTT GCTGCTCACC GTGCCCTCCG CCATCTACCG CCGGCTGCAC CAGGAGCACT  
 TCCAGCCCG GCTGCAGTGT GTGGTGGACT ACGGCGGCTC CTCCAGCACC GAGAATGCGG  
 TGACTGCCAT CCGGTTTCTT TTTGGCTTCC TGGGCCCCCT GGTGGCCGTG GCCAGCTGCC  
 ACAGTGCCCT CCTGTGCTGG GCAGCCCGAC GCTGCCGGCC GCTGGGCACA GCCATTGTGG  
 TGGGGTTTTT TGTCTGCTGG GCACCCTACC ACCTGCTGGG GCTGGTGCTC ACTGTGGCGG  
 CCCCAGACTC CGCACTCCTG GCCAGGGCCC TGCGGGCTGA ACCCCTCATC GTGGGCCTTG  
 CCCTCGCTCA CAGCTGCCTC AATCCCATGC TCTTCTGTA TTTTGGGAGG GCTCAACTCC  
 GCCGGTCACT GCCAGCTGCC TGTCAGTGGG CCCTGAGGGA GTCCAGGGC CAGGACGAAA  
 GTGTGGACAG CAAGAAATCC ACCAGCCATG ACCTGGTCTC GGAGATGGAG GTGTAG

The following amino acid sequence <SEQ ID NO. 115> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 55:

MGNDSVSYEYGDYSDLSDRPVDCLDGACLAIDPLRVAPLPLYAAIFLVGV  
 PGNAMVAWVAGKVARRRVGATWLLHLAVADLLCCLSLPILAVPIARGGHV



PYGAVGCRALPSIILLTMYASVLLLAALSADLCFLALGPAWWSTVQRACG  
 VQVACGAAWTLALLLTVPSTAIYRRLHQEHFARLQCVVDYGGSSSTENAV  
 TAIRFLGFLGPLVAVASCHSALLCWAARRCRPLGTAIIVGFFVCWAPYH  
 LLGLVLTVAAPNSALLARALRAEPLIVGLALAHSLNPMFLFLYFGRAQLR  
 RSLPAACHWALRESQGQDESVDSSKSTSHDLVSEMEV

The following DNA sequence nGPCR-Seq48 <SEQ ID NO. 56> was identified in *H. sapiens*:

ATGGCACCT TCTCATCGGG CATCACAGGT GGGGTTTTGC CCCACCCCTG AACGCCCTCT  
 GTGGCGCCTT CCACCCACCT GTAGGCCAG AAGGATGTCG GTCTGCTACC GTCCCCCAGG  
 GAACGAGACA CTGCTGAGCT GGAAGACTTC GCGGGCCACA GGCACAGCCT TCCTGCTGCT  
 GGGCGGCGTG CTGGGGCTGC CTGGCAACGG CTTCGTGGTG TGGAGCTTGG CGGGCTGGCG  
 GCCTGCACGG GGGCGACCGC TGGCGGCCAC GCTTGTGCTG CACCTGGCGC TGGCCGACGG  
 CGCGGTGCTG CTGCTCACGC CGCTCTTTGT GGCCTTCCTG ACCCGGCAGG CCTGGCCGCT  
 GGGCCAGGCG GGCTGCAAGG CGGTGTACTA CGTGTGCGCG CTCAGCATGT ACGCCAGCGT  
 GCTGCTCACC GGCCTGCTCA GCCTGCAGCG CTGCCTCGCA GTCACCCGCC CCTTCCTGGC  
 GCCTCGGCTG CGCAGCCCGG CCCTGGCCCG CCGCCTGCTG CTGGCGGTCT GGCTGGCCGC  
 CCTGTTGCTC GCCGTCCCGG CCGCGTCTA CCGCCACCTG TGGAGGGACC GCGTATGCCA  
 GCTGTGCCAC CCGTCGCCGG TCCACGCCGC CGCCACCTG AGCCTGGAGA CTCTGACCGC  
 TTTCGTGCTT CCTTTCGGGC TGATGCTCGG CTGCTACAGC GTGACGCTGG CACGGCTGCG  
 GGGCGCCCGC TGGGGCTCCG GGCGGCACGG GGCGCGGGTG GGCCGGCTGG TGAGCGCCAT  
 CGTGCTTGCC TTCGGCTTGC TCTGGGCCCC CTACCACGCA GTCAACCTTC TGCAGGCGGT  
 CGCAGCGCTG GCTCCACCGG AAGGGGCCCT GGCGAAGCTG GGCGGAGCCG GCCAGGCGGC  
 GCGAGCGGGA ACTACGGCCT TGGCCTTCTT CAGTTCTAGC GTCAACCCGG TGCTCTACGT  
 CTTACCCGCT GGAGATCTGC TGCCCCGGGC AGGTCCCCGT TTCCTCACGC GGCTCTTCGA  
 AGGCTCTGGG GAGGCCCGAG GGGCGGCCG CTCTAGGGAA GGGACCATGG AGCTCCGAAC  
 TACCCCTCAG CTGAAAGTGG TGGGGCAGG CCGCGCAAT GGAGACCCGG GGGGTGGGAT  
 GGAGAAGGAC GGTCCGGAAT GGGACCTTTG A

The following amino acid sequence <SEQ ID NO. 116> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 56:

MAPSHRASQVGFCTPERPLWRLPPTCRPRRMSVCYRPPGNETLLSWKTS  
 RATGTAFLLLAALLGLPGNGFVWVSLAGWRPARGRPLAATLVHLALADG  
 AVLLLTPLFVAFLTRQAWPLGQAGCKAVYYVCALSMYASVLLTGLLSLQR  
 CLAVTRPFLAPRLRSPALARRLLAVWLAALLAVPAAVYRHLWRDRVQ  
 LCHPSPVHAAHLSLETLTAFVLPFGLMLGCYSVTLARLRGARWGSGRHG  
 ARVGRLVSAIVLAFGLLWAPYHAVNLLQAVAALAPPEGALAKLGGAGQAA  
 RAGTTALAFFSSSVNPVLYVFTAGDLLPRAGPRFLTRLFEGSGEARGGGR  
 SREGTMELRTPQLKVVGQGRGNGDPGGGMEKDGPEWDL

The following DNA sequence nGPCR-Seq51 <SEQ ID NO. 57> was identified in *H. sapiens*:

CCGCCGCGCG GGGAGGACGC GAGCACCCAG CTTTAATCCC TGGAAAGTCC ACGAACAATG  
 AATCCATTTT ATGCATCTTG TTGGAACACC TCTGCCGAAC TTTTAAACAA ATCCTGGAAT  
 AAAGAGTTTG CTTATCAAAC TGCCAGTGTG GTAGATACAG TCATCCTCCC TTCCATGATT  
 GGGATTATCT GTTCAACAGG GCTGGTTGGC AACATCCTCA TTGTATTAC TATAATAAGA  
 TCCAGGAAAA AAACAGTCCC TGACATCTAT ATCTGCAACC TGGCTGTGGC TGATTTGGTC  
 CACATAGTTG GAATGCCTTT TCTTATTAC CAATGGGCCC GAGGGGGAGA GTGGGTGTTT  
 GGGGGGCCCT TCTGCACCAT CATCACATCC CTGGATACTT GTAACCAATT TGCCTGTAGT  
 GCCATCATGA CTGTAATGAG TGTGGACAGG TACTTTGCC TCGTCCAACC ATTTGCACTG  
 ACACGTTGGA GAACAAGGTA CAAGACCATC CGGATCAATT TGGGCCTTTG GGCAGCTTCC  
 TTTATCCTGG CATTGCCTGT CTGGGTCTAC TCGAAGGTCA TCAAATTTAA AGACGGTGTT  
 GAGAGTTGTG CTTTTGATTT GACATCCCCT GACGATGTAC TCTGGTATAC ACTTTATTTG  
 ACGATAACAA CTTTTTTTTT CCTCTACCC TTGATTTTGG TGTGCTATAT TTTAATTTA  
 TGCTATACTT GGGAGATGTA TCAACAGAAT AAGGATGCCA GATGCTGCAA TCCAGTGTA  
 CCAAAACAGA GAGTGATGAA GTTGACAAAG ATGGTGCTGG TGCTGGTGGT AGTCTTTATC  
 CTGAGTGCTG CCCCTTATCA TGTGATACAA CTGGTGAAC TACAGATGGA ACAGCCCACA  
 CTGGCCTTCT ATGTGGGTGA TTACCTCTCC ATCTGTCTCA GCTATGCCAG CAGCAGCATT  
 AACCTTTTCT TCTACATCCT GCTGAGTGGA AATTTCCAGA AACGTCTGCC TCAAATCCAA

AGAAGAGCGA CTGAGAAGGA AATCAACAAT ATGGGAAACA CTCTGAAATC ACACTTTTAG  
 GAAAGTACAT GGATCACCAT GAGTCTAGAC ATGATTGTCT ATCTTACTGG TATTATTAGA  
 AAGGGCAGGT GTACCGATAT GTTATGCCC ATTCTTCTTG TGTACTTGTG ACTCTTAGCA  
 GCATGGAAGA GAAGTGTAAC CATGCAAATA CAATGAGCTT AATATGCTAA CTTTAGCAAG  
 ATGTAAAATG TTGATCTATA TTGTGGGTAG GGAATGGGAT AGTCTGAGAT ACCCAGGCTT  
 CATGATGGTG TATATTATTT CAGCATATTA TAACTAGTC ACTAATGAAA ATGGCCATCC  
 ATGACCATTG ACTCAAACT CACCAAGGAA CCTGACCTTG CCCTCCACAC TGTGGCCTCA  
 CTGTAACAGT TTCCTCAAGG TTCCTAGGAG GGTATCACCT TAGAGTGAAG TCTAAAATTT  
 GGCTATTTTT TATCTATAAA AAATGTCAGT TTTATATGGT CCAATACTAA TACCCTCAAC  
 AACTAAGCCC CACCTTTTAG AATAAGTTAC CATTTATTGC ACACATGCAA TGTGTAAGAT  
 TACATGTAAC AAACCTGTGA AATAAGTATT ATTACCTTG TTTGCTAAGG CTCAGAAAGG  
 AGAAATGATA GGCCTAATGC TGCAACAGCT ATCTAAGAGC TGAGCTAACA TTCAGCTCTG  
 CCTGTTTCTT TTCTACTGCC GACCTTGACA ACCTTTACTT ATCATACTGG AGAACCCAGT  
 AACTTGGAGT TTCTTTTGCT TTCTCCTGTA GCCCTACAAG AGGAGAACTA AAGTCTGATA  
 GAAATGAGTT GATGTTTTAA GCATCATTTT GGATTATCTT GTTCTCACAC CTGCTAACTG  
 TAGAAACTGG CATCTGGACT TTAATAATAA TACTTTACTT CTGGA

The following amino acid sequence <SEQ ID NO. 117> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 57:

MNPFHASCWNTSAELLNKSWNKEFAYQTASVVDTVILPSMIGIICSTGLVGNILIVFTIIRSRRKKTVP  
 DIYICNLAVADLV HIVGMPFLIHQWARGGEWVFGGPLCTHTSLDTCNQFACSAIMTVMSVDR  
 YFALVQPFRLTRWRTRYKTIRINLGLWAASFILALPVWVYSKVIKFKDGV  
 ESCAFDLTSPDDVLWYTLYLITITFFPLPLILVCYILLCYTWEMYQQN  
 KDARCCNPSPVKQRVMKLTKMVLVLVVVFILSAAPYHVIQLVNLQMEQPT  
 LAFYVGYLSICLSYASSINPFLYILLSGNFQKRLPQIQRRATEKEINN  
 MGNTLKSHF

The following DNA sequence nGPCR-Seq52 <SEQ ID NO. 58> was identified in *H. sapiens*:

CCATGACCAG CAATTTTTCC CAACCTGTTG TGCAGCTTTG CTATGAGGAT GTGAATGGAT  
 CTGTGATTGA AACTCCCTAT TCTCCTGGGT CCCGGGTAAT TCTGTACACG GCGTTTAGCT  
 TTGGGTCITT GCTGGCTGTA TTTGGAATC TCTTAGTAAT GACTTCTGTT CTTCATTTTA  
 AGCAGCTGCA CTCTCCAACC AATTTTCTCA TTGCCTCTCT GGCCTGTGCT GACTTCTTGG  
 TAGGTGTGAC TGTGATGCTT TTCAGCATGG TCAGGACGGT GGAGAGCTGC TGGTATTTTG  
 GAGCCAAATT TTGTACTCTT CACAGTTGCT GTGATGTGGC ATTTTGTTAC TCTTCTGTCC  
 TCCACTTGTG CTTCATCTGC ATCGACAGGT ACATTGTGGT TACTGATCCC CTGGTCTATG  
 CTACCAAGTT CACCGTGTCT GTGTCGGGAA TTTGCATCAG CGTGTCTGCT ATTCTGCCTC  
 TCACGTACAG CGGTGCTGTG TTCTACACAG GTGTCAATGA TGATGGGCTG GAGGAATTAG  
 TAAGTGCTCT CAACTGCGTA GGTGGCTGTC AAATTATTGT AAGTCAAGGC TGGGTGTTGA  
 TAGATTTTCT GTTATTCTTC ATACCTACCC TTGTTATGAT AATTCTTTAC AGTAAGATTT  
 TTCTTATAGC TAAACAACAA GCTATAAAAA TTGAACTAC TAGTAGCAAA GTAGAATCAT  
 CCTCAGAGAG TTATAAAATC AGAGTGGCCA AGAGAGAGAG GAAAGCAGCT AAAACCCCTGG  
 GGGTCACGGT ACTAGCATTT GTTATTTTAT GGTACCGTA TACAGTTGAT ATATTAATTG  
 ATGCCTTTAT GGGCTTCCTG ACCCCTGCCT ATATCTATGA AATTTGCTGT TGGAGTGCTT  
 ATTATAACTC AGCCATGAAT CCTTTGATTT ATGCTCTATT TTATCCTTGG TTTAGGAAAG  
 CCATAAACT TATTTAAAGT GGAGATGTTT TAAAGGCTAG TTCATCAACC ATTAGTTTAT  
 TTTTAGAATA A

The following amino acid sequence <SEQ ID NO. 118> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 58:

MTSNFSQP VQLCYEDVNG SCIETPYSPG SRVILYTAFS FGSLAVFGN LLVMTSVLHF  
 KQLHSPTNFL IASLACADFL VGVTVMLFSM VRTVESCWYF GAKFCTLHSC CDVAFCYSSV  
 LHLFCICIDR YIVVTDPLVY ATKFTVSVSG ICISVSWILP LTYSGAVFYT  
 GVNDGLEEL VSALNCVGGC QIVSQGWVL IDFLFFIPT LVMILYSKI  
 FLIAKQQA K IETTSSKVES SSES YKIRVA KRERKAAKTL GVTVLAFVIS  
 WLPYTVDILI DAFMGFLTPA YIYEICWWSA YYN SAMNPLI YALFYPWFRK  
 AIKLILSGDV LKASSSTISL FLE

The following DNA sequence nGPCR-Seq49 <SEQ ID NO. 59> was identified in *H. sapiens*:

A TGCCACTCCC TGTGCCCCCA GCGGGGGCAC AAAAGACTCC TGAGGACCAT GTCTGTCTCC  
ATCTTGACAGG TCCCTCGCCA GCACCTCTG AGCCGGCCAG GATGTTTGGG CTGTTGGGCC  
TCTGGAGAAC CTTGACAGT GTGGTCTTCT ACCTGACGCT GATCGTGGGC CTCGGGGGAC  
CGGTAGGTAA CGGGCTGGTG CTCTGGAACC TCGGCTTCCG CATCAAGAAG GGCCCTTCT  
CCATCTACCT GCTGCACCTG GCCGCCGCCG ACTTCCTGTT CCTCTCCTGC CGTGTGGGCT  
TCTCCGTGGC TCAGGCTGCC CTGGGCGCCC AGGACACACT CTAATCGTG CTCACCTTCC  
TGTGGTTCGC GGTGGGGCTC TGGCTGCTGG CGGCCTTCAG CGTGGAGCGC TGCCTCTCCG  
ACCTCTTCCC CGCCTGCTAC CAGGGCTGCC GGCCAGACA CGCCTCGGCC GTCCTCTGCG  
CCCTGGTGTG GACCCCGACC CTGCCGGCCG TGCCGCTGCC CGCCAACGCC TGCGGCCTGC  
TGCGCAACAG CGCGTGCCCC CTGGTCTGCC CGCGCTACCA CGTGGCCAGC GTCACCTGGT  
TCCTGGTGTG GGCCCGCGTC GCCTGGACGG CTGGCGTGGT CCTCTTTGTC TGGGTGACCT  
GCTGCTCCAC TCGCCCGCGG CCCAGGCTCT ACGGCATCGT CCTGGGCGCG CTGCTCCTGC  
TCTTCTTCTG TGGCCTGCCC TCGGTCTTCT ACTGGAGCCT GCAGCCCCTG CTGAACCTCC  
TGCTGCCCCG GTTTTCCCCG CTGGCCACGC TGCTGGCCTG CGTCAACAGC AGCTCCAAGC  
CCCTCATCTA CTCGGGGTTG GGCCGACAGC CCGGGAAGCG GGAGCCGCTG AGGTCCGTAC  
TGCGGAGGGC CTTGGGGGAG GCGCCGAGC TGGGTGCCAG GGGACAGTCC CTGCCATGG  
GTCTCCTATA A

The following amino acid sequence <SEQ ID NO. 119> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 59:

MPLPVPPAGA QKTPEDHVCLHLA GSPAPSEPARMFGLFGLWR  
TFDSVVFYLT LIVGLGGP  
VGNGLVLWNLGFRIKGPFSIYLLHLAAADFLFLSCR  
VGFSAQAALGAQDTLYFVL TFLWFAVG  
LWLLAASFVERCLSDLFPACYQGC  
RPRHASAVLCALVWTPLPAVPLPANACGLLRNSACPLVCP  
RYHVASVTWFLVLARVAWTAGVVLFWVTCCSTRPR  
RLYGIVLGALLLFFCGLPSVF YWSLQP  
LLNLLPVFESPLATLLACVNSSSKPLIYSG  
LRQPGKREPLRSVLRRALGEGAE LGARGQSLPMGLL

The following DNA sequence nGPCR-Seq61 <SEQ ID NO. 60> was identified in *H. sapiens*:

ATGAACAACAATACAACATGTATTCAACCATCTATGATCTCTTCCATGGCTTTACCAATC  
ATTACATCCTCCTTTGTATTGTTGGTGTTTTTGGAAACACTCTCTCTCAATGGATATTT  
TTAACAATAATAGGTAAAAAACATCAACGCACATCTACCTGTACACCTTGTGACTGCA  
AACTTACTTGTGTGACAGTGCCATGCCTTTCATGAGTATCTATTTCTGAAAGGTTTCCA  
TGGGAATATCAATCTGCTCAATGCAGAGTGGTCAATTTTCTGGGAACTCTATCCATGCAT  
GCAAGTATGTTTGTGAGTCTCTTAATTTTAAGTTGGATTGCCATAAGCCGCTATGCTACC  
TTAATGCAAAAGGATTCTCGCAAGAGACTACTTCATGCTATGAGAAAATATTTTATGGC  
CATTTACTGAAAAAATTTTCGCCAGCCCACTTTGCTAGAAAACTATGCATTACATATGG  
GGAGTTGTACTGGGCATAATCATTCCAGTTACCGTATACTACTCAGTCATAGAGGCTACA  
GAAGGAGAAGAGAGCCTATGCTACAATCGGCAGATGGAAGTGGAGCCATGATCTCTCAG  
ATTGCAGGTCTCATTGGAACCATTTAATTGGATTTTCTTTTTAGTAGTACTAACATCA  
TACTACTCTTTTGTAGCCATCTGAGAAAAATAAGAACCTGTACGTCCATTATGGAGAAA  
GATTTGACTTACAGTTCTGTGAAAAGACATCTTTTGGTCATCCAGATTCTACTAATAGTT  
TGCTTCCTTCCTTATAGTATTTTAAACCCATTTTATGTTCTACACCAAAGAGATAAC  
TGTCAGCAATTGAATTATTTAATAGAAACAAAAACATTCTCACCTGTCTTGCTTCGGCC  
AGAAGTAGCACAGACCCCATTAATTTCTTTTATTAGACAAAACATTCAAGAAGACACTA  
TATAATCTCTTTACAAAGTCTAATTCAGCACATATGCAATCATATGGTTGA

The following amino acid sequence <SEQ ID NO. 120> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 60:

MNNNTTCIQPSMISSMALPIYILLCIVGVFGNTLSQWIFLTKIGKKTSTHIYLSHLVT  
ANLLVCSAMPFMSIYFLKGFQWEYQSAQCRVVN  
FLGTLMSHASMFSLLILSWIAISRY  
ATLMQKDSSQETTSCEYKIFYGHLKKFRQPNFARKLCIYTWGVVLGHIIPVTVYYSVI  
EATEGEESLCYNRQME LGAMISQIAGLIGTTFIGFSFLVVLTSYYSFVSHLRKIRTCTS  
IMEKDLTYSSVKRHLVIOILLIVCFLPYSIFKPIFYVLHQRDNCQQLNYLIETKNILT  
CLASARSSTDPIIFLLLDKTFKKTLYNLFTKSNSAHMQSYG

**Example 2: Cloning of nGPCR-x**

cDNAs may be sequenced directly using an ABI377 or ABI373A fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM Ready Dye-Deoxy Terminator kit with Taq FS polymerase. Each ABI cycle  
5 sequencing reaction contains about 0.5 µg of plasmid DNA. Cycle-sequencing is performed using an initial denaturation at 98C for 1 minute, followed by 50 cycles: 98C for 30 seconds, annealing at 50C for 30 seconds, and extension at 60C for 4 minutes. Temperature cycles and times are controlled by a Perkin-Elmer 9600 thermocycler. Extension products are purified using Centriflex gel filtration (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each  
10 reaction product is loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B table top centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples are dried under vacuum for about 40 minutes and then dissolved in 5 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples are then heated to 90C for three minutes and loaded  
15 into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence analysis is done by importing ABI373A files into the Sequencer program (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp are obtained. Potential sequencing errors are minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers at different locations until all sequencing ambiguities are removed.

20 To isolate a cDNA clone encoding full length nGPCR, a DNA fragment corresponding to a nucleotide sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60, or a portion thereof, can be used as a probe for hybridization screening of a phage cDNA library. The DNA fragment is amplified by the polymerase chain reaction (PCR) method. The PCR reaction mixture of 50 µl contains polymerase mixture (0.2 mM dNTPs, 1x PCR Buffer and  
25 0.75 µl Expand High Fidelity Polymerase (Roche Biochemicals)), 1 µg of 3206491 plasmid, and 50 pmoles of forward primer and 50 pmoles of reverse primer. The primers are preferably 10 to 25 nucleotides in length and are determined by procedures well known to those skilled in the art. Amplification is performed in an Applied Biosystems PE2400 thermocycler, using the following program: 95C for 15 seconds, 52C for 30 seconds and 72C for 90 seconds; repeated  
30 for 25 cycles. The amplified product is separated from the plasmid by agarose gel electrophoresis, and purified by Qiaquick gel extraction kit (Qiagen).

A lambda phage library containing cDNAs cloned into lambda ZAPII phage-vector is plated with E. coli XL-1 blue host, on 15 cm LB-agar plates at a density of 50,000 pfu per plate, and grown overnight at 37C; (plated as described by Sambrook et al., supra). Phage plaques are

transferred to nylon membranes (Amersham Hybond NJ), denatured for 2 minutes in denaturation solution (0.5 M NaOH, 1.5 M NaCl), renatured for 5 minutes in renaturation solution (1 M Tris pH 7.5, 1.5 M NaCl), and washed briefly in 2xSSC (20x SSC: 3 M NaCl, 0.3 M Na-citrate). Filter membranes are dried and incubated at 80C for 120 minutes to cross link the phage DNA to the membranes.

The membranes are hybridized with a DNA probe prepared as described above. A DNA fragment (25 ng) is labeled with  $\alpha$ -<sup>32</sup>P-dCTP (NEN) using Rediprime random priming (Amersham Pharmacia Biotech), according to manufacturers instructions. Labeled DNA is separated from unincorporated nucleotides by S200 spin columns (Amersham Pharmacia Biotech), denatured at 95C for 5 minutes and kept on ice. The DNA-containing membranes (above) are pre-hybridised in 50 ml ExpressHyb (Clontech) solution at 68C for 90 minutes. Subsequently, the labeled DNA probe is added to the hybridization solution, and the probe is left to hybridise to the membranes at 68C for 70 minutes. The membranes are washed five times in 2x SSC, 0.1% SDS at 42C for 5 minutes each, and finally washed 30 minutes in 0.1x SSC, 0.2% SDS. Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with an intensifying screen at -80C for 16 hours. One positive colony is isolated from the plates, and replated with about 1000 pfu on a 15 cm LB plate. Plating, plaque lift to filters and hybridization are performed as described above. About four positive phage plaques are isolated from this secondary screening.

cDNA containing plasmids (pBluescript SK-) are rescued from the isolated phages by in vivo excision by culturing XL-1 blue cells co-infected with the isolated phages and with the Excision helper phage, as described by manufacturer (Stratagene). XL-blue cells containing the plasmids are plated on LB plates and grown at 37C for 16 hours. Colonies (18) from each plate are replated on LB plates and grown. One colony from each plate is stricken onto a nylon filter in an ordered array, and the filter is placed on a LB plate to raise the colonies. The filter is then hybridized with a labeled probe as described above. About three positive colonies are selected and grown up in LB medium. Plasmid DNA is isolated from the three clones by Qiagen Midi Kit (Qiagen) according to the manufacturer's instructions. The size of the insert is determined by digesting the plasmid with the restriction enzymes NotI and SalI, which establishes an insert size. The sequence of the entire insert is determined by automated sequencing on both strands of the plasmids.

#### *nGPCR-70*

The PCR reaction was performed in 50  $\mu$ l containing 31  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l Buffer II (PE Applied Biosystems AmpliTaq Gold system), 6  $\mu$ l 25 mM MgCl<sub>2</sub>, 2  $\mu$ l 10 mM dNTP mix, 5  $\mu$ l

Marathon Ready whole human brain cDNA (Clontech #7400-1), 0.3 µl primer VR70C (1 µg/µl)(SEQ ID NO:121), 0.3 µl primer VR70D (1 µg/µl)(SEQ ID NO:122), and 0.4 µl AmpliTaq Gold™ DNA Polymerase. The primer sequence for VR70C is 5'-

TTCAAAGCTTATGACGTCCACCTGCACC-3' (SEQ ID NO:121), corresponding to the 5' end of the coding region and containing a HindIII restriction site. The primer sequence for VR70D is 5'- TTCACTCGAGTCAAGGAAAAGTAGCAGAATCGTAG-3' (SEQ ID NO:122), corresponding to the 3' end of the coding region and containing an XhoI restriction site (Genosys).

The PCR reaction was carried out using a GeneAmp PCR 9700 thermocycler (Perkin Elmer Applied Biosystems) and started with 1 cycle of 80C for 20 minutes followed by 95C for 10 minutes, then 12 cycles at 95C for 30 seconds, 72C for 2 minutes decreasing 1C each cycle, 72C for 1 minute, followed by 35 cycles at 95C for 30 seconds, 60C for 30 seconds, 72C for 1 minute. The PCR reaction was loaded on a 0.75% gel. The DNA band was excised from the gel and the DNA was eluted from the agarose using a QIAquick gel extraction kit (Qiagen). The eluted DNA was ethanol-precipitated and resuspended in 4 µl H<sub>2</sub>O for ligation. The ligation reaction consisted of 4 µl of fresh ethanol-precipitated PCR product and 1 µl of pCRII-TOPO vector (Invitrogen). The reaction was gently mixed and allowed to incubate for 5 minutes at room temperature followed by the addition of 1 µl of 6x TOPO cloning stop solution and mixing for 10 seconds at room temperature. The sample was then placed on ice and 2 µl was transformed in 50 µl of One Shot cells (Invitrogen) and plated onto ampicillin plates. Seven white colonies were chosen and the presence of an insert was verified by PCR in the following manner. Each colony was resuspended in 2 ml LB broth and incubated at 37C for 2 hours. A 500 µl aliquot was spun down in the microfuge, the supernatant discarded, and the pellet resuspended in 25 µl of H<sub>2</sub>O. A 16 µl aliquot was removed and boiled for 5 minutes and the sample was placed on ice. The sample was microfuged briefly to pellet any bacterial debris and PCR was carried out as described above with 15 µl of sample using primers VR70C (SEQ ID NO:121) and VR70D (SEQ ID NO:122).

#### *nGPCR-63*

Isolation of a clone for nGPCR-63 from genomic DNA was performed by PCR in a 50 µl reaction containing Herculase DNA Polymerase blend (Stratagene), with buffer recommendations as supplied by the manufacturer, 200 ng each primers PSK16 and 17 (SEQ ID NO:123 and SEQ ID NO:124, respectively), 150 ng of human genomic DNA (Clontech) and 4% DMSO. The PCR reaction was performed on a Robocycler thermocycler (Stratagene) starting with 1 cycle of 94C for 2 minutes followed by 35 cycles of 94C for 30 seconds, 65C for

30 seconds, 72C for 2 minutes. The PCR reaction was purified by the QiaQuick PCR Purification Kit (Qiagen) and eluted in TE. The PCR primer sequences were: PSK16 5'-GATCGAATTCATGATGGAGCCCAGAGAAGCTGGAC-3' (SEQ ID NO:123) and PSK17 5'-GATCCTCGAGTCAGGCTGCTATGTCCACCAGGCC-3' (SEQ ID NO:124). Translation initiation and termination codons are shown above in bold.

The PCR product was ligated into the pCR-BluntII-TOPO vector (Invitrogen) using the Zero Blunt Topo PCR TA cloning kit as follows. 3 µl PCR product DNA, 1 µl pCRII-TOPO vector, and 1 µl TOPOII salt solution (1.2M NaCl, 0.06M MgCl<sub>2</sub>) was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250 µl of SOC was added, then incubated at 37C with shaking for one hour and then plated onto ampicillin plates supplemented with Xgal and IPTG. Single colonies were screened by restriction digestion for the presence of the insert, and a plasmid DNA from colony 63-4-23 was purified using a Qiagen Endo-Free plasmid purification kit.

The clone containing nGPCR-63 was sequenced directly using an ABI377 fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI BigDye<sup>TM</sup> Terminator Cycle Sequencing Ready Reaction kit with Taq FSTM polymerase. Each ABI cycle sequencing reaction contained about 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98C for 1 minute, followed by 50 cycles: 96C for 30 seconds, annealing at 50C for 30 seconds, and extension at 60C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using AGTC (R) gel filtration block (Edge Biosystems, Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which was then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 3 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90C for 3.5 minutes and loaded into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence analysis was performed by importing ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI).

*nGPCR-42*

PCR was performed in a 50  $\mu$ l reaction containing 36.7  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l 10X TT buffer (140 mM Ammonium Sulfate, 0.1 % gelatin, 0.6 M Tris-tricine pH 8.4), 5  $\mu$ l 15 mM MgSO<sub>4</sub>, 2  $\mu$ l 10 mM dNTP, 0.3  $\mu$ l 6541947H1 DNA (1.4 Tg/Tl), 0.3  $\mu$ l of LW1579 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l of LW1580 (1  $\mu$ g/ $\mu$ l), 0.4  $\mu$ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was performed on a on a Robocycler thermocycler (Stratagene) starting with 1 cycle of 94C for 2 minutes followed by 15 cycles of 94C for 30 seconds, 55C for 30 seconds, 68C for 1.3 minutes. The PCR reaction was loaded onto a 1.2 % agarose gel. The DNA band was excised from the gel, placed in GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microcentrifuge. The eluted DNA was EtOH precipitated and resuspended in 4 $\mu$ l H<sub>2</sub>O for ligation. The forward PCR primer was: LW1579 5'-GCATAAGCTTCCATGTGGAGCTGC AGCTGGTTCAACG-3' (SEQ ID NO:125), and the reverse PCR primer was: LW1580 5'-GCATCTCGAGCCTACGCCAGCACCTGCTGCACC-3' (SEQ ID NO:126). The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen) which consisted of 4 $\mu$ l PCR product DNA and 1  $\mu$ l pCRII-TOPO vector that was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250  $\mu$ l of SOC was added, then incubated at 37C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and then sequenced.

#### *nGPCR-46*

PCR was performed in a 50  $\mu$ l reaction using components that come with PLATINUM<sup>®</sup> Pfx DNA Polymerase (GibcoBRL) containing 30.5  $\mu$ l H<sub>2</sub>O, 5  $\mu$ l 10X Pfx Amplification buffer, 5  $\mu$ l 10X Enhancer solution, 1.5  $\mu$ l 50mM MgSO<sub>4</sub>, 2  $\mu$ l 10 mM dNTP, 5  $\mu$ l human genomic DNA (0.3 $\mu$ g/ $\mu$ l)(Clontech), 0.3  $\mu$ l of LW1626 (1  $\mu$ g/ $\mu$ l), 0.3  $\mu$ l of LW1627 (1  $\mu$ g/ $\mu$ l), 0.4  $\mu$ l PLATINUM<sup>®</sup> Pfx DNA Polymerase (2.5 U/ $\mu$ l). The PCR reaction was performed in a Robocycler Gradient 96 (Stratagene) starting with 1 cycle of 94C for 5 minutes followed by 30 cycles at 94C for 40 seconds, 55C for 2 minutes, 68C for 3 minutes. Following the final cycle, 0.5  $\mu$ l of AmpliTaq DNA Polymerase (5 U/ $\mu$ l) was added and the tube was incubated at 72C for 5 minutes. The PCR reaction was loaded onto a 1.2 % agarose gel. The DNA band was excised from the gel, placed in GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microcentrifuge. The eluted DNA was EtOH precipitated and resuspended



in 12µl H<sub>2</sub>O for ligation. The forward PCR primer was: LW1626 5'-

GCATAAGCTTCCATGGGGAACGATTCTGTCAGC-3' (SEQ ID NO:127) and the reverse

PCR primer was: LW1627 5'-GCATCTCGAGCCTACACCTCCATCTCCGAGACC-3' (SEQ ID NO:128). The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen)

5 which consisted of 4µl PCR product DNA and 1 µl pCRII-TOPO vector that was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250  
10 µl of SOC was added, then incubated at 37C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and then sequenced.

#### *nGPCR-48*

15 PCR was performed in a 50 µl reaction using components that come with PLATINUM<sup>®</sup> Pfx DNA Polymerase (GibcoBRL) containing 30.5 µl H<sub>2</sub>O, 5 µl 10X Pfx Amplification buffer, 5 µl 10X Enhancer solution, 1.5 µl 50mM MgSO<sub>4</sub>, 2 µl 10 mM dNTP, 5 µl human genomic DNA (0.3µg /µl)(Clontech), 0.3 µl of LW1572 (1 µg/µl), 0.3 µl of LW1573 (1 µg/µl), 0.4 µl PLATINUM<sup>®</sup> Pfx DNA Polymerase (2.5 U/µl). The PCR reaction was performed in a  
20 Robocycler Gradient 96 (Stratagene) starting with 1 cycle of 94C for 5 minutes followed by 30 cycles at 94C for 40 seconds, 55C for 2 minutes, 68C for 3 minutes. Following the final cycle, 0.5 µl of AmpliTaq DNA Polymerase (5 U/Tl) was added and the tube was incubated at 72C for 5 minutes. The PCR reaction was loaded onto a 1.2 % agarose gel. The DNA band was excised from the gel, placed in GenElute Agarose spin column (Supelco) and spun for 10 minutes at  
25 maximum speed in a microcentrifuge. The eluted DNA was EtOH precipitated and resuspended in 12µl H<sub>2</sub>O for ligation. The forward primer for PCR was: LW1572 5'-GATCAAGCTTGCA TGGCACCTTCTCATCGGG-3' (SEQ ID NO:129) and the reverse primer was: LW1573 5'-GATCCTCGAGTCAAAGGTCCCATTCCGACC-3' (SEQ ID NO:130). The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen) which consisted of 4µl PCR product  
30 DNA and 1 µl pCRII-TOPO vector that was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250 µl of SOC was added, then incubated

at 37C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and then sequenced. *nGPCR-49*

5        The PCR reaction used components that come with PLATINUM<sup>®</sup> Pfx DNA Polymerase (GibcoBRL) containing 30.5 µl H<sub>2</sub>O, 5 µl 10X Pfx Amplification buffer, 5 µl 10X Enhancer Solution, 1.5 µl 50mM MgSO<sub>4</sub>, 2 µl 10 mM dNTP, 5 µl human genomic DNA (0.3µg/µl) (Clontech), 0.3 µl of LW1726 (1 µg/µl) (SEQ ID NO:131), 0.3 µl of LW1727 (1 µg/µl) (SEQ ID NO:132), 0.4 µl PLATINUM<sup>®</sup> Pfx DNA Polymerase (2.5 U/µl). The PCR reaction was  
10 performed in a Robocycler Gradient 96 (Stratagene) starting with 1 cycle of 94C for 2 minutes followed by 35 cycles at 94C for 30 seconds, 55C for 30 seconds, 68C for 2 minutes. Following the final cycle, 0.5 µl of AmpliTaq DNA Polymerase (5 U/µl) was added and the tube was incubated at 72C for 5 minutes. The sequence of LW1726 is: 5'-CATAAGCTTTGGATGCCA  
CTCCCTGTGCCCCC-3' (SEQ ID NO:131) and for LW1727 is: 5'-GCATCTCGAGTTATAG  
15 GAGACCCATGGGCAGGG-3' (SEQ ID NO:132). The underlined portion of the primer matches the 5' and 3' areas, respectively, of the coding region. The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen) which consisted of 4 µl PCR product DNA, 1 µl of salt solution and 1 µl pCRII-TOPO vector that was incubated for 5 minutes at room temperature then the reaction was placed on ice. Two microliters of the ligation reaction  
20 was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250 µl of SOC was added, then incubated at 37C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and  
25 then sequenced.

      The mutation in SEQ-49 in the plasmid above was repaired using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). The PCR reaction contained 40 µ H<sub>2</sub>O, 5 µl 10x Reaction buffer, 1 µl mini-prep DNA, 1 µl LW1741 (125 ng/µl) (SEQ ID NO:133), 1µl LW1742 (125 ng/µl) (SEQ ID NO:134), 1µl 10 mM dNTP, 1 µl Pfu DNA polymerase. The  
30 cycle conditions were 95C for 30 seconds then 14 cycles at 95C for 30 seconds, 55C for 1 minute, 68C for 12 minutes. The tube was placed on ice for 2 minutes, then 1 µl of DpnI was added and the tube incubated at 37C for one hour. One microliter of the DpnI-treated DNA was transformed into Epicurian coli XL1-Blue supercompetent cells and the entire insert was re-sequenced. The primer sequences are for LW1741: 5'-

GGTGCTCTGGAACCTCGGCTTCCGCATCAAGAAGGGCCCC-3' (SEQ ID NO:133) and for LW1742 was: 5'-GGGGCCCTTCTTGATGCGGAAGCCGAGGTTCAGAGCACC-3' (SEQ ID NO:134).

*nGPCR-61*

5           The PCR reaction was performed in 50 µl containing 34.5 µl H<sub>2</sub>O, 5 µl Buffer II (PE Applied Biosystems AmpliTaq Gold system), 6 µl 25 mM MgCl<sub>2</sub>, 2 µl 10 mM dNTP mix, 1.5 µl human genomic DNA (Clontech #6550-1, 0.1 µg/µl), 0.3 µl primer VR61C (1 µg/µl) (SEQ ID NO:135), 0.3 µl primer VR61D (1 µg/µl) (SEQ ID NO:136), and 0.4 µl AmpliTaq Gold™ DNA Polymerase. The primer sequence for VR61C was: 5'-TTCAAAGCTTATGAACAAC  
10 AATACAACATGTATTCAAC-3' (SEQ ID NO:135), corresponding to the 5' end of the coding region and containing a HindIII restriction site, and the primer sequence for VR61D was: 5'-TTCACTCGAGTCAAACATATGATTGCATATGTG-3' (SEQ ID NO:136), corresponding to the 3' end of the coding region and containing an XhoI restriction site(Genosys). The PCR reaction was carried out using a GeneAmp PCR9700 thermocycler (Perkin Elmer Applied  
15 Biosystems) and started with 1 cycle of 95C for 10 minutes, then 14 cycles at 95C for 30 seconds, 72C for 2 minutes decreasing 1C each cycle, 72C for 1 minute, followed by 30 cycles at 95C for 30 seconds, 60C for 30 seconds, 72C for 1 minute. The PCR reaction was loaded on a 0.75% agarose gel. The DNA band was excised from the gel and the DNA was eluted from the agarose using a QIAquick gel extraction kit (Qiagen). The eluted DNA was ethanol-precipitated  
20 and resuspended in 4µl H<sub>2</sub>O for ligation. The ligation reaction consisted of 4µl of fresh ethanol-precipitated PCR product and 1 µl of pCRII-TOPO vector (Invitrogen). The reaction was gently mixed and allowed to incubate for 5 minutes at room temperature followed by the addition of 1 µl of 6x TOPO cloning stop solution and mixing for 10 seconds at room temperature. The sample was then placed on ice and 2 µl was transformed in 50µl of One Shot cells (Invitrogen)  
25 and plated onto ampicillin plates. Four white colonies were chosen and the presence of an insert was verified by PCR in the following manner. Each colony was resuspended in 50 µL H<sub>2</sub>O. A 16 µl aliquot was removed and boiled for 5 minutes and the sample was placed on ice for 5 minutes. The sample was microfuged briefly to pellet any bacterial debris and PCR was carried out with 15 µl sample using primers VR61C and VR61D, above.

30           Colonies from the positive clones identified by PCR were used to inoculate a 4 ml culture of LB medium containing 100 µg/ml ampicillin. Plasmid DNA was purified using the Wizard Plus Minipreps DNA purification system (Promega). Since the primers used to PCR SEQ-61 from genomic DNA were engineered to have HindIII and XhoI sites, the cDNA obtained from the minipreps was digested with these restriction enzymes. Clones were verified

as having an insert of the correct size by gel electrophoresis. cDNA from one of the clones was then submitted for sequencing. Two mutations were found (bp 939 T→C and bp1004 G→T). The mutation at bp 939 was found to be a silent mutation and was not repaired. The mutation at bp 1004 was repaired as described below.

5       The mutation at bp 1004 in SEQ-61 was repaired using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). The PCR reaction contained 38.1 µl H<sub>2</sub>O, 5 µl 10x reaction buffer, 50 ng mini-prep cDNA from above, 1.25 µl primer VR61G (100 ng/µl) (SEQ ID NO:137), 1.25 µl primer VR61H (100 ng/µl) (SEQ ID NO:138), 2 µl 10 mM dNTP mix, and 1 µl Pfu DNA polymerase. The primer sequence for VR61G was: 5'-GCACATATGCAATCATA  
10   TGGTTGACTCGAGTGAAAAGGG-3' (SEQ ID NO:137) and the primer sequence for VR61H was: 5'- CCCTTTTCACTCGAGTCAACCCATATGATTGCATATGTGC-3' (SEQ ID NO:138), where the base highlighted and underlined is the one being corrected. The PCR cycle conditions were 95C for 30 seconds, then 12 cycles at 95C for 30 seconds, 55C for 1 minute, 68C for 11 minutes. One µl of DpnI was added and the reaction was incubated at 37C for 1  
15   hour. One µl of the DpnI-treated DNA was transformed into 50 µl Epicurian coli XL1-Blue supercompetent cells and plated onto ampicillin plates. Two colonies were chosen and miniprep DNA was purified as described above. The presence of an insert was verified by restriction digest/gel electrophoresis. DNA from both of the samples was submitted for sequencing. Sequencing information showed that the mutation was repaired in both clones. One of the clones  
20   was chosen for use in subsequent work.

      The clone described above was sequenced directly using an ABI377 fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI BigDye™ Terminator Cycle Sequencing Ready Reaction kit with Taq FSTM polymerase. Each ABI cycle sequencing reaction contained 0.5 µg of plasmid DNA. Cycle-sequencing was  
25   performed using an initial denaturation at 98C for 1 minute, followed by 50 cycles: 96C for 30 seconds, annealing at 50C for 30 seconds, and extension at 60C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using AGTC (R) gel filtration block (Edge Biosystems, Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which was then centrifuged in a  
30   swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 3 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90C for 3.5 minutes and loaded into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence

analysis was performed by importing ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI).

#### *nGPCR-51*

The cDNA clone for nGPCR-51 was isolated by Life Technologies, Inc. using their  
5 GENETRAPPER® cDNA Positive Selection System (Cat. No. 10356-020). Double-stranded plasmid DNA from the SUPERScript Human Fetal Brain cDNA Library (Cat. No. 10662-013) was used as a substrate. The library was made with fetal brain cDNA directionally cloned into the *NotI-Sall* site of pCMV-SPORT2. An oligonucleotide probe 5'-CTCCATCTGTCTC  
AGCTATGCCAGCAGCAG-3' (SEQ ID NO:139) was designed from Celera sequence  
10 GA\_11007426 and then biotinylated for solution hybridization to single-stranded fetal brain DNA (prepared from total cDNA library) and selection with paramagnetic beads.

#### *Preparation of dsDNA from a Plasmid cDNA Library*

One hundred ml of Terrific Broth containing 100 mg/ml ampicillin was inoculated with cells from an amplified library. The cells were grown to saturation at 30C. The sample was  
15 centrifuged at 4,800 g for 15 minutes at 4C. The supernatant was decanted, and the cell pellet was resuspended in a total volume of 10 ml of buffer I with RNase (15 mM Tris-HCl (pH 8.0), 10 mM EDTA, RNase A(100 µg/ml), RNase T1 (1,200 units/ml)). Ten ml of buffer II (0.2 M NaOH, 1% SDS) was added to the resuspended cells, and the sample was inverted to mix and allowed to incubate for 5 minutes at room temperature. Ten ml of cold 7.5 M NH<sub>4</sub>OAc was  
20 added to the cell mixture, and the sample was inverted to mix and allowed to incubate on ice for 10 minutes. The sample was centrifuged at 3,000 g (4,000 rpm in an HS4 rotor) for 15 minutes at 4C. The supernatant was poured through cheesecloth into a fresh 50-ml centrifuge tube. An equal volume of cold isopropanol (-20C) was added to the tube, mixed well, and the sample was centrifuged at 3,000 g for 15 minutes at 4C. The supernatant was discarded, and the pellet was  
25 resuspended in 1 ml of buffer I with RNase and transferred to a microcentrifuge tube. The solution was clarified by centrifugation at 4C for 1 minute at 14,000 g. The supernatant was transferred to a fresh microcentrifuge tube, and incubated at 37C for 30 minutes, after which the tube was incubated at 65C for 5 minutes. The sample was split into two equal parts (~500 µl each) in 1.5-ml microcentrifuge tubes. An equal volume of phenol: chloroform:isoamyl alcohol  
30 (25:24:1) was added to each sample and the tube was vortexed. The sample was centrifuged at room temperature for 5 minutes at 14,000 g. Four hundred fifty µl of the upper aqueous phase was transferred to fresh microcentrifuge tubes. The phenol:chloroform extraction procedure above was repeated 3 times. An equal volume of isopropanol (-20C) was added to each tube. The tube was centrifuged at 4C for 15 minutes at 14,000 g, and the supernatant was discarded.

Five hundred  $\mu\text{l}$  of 70% ethanol was carefully added to each tube. The tubes were centrifuged at 4C for 5 minutes at 14,000 g. The supernatant was discarded, and the pellet was dried at room temperature for 10 minutes. The pellets were resuspended in a total of 200 $\mu\text{l}$  TE buffer.

#### *Generation of ssDNA*

5           Five  $\mu\text{g}$  ds phagemid cDNA (1  $\mu\text{g}/\mu\text{l}$ ), 2  $\mu\text{l}$  10X Gene II Buffer, and sufficient autoclaved, distilled water to bring the volume to 19  $\mu\text{l}$  was added to a 1.5 ml microcentrifuge tube at room temperature. To each sample, 1  $\mu\text{l}$  of Gene II (phage F1 endonuclease) was added. The sample was vortexed and centrifuged at room temperature for 2 s at 14,000 g to collect the contents to the bottom of the tube. The tubes were incubated in a 30C water bath for 25  
10   minutes. The mixture was heated at 65C for 5 minutes and immediately chilled on ice for 1 minute. Two  $\mu\text{l}$  of Exo III were added, and the tube was vortexed and centrifuged at room temperature for 2 s at 14,000 g. The tubes were incubated at 37C for 60 minutes. An equal volume (20  $\mu\text{l}$ ) of phenol:chloroform:isoamyl alcohol (25:24:1) was added to each tube, vortexed thoroughly, and centrifuged at room temperature for 5 minutes at 14,000 g to separate  
15   the phases. Eighteen  $\mu\text{l}$  of the upper aqueous phase was transferred to a fresh 1.5-ml microcentrifuge tube.

#### *Biotinylation Reaction*

          The following components were added to a 1.5 ml microcentrifuge tube:  
5  $\mu\text{l}$  5X TdT Buffer; 3  $\mu\text{g}$  oligonucleotide (SEQ ID NO:139), 5  $\mu\text{l}$  Biotin-14-dCTP, sufficient  
20   autoclaved, distilled water to bring the volume to 23  $\mu\text{l}$ , and 2  $\mu\text{l}$  TdT (Terminal Deoxynucleotidyl Transferase). Each tube is vortexed gently and centrifuged for 2 s at 14,000 g. The tubes were incubated for 1 hour at 30C. After 1 hour, 1  $\mu\text{l}$  of Glycogen (20  $\mu\text{g}/\mu\text{l}$ ), 26  $\mu\text{l}$  of 1 M Tris-HCl (pH 7.5), and 120  $\mu\text{l}$  of ethanol were added to the oligonucleotide biotinylation reaction. Each sample was vortexed and stored on dry ice for 10 minutes. Each tube was  
25   centrifuged at 4C for 30 minutes at 14,000 g. The supernatant was removed from the microcentrifuge tubes, and 200  $\mu\text{l}$  of 70% ethanol (-20C) was layered over the pellet. Each tube was centrifuged at 4C for 2 minutes at 14,000 g. The ethanol wash was removed from the microcentrifuge tubes, and the ethanol wash was repeated once. The pellets were dried at room temperature for 10 minutes or until completely dry. The biotinylated oligonucleotide was  
30   resuspended in 20  $\mu\text{l}$  TE buffer.

#### *Oligonucleotide Hybridization*

          The biotinylated oligonucleotide was diluted to 20 ng/ $\mu\text{l}$  in TE buffer. The 4X Hybridization Buffer was incubated for 2 minutes at 37C, mixed well, and 6  $\mu\text{l}$  was added to the remaining 17  $\mu\text{l}$  of Gene II/Exo III-treated DNA. The DNA was denatured in a 95C water bath

for 1 minutes, and then chilled immediately on ice for 1 minute. One  $\mu$ l of diluted biotinylated oligonucleotide (20 ng) was added to the denatured DNA. The tube was incubated at 37C in a water bath or incubator for 1 hour.

#### *Streptavidin Paramagnetic Bead Preparation*

5           The beads were gently mixed by pipetting until the beads at the bottom of the tube were completely resuspended. For each reaction, 45  $\mu$ l of the mixed beads were transferred to the bottom of a microcentrifuge tube. The tubes were placed in the magnet and allowed to sit for 2 minutes. With the tubes still in the magnet, the supernatant was removed by pipetting. One hundred  $\mu$ l of TE buffer was immediately added to the beads. The tubes were removed from the magnet and the beads were resuspended by finger-tapping or vortexing at the lowest setting. The 10 tubes were reinserted into the magnet, and after 2 min, the supernatant was removed. The beads were resuspended in 30  $\mu$ l of TE buffer.

#### *cDNA Capture*

          The hybridization mixture was removed from the 37C water bath and centrifuged at 15 room temperature for 2 s at 14,000 g. The prepared paramagnetic beads were pipetted into the mixture and gently mixed by pipetting. The suspension was incubated for 30 minutes at room temperature. The suspension was gently mixed frequently to resuspend the beads. The tubes were inserted into the magnet, and after 2 min, the supernatant was removed and discarded. One hundred  $\mu$ l of Wash Buffer was then added to the beads. The beads were resuspended and the 20 tubes reinserted into the magnet for 2 minutes. The supernatant was removed and discarded. The washing step was repeated one more time. One hundred  $\mu$ l of Wash Buffer was added to the beads. The beads were gently resuspended by pipetting up and down (not by vortexing) and the solution was transferred to a new tube. The tubes were inserted into the magnet for 5 minutes. The supernatant was removed from the paramagnetic beads and discarded. One hundred  $\mu$ l of 25 Wash Buffer was immediately added and the tubes mixed by finger tapping or gently vortexing. The tubes were placed into the magnet for 5 minutes. For each elution, 14  $\mu$ l of TE buffer (pH 8.0) was mixed with 7  $\mu$ l of the 3X Elution Buffer. After the 5 minute incubation, the supernatant was removed and discarded from the paramagnetic beads; 20  $\mu$ l of 1X elution buffer was added to the beads, and mixed well. The beads were incubated for 5 minutes at room 30 temperature. During the incubation, the beads were mixed for 10 seconds every minute. The tube was inserted into the magnet and allowed to sit for 5 minutes. The supernatant (containing the captured cDNA clone) was collected and transferred to a fresh tube, and the beads were resuspended in 15  $\mu$ l of TE buffer. The tube was inserted into the magnet and allowed to sit for 5 minutes. The supernatant was transferred from the tube and combined with the previous

supernatant. The tube containing the combined supernatants was inserted into the magnet for 10 minutes to remove any remaining paramagnetic beads and the supernatant was transferred to a fresh microcentrifuge tube. To the supernatant (~35 µl), 1 µl of Glycogen, 18 µl of 7.5 M NH<sub>4</sub>OAc, and 135 µl of ethanol (-20C) were added. The tube was mixed well and stored on ice for 5 10 minutes. The tube was centrifuged at 4C for 30 minutes at 14,000 g. The supernatant was removed from the small pellets. One hundred µl of 70% ethanol was added to each tube. The tubes were centrifuged at room temperature for 2 minutes at 14,000 g. The ethanol was removed and the pellets dried at room temperature until dry. The pellets were resuspended in 5 µl of TE buffer and stored at 4C.

#### 10 *Repair of Captured cDNA: Repair Reaction*

The thermal cycler is programmed for one cycle as described below:

90C denature step for 1 minute; 55C annealing step for 30 seconds; 70C extension step for 15 minutes. A DNA primer/repair mix was prepared for each capture reaction by adding the following to the captured cDNA: captured DNA (5 µl), autoclaved, distilled water (11 µl), 50 ng 15 oligonucleotide (SEQ ID NO:139; not biotinylated)( 1 µl), 10 mM dNTP Mix (0.5 µl), 10X Repair Buffer (2 µl), Repair Enzyme (0.5 µl). The tubes were mixed and centrifuged at room temperature for 2 s at 14,000 g. The DNA primer/repair mix was incubated at 90C for 1 minute. The mix was transferred to 55C and incubated for 30 seconds. The mix was transferred to 70C and incubated for 15 minutes to allow primer extension. The mix was centrifuged at room 20 temperature for 2 s at 14,000 g. The repaired DNA was precipitated by adding 1 µl Glycogen, 11 µl 7.5 M ammonium acetate, and 90 µl of -20C ethanol to each tube. The tubes were vortexed and placed in ice for 10 minutes. The tubes were centrifuged at 4C for 30 minutes at 14,000 g. The ethanol was removed from the small pellet and 100 µl of 70% ethanol (-20C) was layered on the pellet. The tubes were centrifuged at 4C for 2 minutes at 14,000 g. The ethanol 25 was removed and the pellets dried at room temperature until dry. The pellets were dissolved in 10 µl of TE buffer.

#### *Transformation with ULTRAMAX DH5a a-FT Cells*

Competent cells were removed from -70C freezer and thawed on wet ice. The required number of 17 × 100 polypropylene tubes (Falcon 2059) were placed on ice. Immediately after 30 thawing, the cells were gently mixed, then 100 µl of cells were aliquotted into chilled polypropylene tubes. Three µl of the DNA were mixed into each individual tube of cells. The cells were incubated on ice for 30 minutes. Cells were heat-shocked for 45 seconds in a 42C water bath, and then placed on ice for 2 minutes. To each tube, 0.9 ml of room temperature S.O.C medium was added. The tubes were shaken at 225 rpm (37C) for 1 hour. For captured or



repaired cDNA samples, 100 µl and 200 µl aliquots were plated onto LB plates containing 100 µg/ml ampicillin. The plates were incubated overnight in a 37C incubator.

*Identification of Desired cDNA Clones by Colony PCR*

A PCR master mix was prepared with the following components: 22.5 µl/clone of 1X  
5 PCR SUPERMIX, forward and reverse primers at a final concentration of 200 nM/primer, and autoclaved, distilled water to a final volume of 25 µl. Using a micropipette tip or sterile toothpick, each colony was picked and placed into an individual tube containing master mix. The PCR reaction was as follows: 1 cycle at 94C, 1 minute, then 30 cycles of 94C, 30 seconds; 55C, 30 seconds; 72C, 1 minute. The PCR primer pairs used in colony screening were, forward  
10 primer: 5'-CCATCTGTCTCAGCTATGCC-3' (SEQ ID NO:140), and the reverse primer: 5'-TCCTTCTCAGTCGCTCTTC-3' (SEQ ID NO:141).

*nGPCR-52*

Two microliters of a human genomic library (~10<sup>8</sup> PFU/ml) (Clontech) were added to 6 ml of an overnight culture of K802 cells (Clontech), then distributed as 250 µl aliquots into each  
15 of 24 tubes. The tubes were incubated at 37C for 15 minutes. Seven milliliters of 0.8% agarose was added to each tube, mixed, then poured onto LB agar + 10 mM MgSO<sub>4</sub> plates and incubated overnight at 37C. To each plate 5 ml of SM (0.1M NaCl, 8.1 mM MgSO<sub>4</sub>·7H<sub>2</sub>O, 50mM Tris-Cl (pH 7.5), 0.0001% gelatin) phage buffer was added and the top agarose was removed with a microscope slide and placed in a 50 ml centrifuge tube. A drop of chloroform was added and the  
20 tube was placed in a 37C shaker for 15 minutes, then centrifuged for 20 minutes at 4000 RPM (Sorvall RT6000 table top centrifuge) and the supernatant stored at 4C as a stock solution.

The PCR reaction was performed in 20 µl containing 8.8 µl H<sub>2</sub>O, 4 µl 5X Rapid-Load Buffer (Origene), 2 µl 10xPCR buffer II (Perkin-Elmer), 2 µl 25 mM MgCl<sub>2</sub>, 0.8 µl 10 mM dNTP, 0.12 µl LW1632 (SEQ ID NO:142) (1 µg/µl), 0.12 µl LW1633 (SEQ ID NO:143) (1  
25 µg/µl), 0.2 µl AmpliTaq Gold polymerase (Perkin Elmer) and 2 µl of phage from each of the 24 tubes. The PCR reaction involved 1 cycle at 80C for 20 minutes, 95C for 10 minutes, then 22 cycles at 95C for 30 seconds, 72C for 4 minutes decreasing 1C each cycle, 68C for 2 minutes, followed by 30 cycles at 95C for 30 seconds, 55C for 30 seconds, 68C for 60 seconds. The reaction was loaded onto a 2 % agarose gel. From the tube that gave a PCR product of the  
30 correct size, 5 µl was used to set up five 1:10 dilutions that were plated onto LB agar + 10 mM MgSO<sub>4</sub> plates and incubated overnight. A BA85 nitrocellulose filter (Schleicher & Schuell) was placed on top of each plate for 1 hour. The filter was removed, placed phage side up in a petri dish, and covered with 4 ml of SM for 15 minutes to elute the phage. One milliliter of SM was removed from each plate and used to set up a PCR reaction as above. The plate of the lowest

dilution to give a PCR product was subdivided, filter-lifted and the PCR reaction was repeated. The series of dilutions and subdividing of the plate was continued until a single plaque was isolated that gave a positive PCR band. Once a single plaque was isolated, 10 µl phage supernatant was added to 100 µl SM and 200 µl of K802 cells per plate with a total of 8 plates set up. The plates were incubated overnight at 37C. The top agarose was removed by adding 8 ml of SM, then scrapping off the agarose with a microscope slide and collected in a centrifuge tube. To the tube, 3 drops of chloroform was added, vortexed, incubated at 37C for 15 minutes then centrifuged for 20 minutes at 4000 RPM (Sorvall RT6000 table top centrifuge) to recover the phage, which was used to isolate genomic phage DNA using the Qiagen Lambda Midi Kit. The sequence for primer LW1632 was 5'-CCTCCACTTGTGCTTCATC-3' (SEQ ID NO:142) and for LW1633: 5'-AAAATCTATCAACACCCAGCC-3' (SEQ ID NO:143).

For subcloning the coding region of nGPCR-52, PCR was performed in a 50 µl reaction containing 33 µl H<sub>2</sub>O, 5 µl 10X TT buffer (140 mM Ammonium Sulfate, 0.1 % gelatin, 0.6 M Tris-tricine pH 8.4), 5 µl 15 mM MgSO<sub>4</sub>, 2 µl 10 mM dNTP, 4 µl genomic phage DNA (0.1 µg/µl), 0.3 µl LW1643 (SEQ ID NO:144) (1 µg/µl), 0.3 µl LW1644 (SEQ ID NO:145) (1 µg/µl), 0.4 µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94C for 2 minutes followed by 15 cycles at 94C for 30 seconds, 55C for 60 seconds, and 68C for 2 minutes. The PCR reaction was loaded onto a 2 % agarose gel. The DNA band was excised from the gel, placed in GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed. The eluted DNA was EtOH precipitated and resuspended in 12µl H<sub>2</sub>O for ligation. The PCR primer sequence for LW1643: 5'-GATCAAGCTTACCATGACCAGCAATTTTCCC-3' (SEQ ID NO:144) and for LW1644 was: 5'-GATCCTCGAGCTTATTCTAAAAATAAACTAATGG-3' (SEQ ID NO:145). The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen) which consisted of 4µl PCR product DNA and 1 µl pCRII-TOPO vector that was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice for 30 minutes. The cells were heat-shocked for 30 seconds at 42C, placed on ice for two minutes, 250 µl of SOC was added, then incubated at 37C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid Plasmid Miniprep System (GibcoBRL) and then sequenced.

nGPCR-52 genomic phage DNA was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. The cycle-sequencing reaction contained 14 µl of H<sub>2</sub>O, 16 µl of BigDye Terminator mix, 7 µl  
5 genomic phage DNA (0.1 µg/µl), and 3 µl primer (25 ng/µl). The reaction was performed in a Perkin-Elmer 9600 thermocycler at 95C for 5 minutes, followed by 99 cycles of 95C for 30 seconds, 55C for 20 seconds, and 60C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridges, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent. The samples were heated at 95C for 5 minutes then placed in the 310  
10 Genetic Analyzer

### Example 3: Subcloning of the Coding Region of nGPCR-x Via PCR

Additional experiments may be conducted to subclone the coding region of nGPCR and place the isolated coding region into a useful vector. Two additional PCR primers are designed based on the coding region of nGPCR, corresponding to either end. To protect against  
15 exonucleolytic attack during subsequent exposure to enzymes, *e.g.*, Taq polymerase, primers are routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target.

PCR is performed in a 50 µl reaction containing 34 µl H<sub>2</sub>O, 5 µl 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl 15 mM MgSO<sub>4</sub>, 2 µl dNTP  
20 mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 3 µl genomic phage DNA (0.25 µg/µl), 0.3 µl Primer 1 (1 µg/µl), 0.3 µl Primer 2 (1 µg/µl), 0.4 µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94C for 2 minutes; followed by 25 cycles at 94C for 30 seconds, 55C for 30 seconds, and 72C for 1.3 minutes.

The contents from the PCR reaction are loaded onto a 2% agarose gel and fractionated.  
25 The DNA band of expected size is excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA is precipitated with ethanol and resuspended in 6 µl H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the coding region is cloned into pCR2.1 using a protocol standard in the art. In particular, the ligation reaction consists of 6 µl of GPCR  
30 DNA, 1 µl 10X ligation buffer, 2 µl pCR2.1 (25 ng/µl, Invitrogen), and 1 µl T4 DNA ligase (Invitrogen). The reaction mixture is incubated overnight at 14C and the reaction is then stopped by heating at 65C for 10 minutes. Two microliters of the ligation reaction are transformed into One Shot cells (Invitrogen) and plated onto ampicillin plates. A single colony containing a recombinant pCR2.1 bearing an insert is used to inoculate a 5 ml culture of LB medium.

Plasmid DNA is purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, a 50 ml culture of LB medium is inoculated with the transformed One Shot cells, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-GPCR.

5 *nGPCR-70*

Colonies from the positive clones identified by PCR were used to inoculate a 4 ml culture of LB medium containing 100 µg/ml ampicillin. Plasmid DNA was purified using the Wizard Plus Minipreps DNA purification system (Promega). Since the primers used to PCR SEQ-70 from human brain cDNA were engineered to have HindIII and XhoI sites, the DNA  
10 obtained from the minipreps was digested with these restriction enzymes. Five of the clones were verified by gel electrophoresis to give a DNA band of the correct size. cDNA from each of these clones was then submitted for sequencing. Each of the clones was found to have two or more mutations. The clone containing the fewest mutations (clone #7-2 mutations at bp 561 G→A and bp 1093 G→A) was repaired as described as below.

15 The mutations in SEQ-70 were repaired sequentially using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). The mutation at bp 561 was repaired first and the primer sequences used were: VR70G 5'-CATGATCTGGGGGGCCAGCCCCAGC-3' (SEQ ID NO:146) and VR70H 5'-GCTGGGGCTGGCCCCCCCAGATCATG-3' (SEQ ID NO:147) where the base highlighted and underlined is the one being corrected. The PCR reaction contained 38.4  
20 µl H<sub>2</sub>O, 5 µl 10x reaction buffer, 50 ng mini-prep cDNA from clone #7, 2 µl 10 mM dNTP mix, and 1 µl Pfu DNA polymerase. The cycle conditions were 95C for 30 seconds, then 12 cycles at 95C for 30 seconds, 55C for 1 minute, 68C for 11 minutes. One µl of DpnI was added and the reaction was incubated at 37C for 1 hour. One µl of the DpnI-treated DNA was transformed into 50 µl Epicurian coli XL1-Blue supercompetent cells and plated onto ampicillin  
25 plates. Four colonies were chosen and miniprep DNA was purified as described above. DNA from one of the preps was used as a template to repair the second mutation at bp 1093. The primer sequences used were: VR70I 5'-GACATCAATTTCAGTGAGGATGACGTCGAGGCAG-3' (SEQ ID NO:148) and VR70J 5'-CTGCCTCGACGTCATCCCTCACTGAAATTGATGTC-3' (SEQ ID NO:149), where the base highlighted and underlined is the one being corrected.  
30 The PCR and transformation reactions were carried out as described above. Mini-prep DNA was prepared from three colonies, and the presence of an insert was verified by restriction digest/gel electrophoresis. DNA from one of the samples was submitted for sequencing. Sequencing information showed that both mutations were repaired but that an additional mutation at bp 560 (G→A) had been introduced. This mutation was adjacent to the original mutation at bp561 and

was encompassed in the sequence of the primer that was used. Since it could not be ruled out that there was an error in the synthesis of the primers, the same exact sequence corresponding to primers VR70G (SEQ ID NO:146) and VR70H (SEQ ID NO:147) was reordered and repair of the mutation with the new primers was carried out as above. DNA from 2 clones was submitted for sequencing. One of the clones had no mutations and was used for subsequent work.

The clone described above was sequenced directly using an ABI377 fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI BigDye™ Terminator Cycle Sequencing Ready Reaction kit with Taq FST™ polymerase. Each ABI cycle sequencing reaction contained 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98C for 1 minute, followed by 50 cycles: 96C for 30 seconds, annealing at 50C for 30 seconds, and extension at 60C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using AGTC (R) gel filtration block (Edge Biosystems, Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which was then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 3 µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90C for 3.5 minutes and loaded into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence analysis was performed by importing ABI377 files into the Sequencer program (Gene Codes, Ann Arbor, MI).

#### *nGPCR-42*

The DNA subcloned into pCRII was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) and was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96C for 10 seconds, 50C for 10 seconds, and 60C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95C for 5 minutes then placed in the 310 Genetic Analyzer.

#### *nGPCR-46*

The DNA subcloned into pCRII was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology

and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) and was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96C for 10 seconds, 50C for 10 seconds, and 60C for 4 minutes.

5 The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95C for 5 minutes then placed in the 310 Genetic Analyzer.

The mutation found in the clone was repaired using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). The PCR reaction contained 39 µl H<sub>2</sub>O, 5 µl 10xReaction buffer,

10 1 µl mini-prep DNA (150 ng/µl), 2µl 10 mM dNTP, 1 µl Pfu DNA polymerase, 1 µl LW1638 (SEQ ID NO:150) (125 ng/µl), and 1 µl LW1639 (SEQ ID NO:151) (125 ng/µl). The cycle conditions were 95C for 30 seconds then 12 cycles at 95C for 30 seconds, 55C for 1 minute, 68C for 12 minutes. The tube was placed on ice for 2 minutes, then 1 µl of DpnI was added and the tube incubated at 37C for one hour. Two microliters of the *DpnI*-treated DNA was

15 transformed into Epicurian coli XL1-Blue supercompetent cells and the entire insert was re-sequenced. The forward PCR primer sequence was: LW1638 5'-CGAACTCCGCACTCCTGG CCAGGGCCCTGCGGGC-3' (SEQ ID NO:150) and the reverse PCR primer sequence was: LW1639 5'-GCCCCGAGGGCCCTGGCCAGGAGTGCGGAGTTC-3' (SEQ ID NO:151).

*nGPCR-48*

20 The DNA subcloned into pCRII was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) and was performed in a Perkin-Elmer 9600

25 thermocycler with 25 cycles of 96C for 10 seconds, 50C for 10 seconds, and 60C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95C for 5 minutes then placed in the 310 Genetic Analyzer.

*nGPCR-49*

30 The DNA subcloned into pCRII was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) (SEQ ID NO:131 and SEQ ID NO:132) and was

performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96C for 10 seconds, 50C for 10 seconds, and 60C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95C for 5 minutes then placed in the 310

5 Genetic Analyzer.

#### *nGPCR-52*

The DNA subcloned into pCRII was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each  
10 cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) and was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96C for 10 seconds, 50C for 10 seconds, and 60C for 4 minutes. The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples  
15 were heated at 95C for 5 minutes then placed in the 310 Genetic Analyzer.

#### **Example 4: Hybridization Analysis to Demonstrate nGPCR-X Expression in Brain**

The expression of nGPCR-x in mammals, such as the rat, may be investigated by *in situ* hybridization histochemistry. To investigate expression in the brain, for example, coronal and sagittal rat brain cryosections (20 µm thick) are prepared using a Reichert-Jung cryostat.  
20 Individual sections are thaw-mounted onto silanized, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80C. Sections are processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold phosphate-buffered saline (PBS), acetylated using acetic anhydride in triethanolamine buffer, and dehydrated through a series of alcohol washes in 70%, 95%, and 100% alcohol at room temperature. Subsequently, sections are delipidated in  
25 chloroform, followed by rehydration through successive exposure to 100% and 95% alcohol at room temperature. Microscope slides containing processed cryosections are allowed to air dry prior to hybridization. Other tissues may be assayed in a similar fashion.

A nGPCR-x-specific probe is generated using PCR. Following PCR amplification, the fragment is digested with restriction enzymes and cloned into pBluescript II cleaved with the  
30 same enzymes. For production of a probe specific for the sense strand of nGPCR-x, the nGPCR-x clone in pBluescript II is linearized with a suitable restriction enzyme, which provides a substrate for labeled run-off transcripts (*i.e.*, cRNA riboprobes) using the vector-borne T7 promoter and commercially available T7 RNA polymerase. A probe specific for the antisense strand of nGPCR-x is also readily prepared using the nGPCR-x clone in pBluescript II by

cleaving the recombinant plasmid with a suitable restriction enzyme to generate a linearized substrate for the production of labeled run-off cRNA transcripts using the T3 promoter and cognate polymerase. The riboprobes are labeled with [<sup>35</sup>S]-UTP to yield a specific activity of about  $0.40 \times 10^6$  cpm/pmol for antisense riboprobes and about  $0.65 \times 10^6$  cpm/pmol for sense-strand riboprobes. Each riboprobe is subsequently denatured and added (2 pmol/ml) to hybridization buffer which contained 50% formamide, 10% dextran, 0.3 M NaCl, 10 mM Tris (pH 8.0), 1 mM EDTA, 1X Denhardt's Solution, and 10 mM dithiothreitol. Microscope slides containing sequential brain cryosections are independently exposed to 45  $\mu$ l of hybridization solution per slide and silanized cover slips are placed over the sections being exposed to hybridization solution. Sections are incubated overnight (15-18 hours) at 52C to allow hybridization to occur. Equivalent series of cryosections are exposed to sense or antisense nGPCR-x-specific cRNA riboprobes.

Following the hybridization period, coverslips are washed off the slides in 1X SSC, followed by RNase A treatment involving the exposure of slides to 20  $\mu$ g/ml RNase A in a buffer containing 10 mM Tris-HCl (pH 7.4), 0.5 M EDTA, and 0.5 M NaCl for 45 minutes at 37C. The cryosections are then subjected to three high-stringency washes in 0.1 X SSC at 52C for 20 minutes each. Following the series of washes, cryosections are dehydrated by consecutive exposure to 70%, 95%, and 100% ammonium acetate in alcohol, followed by air drying and exposure to Kodak BioMax™ MR-1 film. After 13 days of exposure, the film is developed.

Based on these results, slides containing tissue that hybridized, as shown by film autoradiograms, are coated with Kodak NTB-2 nuclear track emulsion and the slides are stored in the dark for 32 days. The slides are then developed and counterstained with hematoxylin. Emulsion-coated sections are analyzed microscopically to determine the specificity of labeling. The signal is determined to be specific if autoradiographic grains (generated by antisense probe hybridization) are clearly associated with cresyl violet-stained cell bodies. Autoradiographic grains found between cell bodies indicates non-specific binding of the probe.

Expression of nGPCR-x in the brain provides an indication that modulators of nGPCR-x activity have utility for treating neurological disorders, including but not limited to, mental disorder, affective disorders, ADHD/ADD (*i.e.*, Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), and neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, and senile dementia. Some other diseases for which modulators of nGPCR-x may have utility include depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of nGPCR-x modulators, including



nGPCR-x ligands and anti-nGPCR-x antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### *In Situ Hybridization of SEQ-51*

Coronal and sagittal oriented rat brain sections were cryosectioned (20  $\mu$ m thick) using a  
5 Leica CM3050 cryostat. The individual sections were thaw-mounted onto silanated, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80C. The sections were processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold PBS, acetylated using acetic anhydride in triethanolamine buffer and dehydrated through 70%, 95%, and 100% alcohols at room temperature (RT). This was followed with delipidation in  
10 chloroform then rehydration in 100% and 95% alcohol at RT. Sections were allowed to air dry prior to hybridization. For nGPCR-51, a 250 bp PCR fragment (spanning nt 83 to 332) was generated that contained T7 polymerase on either the 5' end (sense) with primers LW1743 5'-GCGTAATACGACTCACTATAGGGAGACCTGCCAGTGTGGTAGATACAG-3' (SEQ ID NO:152) and LW1746 5'-GGATGTGATGATGGTGCAG-3' (SEQ ID NO:153) or the 3' end  
15 (antisense) with primers LW1744 5'-GCGTAATACGACTCACTATAGGGAGACCGGATGTGATGATGGTGCAG-3' (SEQ ID NO:154) and LW1745 5'-TGCCAGTGTGGTAGATACAG-3' (SEQ ID NO:155). nGPCR-51 was labeled with  $^{35}$ S-UTP to yield a specific activity of  $0.655 \times 10^6$  cpm/pmol for antisense and  $0.675 \times 10^6$  cpm/pmol for sense probe. Both riboprobes were denatured and added to hybridization buffer which contained 50% formamide, 10%  
20 dextran, 0.3M NaCl, 10 mM Tris, 1 mM EDTA, 1X Denhardt's, and 10 mM DTT. Sequential brain cryosections were hybridized with 45  $\mu$ l/slide of the sense and antisense riboprobe hybridization mixture then coverslipped with silanized glass coverslips. The sections were hybridized overnight (15-18 hours) at 50C in an incubator.

Coverslips were washed off the slides in 1X SSC, followed by RNase A treatment, and  
25 high temperature stringency washes (3X, 20 minutes at 49C) in 0.1X SSC. Slides were dehydrated with 70%, 95% and 100%  $\text{NH}_4\text{OAc}$  alcohols, air dried and exposed to Kodak BioMax MR-1 film. After 8 days of exposure, the film was developed. This was followed with coating selected tissue slides with Kodak NTB-2 nuclear track emulsion and storing the slides in the dark for 16 days. The slides were then developed and counterstained with hematoxylin.  
30 Emulsion-coated sections were analyzed microscopically to determine the specificity of labeling. Presence of autoradiographic grains (generated by antisense probe hybridization) over cell bodies (versus between cell bodies) was used as an index of specific hybridization.

Sense and antisense  $^{35}$ S-labeled RNA probes were generated using a 250-bp fragment of nGPCR-51 for *in situ* hybridization histochemistry. Specific labeling with the antisense probe

showed wide spread distribution of nGPCR-51 mRNA. Localization appears in the piriform ctx, habenula, bed nucleus of stria terminalis, islands of Calleja, olfactory tubercle, hippocampus, hypothalamus, PVN, red nucleus, interpeduncular nucleus, dorsal raphe, substantia nigra pars compacta, and reticular thalamus. Expression of nGPCR-51 in these brain regions provided an indication that modulators of nGPCR-51 activity have utility for treating disorders, including but not limited to, schizophrenia, major depression, bipolar disease, anxiety disorder, Parkinson's disease, endocrine disorders, Alzheimer's disease and the like.

#### *In Situ Hybridization SEQ-52*

Coronal and sagittal oriented rat brain sections were cryosectioned (20 um thick) using a Leica CM3050 cryostat. The individual sections were thaw-mounted onto silanated, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80C. The sections were processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold PBS, acetylated using acetic anhydride in triethanolamine buffer and dehydrated through 70%, 95%, and 100% alcohols at room temperature. This was followed with delipidation in chloroform then rehydration in 100% and 95% alcohol at room temperature. Sections were allowed to air dry prior to hybridization. For nGPCR-52, a 292 bp PCR fragment (spanning nt 2920 to 3211) was generated that contained T7 polymerase on either the 5' end (sense) with primers LW1682 5'-GCGTAATACGACTCACTATAGGGAGACCACCAGCAATTTTCCCAACC-3' (SEQ ID NO:156) and LW1685 5'-AATACCAGCAGCTCTCCAC-3' (SEQ ID NO:157) or the 3' end (antisense) with primers LW1683 5'-GCGTAATACGACTCACTATAGGGAGACCAATACCAGCAGCTCTCCAC-3' (SEQ ID NO:158) and LW1784 5'-ACCAGCAATTTTCCCAACC-3' (SEQ ID NO:159). nGPCR-52 was labeled with <sup>35</sup>S-UTP to yield a specific activity of 0.686 x 10<sup>6</sup> cpm/pmol for the antisense probe and 0.601 x 10<sup>6</sup> cpm/pmol for the sense probe. Both riboprobes were denatured and added to hybridization buffer which contained 50% formamide, 10% dextran, 0.3M NaCl, 10 mM Tris, 1 mM EDTA, 1X Denhardt's, and 10 mM DTT. Sequential brain cryosections were hybridized with 45 TI/slide of the sense and antisense riboprobe hybridization mixture then coverslipped with silanized glass coverslips. The sections were hybridized overnight (15-18 hours) at 49C in an incubator.

Coverslips were washed off the slides in 1X SSC, followed by RNase A treatment, and high temperature stringency washes (3X, 20 minutes at 48C) in 0.1X SSC. Slides were dehydrated with 70%, 95% and 100% NH<sub>4</sub>OAc alcohols, air dried and exposed to Kodak BioMax MR-1 film. After 8 days of exposure, the film was developed. This was followed with coating selected tissue slides with Kodak NTB-2 nuclear track emulsion and storing the slides in the dark for 16 days. The slides were then developed and counterstained with hematoxylin.

Emulsion-coated sections were analyzed microscopically to determine the specificity of labeling. Presence of autoradiographic grains (generated by antisense probe hybridization) over cell bodies (versus between cell bodies) was used as an index of specific hybridization.

To determine which regions of the brain nGPCR-52 is expressed in, a 292 bp fragment of nGPCR-52 was used to generate sense and antisense <sup>35</sup>S-labeled RNA probes for *in situ* hybridization histochemistry. Specific labeling with the antisense probe showed wide spread distribution of nGPCR-52 mRNA. Localization appears in the cortex, piriform ctx, habenula, islands of Calleja, hippocampus, hypothalamus, red nucleus, dorsal raphe, substantia nigra pars compacta, and reticular thalamus. The regions where nGPCR-52 is expressed are within the limbic and neuroendocrine circuitry of the brain.

#### **Example 5: Tissue Expression Profiling**

A PCR-based system (RapidScan™ Gene Expression Panel, OriGene Technologies, Rockville, MD) may be used to generate a comprehensive expression profile of the putative nGPCR-x in human tissue, and in human brain regions. The RapidScan Expression Panel is comprised of first-strand cDNAs from various human tissues and brain regions that are serially diluted over a 4-log range and arrayed into a multi-well PCR plate. Human tissues in the array may include: brain, heart, kidney, spleen, liver, colon, lung, small intestine, muscle, stomach, testis, placenta, salivary gland, thyroid, adrenal gland, pancreas, ovary, uterus, prostate, skin, PBL, bone marrow, fetal brain, and fetal liver.

Expression of nGPCR-x in various tissues is detected using PCR primers designed based on the available sequence of the receptor that will prime the synthesis of a predetermined size fragment in the presence of the appropriate cDNA.

PCR is performed in a 50 µl reaction containing 34 µl H<sub>2</sub>O, 5µl 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 µl 15 mM MgSO<sub>4</sub>, 2 µl dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 0.3 µl forward primer (1 µg/µl), 0.3 µl reverse primer (1 µg/µl), 0.4 µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction mixture is added to each well of the PCR plate. The plate is placed in a MJ Research PTC100 thermocycler, and is then exposed to the following cycling parameters: Pre-soak 94C for 3 minutes; denaturation at 94C for 30 seconds; annealing at primer 57C for 45 seconds; extension 72C for 2 minutes; for 35 cycles. PCR productions are then separated and analyzed by electrophoresis on a 1.2% agarose gel stained with ethidium bromide. The 4-log dilution range of cDNA deposited on the plate ensures that the amplification reaction is within the linear range and, hence, facilitates semi-quantitative determination of relative mRNA accumulation in the various tissues or brain regions examined.

*nGPCR-70*

Tissue specific expression of the cDNA encoding nGPCR-70 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 6 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of the putative GPCR in human tissues. Human tissues arrayed included: brain, heart, kidney, peripheral blood leukocytes, lung and testis. PCR primers were designed based on the available sequence of the Celera sequence GA\_16417344. The forward primer used was: 5'-GTGACTAACTCTGCCT GCG-3' (SEQ ID NO:160). The reverse primer used was: 5'-TTGCGCTGCAACACTAGCG-3' (SEQ ID NO:161). This primer set primed the synthesis of a 286 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan<sup>TM</sup> Panel (OriGene Technologies, Rockville, MD). This panel represented serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at (94C for 3 minutes) followed by 35 cycles of 94C for 45 seconds, 53C for 2 minutes and 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-70 was expressed in the brain, heart and lung. Within the brain, nGPCR-70 was expressed in regions including but not limited to, cerebellum, hippocampus, substantia nigra, thalamus, frontal lobe, caudate nucleus, and spinal cord. Expression of the nGPCR-70 in the brain provides an indication that modulators of nGPCR-70 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-70 modulators, including nGPCR-70 ligands and anti-nGPCR-70 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

*nGPCR-63*

Tissue specific expression of the putative nGPCR-63 was detected using a PCR-based RapidScan™ Gene Expression Panel (OriGene Technologies, Rockville, MD). The RapidScan Expression Panel is comprised of first-strand cDNAs from 12 human tissues that are serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of the putative GPCR in human tissues. Human tissues arrayed included: brain, heart, kidney, spleen, liver, colon, lung, small intestine, muscle, stomach, testis, placenta, salivary gland, thyroid, adrenal gland, pancreas, ovary, uterus, prostate, skin, PBL, bone marrow, fetal brain, fetal liver. PCR primers were designed based on the available sequence of the Celera sequence HUM\_IDS|Contig|11000258115466. The forward primer used was: 5'-ACAGCCCCAAAGCCAAACAC-3' (SEQ ID NO:162). The reverse primer was: 5'-CCGCAGGAGCAATGAAAATCAG-3' (SEQ ID NO:163). This primer set primed the synthesis of a 220 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain RapidScan™ Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System™ (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at (94 for 3minutes) followed by 35 cycles of 94C for 45 seconds, 54C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel, stained with ethidium bromide.

nGPCR-63 was expressed in the brain, heart, kidney, liver, muscle, ovary, prostate, small intestine, spleen, testis, peripheral blood leukocytes, and lung. Within the brain, nGPCR-63 was expressed in regions including but not limited to, cerebellum, amygdala, hypothalamus, medulla, temporal lobe, pons, hippocampus, substantia nigra, thalamus, frontal lobe, caudate nucleus, and spinal cord. Expression of the nGPCR-63 in the brain provides an indication that modulators of nGPCR-63 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-63 modulators, including nGPCR-63 ligands and anti-

nGPCR-63 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

*nGPCR-42*

Expression of nGPCR-42 in the various tissues was detected by using PCR primers  
5 designed based on the sequence of the receptor that prime the synthesis of a 110 bp fragment in  
the presence of the appropriate cDNA. The forward primer used to detect expression of nGPCR-  
42 was: 5'-TCTCCAAACTCCTGGCCTTC-3' (SEQ ID NO:164) and the reverse primer was:  
5'-GCAGGGCAGCTTTTTCATCC-3' (SEQ ID NO:165). Primers were synthesized by  
Genosys Corp., The Woodlands, TX. The primer set was assembled into a PCR reaction using  
10 the components of the Expand Hi-Fi PCR System™ (Roche Molecular Biochemicals). Twenty-  
five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR  
plate. The plate was placed in a GeneAmp PCR9700 PCR thermocycler (Perkin Elmer Applied  
Biosystems). The plate was then exposed to the following cycling parameters: Pre-soak 94C for  
3 minutes; denaturation at 94C for 30 seconds; annealing at primer T<sub>m</sub> for 45 seconds; extension  
15 72C for 2 minutes; for 35 cycles. PCR productions were then separated and analyzed by  
electrophoresis on a 2.0% agarose gel.

The 4-log dilution range of cDNA deposited on the plate ensured that the amplification  
reaction was within the linear range and, facilitated semi-quantitative determination of relative  
mRNA accumulation in the various tissues or brain regions examined.

20 nGPCR-42 was expressed in the brain, peripheral blood leukocytes, bone marrow,  
placenta, salivary gland, liver, ovary, uterus, testis, fetal liver, heart, thyroid gland, kidney,  
adrenal gland, spleen, pancreas, colon, lung, prostate, small intestine, skin, muscle, fetal brain,  
and stomach. Within the brain, GPCR-42 was expressed in the temporal lobe, cerebellum,  
substantia nigra, caudate nucleus, amygdala, frontal lobe, thalamus, hippocampus,  
25 hypothalamus, pons, medulla, and spinal cord. Expression of the nGPCR-42 in the brain  
provided an indication that modulators of nGPCR-42 activity have utility for treating disorders,  
including but not limited to, schizophrenia, affective disorders, metabolic disorders,  
inflammatory disorders, cancers, ADHD/ADD (i.e., Attention Deficit-Hyperactivity  
Disorder/Attention Deficit Disorder), and neural disorders such as Alzheimer's disease,  
30 Parkinson's disease, migraine, and senile dementia. Some other diseases for which modulators  
of nGPCR-42 may have utility include depression, anxiety, bipolar disease, epilepsy, neuritis,  
neurasthenia, neuropathy, neuroses, and the like. Use of nGPCR-42 modulators, including  
nGPCR-42 ligands and anti-nGPCR-42 antibodies, to treat individuals having such disease  
states is intended as an aspect of the invention.

*nGPCR-46*

Tissue specific expression of the cDNA encoding nGPCR-46 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 12 human tissues are serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array is used to generate a comprehensive expression profile of nGPCR-46 in human tissues. Human tissues arrayed include: brain, heart, kidney, peripheral blood leukocytes, liver, lung, muscle, ovary, prostate, small intestine, spleen and testis. The forward PCR primer was: 5'-ACGCCCCGCTGAACCGTATAC-3' (SEQ ID NO:166) and the reverse primer used was: 5'-GGGTGCCACCTGGTTGCTC-3' (SEQ ID NO:167). This primer set will prime the synthesis of a 242 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan<sup>TM</sup> Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at (94C for 3 minutes) followed by 35 cycles of 94C for 45 seconds, 53C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-46 was expressed in the brain, peripheral blood lymphocytes, testis, heart, kidney, spleen, prostate, ovary, liver, lung, small intestine, and muscle. Within the brain, nGPCR-46 was expressed in frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla, and spinal cord. Expression of the nGPCR-46 in the brain indicates that modulators of nGPCR-46 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-46 modulators, including nGPCR-46 ligands and anti-nGPCR-46 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

*nGPCR-48*

Tissue specific expression of the cDNA encoding nGPCR-48 was detected using a PCR-based RapidScan™ Gene Expression Panel (OriGene Technologies, Rockville, MD). The RapidScan Expression Panel is comprised of first-strand cDNAs from 24 human tissues that are serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array is used to generate a comprehensive expression profile of the putative GPCR in human tissues. Human tissues arrayed include: brain, heart, kidney, spleen, liver, colon, lung, small intestine, muscle, stomach, testis, placenta, salivary gland, thyroid, adrenal gland, pancreas, ovary, uterus, prostate, skin, PBL, bone marrow, fetal brain, fetal liver. The forward PCR primer was 5'-ATGGCACCTTCTCATCGG-3' (SEQ ID NO:168) and the reverse primer was 5'-ACGTAGTACACCGCCTTG-3' (SEQ ID NO:169). This primer set will prime the synthesis of a 392 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan™ Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System™ (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR Thermocycler (PE Applied Biosystems). The following cycling program was executed: Pre-soak at (94°C for 3min.) followed by 35 cycles of 94°C for 45 seconds, 53°C for 2 minutes, 72°C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel, stained with ethidium bromide.

The 4-log dilution range of cDNA deposited on the plate ensured that the amplification reaction was within the linear range and, facilitated semi-quantitative determination of relative mRNA accumulation in the various tissues or brain regions examined.

nGPCR-48 was expressed in the ovary, prostate, skin, stomach bone marrow, placenta, fetal liver, brain, heart, spleen, liver, colon, uterus, lung, small intestine, peripheral blood leukocytes, testis, and fetal brain. Within the brain, nGPCR-48 was expressed in the substantia nigra, frontal lobe, temporal lobe, cerebellum, hippocampus, caudate nucleus, thalamus and spinal cord. Expression of nGPCR-48 in the brain provided an indication that modulators of nGPCR-48 activity have utility for treating disorders, including but not limited to, schizophrenia, affective disorders, ADHD/ADD (i.e., Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), neural disorders such as Alzheimer's disease, Parkinson's



disease, migraine, senile dementia, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, metabolic disorders, inflammatory disorders, cancers and the like. Use of nGPCR-48 modulators, including nGPCR-48 ligands and anti-nGPCR-48 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### *nGPCR-49*

Tissue specific expression of the cDNA encoding SEQ-49 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 6 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of the putative GPCRs in human tissues. Human tissues arrayed included: brain, heart, kidney, peripheral blood leukocytes, lung, and testis. PCR primers were designed based on the available sequence of the Celera sequence GA\_11585051. The forward primer used was 5'-AGCAGGTAGATGGAGAA GG-3' (SEQ ID NO:170). The reverse primer used was 5'-GACTCCTGAGGACCATGTC- 3' (SEQ ID NO:171). This primer set primed the synthesis of a 320 base pair fragment in the presence of the appropriate cDNA. PCR reactions were performed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at 94C for 3minutes followed by 35 cycles of 94C for 45 seconds, 53.5C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-49 was expressed in the brain, heart, testis, peripheral blood leukocytes, kidney, liver, muscle, ovary, prostate, small intestine, spleen, and lung. Within the brain, nGPCR-49 was expressed in regions including but not limited to, cerebellum, hippocampus, substantia nigra, thalamus, hypothalamus, frontal lobe, temporal lobe, amygdala, pons, medulla, caudate nucleus, and spinal cord. Expression of the nGPCR-49 in the brain provides an indication that modulators of nGPCR-49 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-49 modulators, including nGPCR-49 ligands and anti-nGPCR-49 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### *nGPCR-61*

Tissue specific expression of the cDNA encoding SEQ-61 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 6 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR

plate. This array was used to generate a comprehensive expression profile of the putative GPCRs in human tissues. Human tissues arrayed included: brain, heart, kidney, peripheral blood leukocytes, lung, and testis. PCR primers were designed based on the available sequence of the Celera sequence GA\_13368549. The forward primer used was 5'-CATTGGAACCACATTTAT TGG-3' (SEQ ID NO:172). The reverse primer used was 5'-AAGCAAGACAGGTGAGAATG-3' (SEQ ID NO:173). This primer set primed the synthesis of a 284 base pair fragment in the presence of the appropriate cDNA. PCR reactions were performed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at 94C for 3 minutes followed by 35 cycles of 94C for 45 seconds, 53.5C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-61 was expressed in the brain, heart, testis, peripheral blood leukocytes, and lung. Expression of the nGPCR-61 in the brain provides an indication that modulators of nGPCR-61 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-61 modulators, including nGPCR-61 ligands and anti-nGPCR-61 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### *nGPCR-51*

Tissue specific expression of the cDNA encoding SEQ-51 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 12 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of the putative GPCR in human tissues. Human tissues arrayed include: brain, heart, kidney, peripheral blood leukocytes, liver, lung, muscle, ovary, prostate, small intestine, spleen and testis. PCR primers were designed based on the sequence of nGPCR-51 provided herein as SEQ ID NO:39 and SEQ ID NO:57. The forward primer used was 5'-CCATCTGTCTCAGCTATGCC-3' (SEQ ID NO:174), corresponding base pairs 929 through 948 of SEQ ID NO:57. The reverse primer used was 5'-TCCTTCTCAGTCGCTCTTC-3' (SEQ ID NO:175) corresponding to base pairs 1022 through 1038 of SEQ ID NO:57. This primer set primes the synthesis of a 112 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan<sup>TM</sup> Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 3 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe,

temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at 94C for 3 minutes followed by 35 cycles of 94C for 45 seconds, 52C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-51 was expressed in the brain, heart, peripheral blood leukocytes, liver, prostate, testis, lung, small intestine, and spleen. Within the brain, nGPCR-51 was expressed in the cerebellum, hippocampus, substantia nigra, thalamus, hypothalamus, pons, frontal lobe, temporal lobe, caudate nucleus, medulla, spinal cord, and amygdala. Expression of the nGPCR-51 in the brain provides an indication that modulators of nGPCR-51 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders, cancers, attention disorders, anxiety, depression, and obesity. Use of nGPCR-51 modulators, including nGPCR-51 ligands and anti-nGPCR-51 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### *nGPCR-52*

Tissue specific expression of the cDNA encoding SEQ-52 was detected using a PCR-based method. Multiple Choice<sup>TM</sup> first strand cDNAs (OriGene Technologies, Rockville, MD) from 12 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of nGPCR-52 in human tissues. Human tissues arrayed include: brain, heart, kidney, peripheral blood leukocytes, liver, lung, muscle, ovary, prostate, small intestine, spleen and testis. The forward primer used was 5'-CAAACAACAAACAGCAGAACC-3' (SEQ ID NO:176) and the reverse primer was 5'-TCACAGTCACACCTACCAAG-3' (SEQ ID NO:177). This primer set primed the synthesis of a 274 base pair fragment in the presence of the appropriate cDNA. For detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan<sup>TM</sup> Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized

by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN).

Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed; Pre-soak at 94C for 3 minutes followed by 35 cycles of 94C for 45 seconds, 53C for 2 minutes, 72C for 45 seconds. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-52 was expressed in the brain, the lungs, muscle, small intestine, spleen, testis, heart, peripheral blood leukocytes, and liver. Within the brain, nGPCR-52 was expressed in the frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, hypothalamus, and medulla. Expression of nGPCR-52 in the brain provides an indication that modulators of nGPCR-52 activity have utility for treating neurological disorders, including but not limited to, movement disorders, affective disorders, metabolic disorders, inflammatory disorders and cancers. Use of nGPCR-52 modulators, including nGPCR-52 ligands and anti-nGPCR-52 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### **Example 6: Chromosomal Localization**

##### *nGPCR-51*

The chromosomal location of the gene encoding SEQ51 was determined using the Stanford G3 Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL). This panel contains 83 radiation hybrid clones of the entire human genome created by the Stanford Human Genome Center. PCR reactions were assembled containing 25ng of DNA from each clone and the the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN) in a final reaction volume of 15 ul. PCR primers were synthesized by Genosys Corp., The Woodlands, TX. PCR primers were designed based on the available sequence of the Celera sequence GA\_11007426. The forward primer used was 5'-CCATCTGTCTCAGCTATGCC-3' (SEQ ID NO:178) corresponding base pairs 241 through 260 of GA\_11007426. The reverse primer used was 5'-TCCTTCTCAGTCGCTCTTC-3' (SEQ ID NO:179) corresponding to base pairs 334 through 352 of GA\_11007426. This primer set will prime the synthesis of a 112 base pair fragment in the presence of the appropriate genomic DNA. PCR reactions were incubated in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at 94C for 3 minutes, and 35 cycles at 94C for 30 seconds, 52C for 60 seconds, 72C for 2 minutes. PCR reaction products were then

separated and analyzed by electrophoresis on a 2.0% agarose gel, stained with ethidium bromide. Lanes were scored for the presence or absence of the expected PCR product and the results submitted to the Stanford Human Genome Center via e-mail for analysis (world wide web of the Internet at, for example, [shgc.stanford.edu/RH/rhserverformnew.html](http://shgc.stanford.edu/RH/rhserverformnew.html)). G3

- 5 Radiation Hybrid Panel Analysis of the SEQ51 places it on chromosome 6, most nearly linked to Stanford marker SHGC-34355 with a LOD score of 12.32. This marker lies at position 6q21. Chromosome 6q21-q22.3 contains a highly significant disease locus for schizophrenia (Cao *et al.*, Genomics 43: 1-8, (1997)). Genes which map to this region of the chromosome are candidate genes for schizophrenia. Any genes localized to chromosomal regions in linkage with
- 10 schizophrenia are candidate genes for disease susceptibility. Genes in these regions with the potential to play a biochemical/functional role in the disease process (like G protein coupled receptors) have a high probability of being a disease modifying locus. nGPCR-51, because of its chromosomal location, is an attractive target therefor for screening ligands useful in modulating cellular processes involved in schizophrenia.

15 *nGPCR-52*

- The chromosomal location of the gene encoding Seq-52 was determined using the Stanford G3 Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL). This panel contains 83 radiation hybrid clones of the entire human genome created by the Stanford Human Genome Center. PCR reactions were assembled containing 25ng of DNA from each clone and
- 20 the components of the Expand Hi-Fi PCR System<sup>TM</sup> (Roche Molecular Biochemicals, Indianapolis, IN) in a final reaction volume of 15 µl. PCR primers were synthesized by Genosys Corp., The Woodlands, TX. The forward primer used was RRH421: 5'-GTCTATG CTACCAAGTTCACC-3' (SEQ ID NO:180) corresponding to base pairs 3328 through 3348. The reverse primer used was RRH422: 5'-ATTCCTCCAGCCCATCATC-3' (SEQ ID NO:181)
- 25 corresponding to base pairs 3433 through 3451. This primer set will prime the synthesis of a 124 base pair fragment in the presence of the appropriate genomic DNA. PCR reactions were incubated in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at 94C for 3 minutes, and 35 cycles of 94C for 30 seconds, 55C for 60 seconds, 72C for 2 minutes. PCR reaction products were then
- 30 separated and analyzed by electrophoresis on a 2.0% agarose gel, stained with ethidium bromide. Lanes were scored for the presence or absence of the expected PCR product and the results submitted to the Stanford Human Genome Center via e-mail for analysis (world wide web of the Internet at, for example, [shgc.stanford.edu/RH/rhserverformnew.html](http://shgc.stanford.edu/RH/rhserverformnew.html)). The chromosomal location of the gene encoding nGPCR-52 was determined using the Stanford G3

Radiation Hybrid Panel (Research Genetics, Inc., Huntsville, AL). This panel contains 83 radiation hybrid clones of the entire human genome created by the Stanford Human Genome Center. Lanes were scored for the presence or absence of the expected PCR product and the results submitted to the Stanford Human Genome Center via e-mail for analysis. The analysis of nGPCR-52 places it on chromosome 6, most nearly linked to Stanford marker SHGC-1836 with a LOD score of 15.65. This marker lies at position 6q21. Chromosome 6q21-q22.3 contains a highly significant disease locus for schizophrenia (Cao *et al.*, Genomics 43: 1-8, (1997)). Genes which map to this region of the chromosome are candidate genes for schizophrenia. Any genes localized to chromosomal regions in linkage with schizophrenia are candidate genes for disease susceptibility. Genes in these regions with the potential to play a biochemical/functional role in the disease process (like G protein coupled receptors) have a high probability of being a disease modifying locus. nGPCR-52 because of its chromosomal location is an attractive target therefor for screening ligands useful in modulating cellular processes involved in schizophrenia.

#### **Example 7: Northern Blot Analysis**

Northern blots are performed to examine the expression of nGPCR-x mRNA. The sense orientation oligonucleotide and the antisense-orientation oligonucleotide, described above, are used as primers to amplify a portion of the GPCR-x cDNA sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60. Multiple human tissue northern blots from Clontech (Human II # 7767-1) are hybridized with the probe. Pre-hybridization is carried out at 42C for 4 hours in 5xSSC, 1X Denhardt's reagent, 0.1% SDS, 50% formamide, 250 mg/ml salmon sperm DNA. Hybridization is performed overnight at 42C in the same mixture with the addition of about  $1.5 \times 10^6$  cpm/ml of labeled probe. The probe is labeled with  $\alpha$ - $^{32}\text{P}$ -dCTP by Rediprime™ DNA labeling system (Amersham Pharmacia), purified on Nick Column™ (Amersham Pharmacia) and added to the hybridization solution. The filters are washed several times at 42C in 0.2x SSC, 0.1% SDS. Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with intensifying screen at -80C.

#### **Example 8: Recombinant Expression of nGPCR-x in Eukaryotic Host Cells**

##### *Expression of nGPCR-x in Mammalian Cells*

To produce nGPCR-x protein, a nGPCR-x-encoding polynucleotide is expressed in a suitable host cell using a suitable expression vector and standard genetic engineering techniques. For example, the nGPCR-x-encoding sequence described in Example 1 is subcloned into the commercial expression vector pzeoSV2 (Invitrogen, San Diego, CA) and transfected into Chinese Hamster Ovary (CHO) cells using the transfection reagent FuGENE6™ (Boehringer-Mannheim) and the transfection protocol provided in the product insert. Other

eukaryotic cell lines, including human embryonic kidney (HEK293) and COS cells, are suitable as well. Cells stably expressing nGPCR-x are selected by growth in the presence of 100 µg/ml zeocin (Stratagene, LaJolla, CA). Optionally, nGPCR-x may be purified from the cells using standard chromatographic techniques. To facilitate purification, antisera is raised against one or  
5 more synthetic peptide sequences that correspond to portions of the nGPCR-x amino acid sequence, and the antisera is used to affinity purify nGPCR-x. The nGPCR-x also may be expressed in-frame with a tag sequence (*e.g.*, polyhistidine, hemagglutinin, FLAG) to facilitate purification. Moreover, it will be appreciated that many of the uses for nGPCR-x polypeptides, such as assays described below, do not require purification of nGPCR-x from the host cell.

#### 10 *Expression of nGPCR-x in 293 cells*

For expression of nGPCR-x in mammalian cells HEK293 (transformed human, primary embryonic kidney cells), a plasmid bearing the relevant nGPCR-x coding sequence is prepared, using vector pSecTag2A (Invitrogen). Vector pSecTag2A contains the murine IgK chain leader sequence for secretion, the c-myc epitope for detection of the recombinant protein with the anti-  
15 myc antibody, a C-terminal polyhistidine for purification with nickel chelate chromatography, and a Zeocin resistant gene for selection of stable transfectants. The forward primer for amplification of this GPCR cDNA is determined by routine procedures and preferably contains a 5' extension of nucleotides to introduce the *HindIII* cloning site and nucleotides matching the GPCR sequence. The reverse primer is also determined by routine procedures and preferably  
20 contains a 5' extension of nucleotides to introduce an *XhoI* restriction site for cloning and nucleotides corresponding to the reverse complement of the nGPCR-x sequence. The PCR conditions are 55°C as the annealing temperature. The PCR product is gel purified and cloned into the *HindIII-XhoI* sites of the vector.

The DNA is purified using Qiagen chromatography columns and transfected into 293  
25 cells using DOTAP™ transfection media (Boehringer Mannheim, Indianapolis, IN). Transiently transfected cells are tested for expression after 24 hours of transfection, using western blots probed with anti-His and anti-nGPCR-x peptide antibodies. Permanently transfected cells are selected with Zeocin and propagated. Production of the recombinant protein is detected from both cells and media by western blots probed with anti-His, anti-Myc or anti-GPCR peptide  
30 antibodies.

#### *Expression of nGPCR-x in COS cells*

For expression of the nGPCR-x in COS7 cells, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60 can be cloned into vector p3-CI. This vector is a pUC18-derived plasmid that contains the HCMV (human

cytomegalovirus) promoter-intron located upstream from the bGH (bovine growth hormone) polyadenylation sequence and a multiple cloning site. In addition, the plasmid contains the dhfr (dihydrofolate reductase) gene which provides selection in the presence of the drug methotrexane (MTX) for selection of stable transformants.

5       The forward primer is determined by routine procedures and preferably contains a 5' extension which introduces an *XbaI* restriction site for cloning, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60. The reverse primer is also determined by routine procedures and preferably contains 5'-extension of nucleotides which introduces a *SalI* cloning site followed by nucleotides which  
10       correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60. The PCR consists of an initial denaturation step of 5 minutes at 95C, 30 cycles of 30 seconds denaturation at 95C, 30 seconds annealing at 58C and 30 seconds extension at 72C, followed by 5 minutes extension at 72C. The PCR product is gel purified and ligated into the *XbaI* and *SalI* sites of vector p3-CI. This construct is transformed into *E. coli*  
15       cells for amplification and DNA purification. The DNA is purified with Qiagen chromatography columns and transfected into COS 7 cells using Lipofectamine™ reagent from BRL, following the manufacturer's protocols. Forty-eight and 72 hours after transfection, the media and the cells are tested for recombinant protein expression.

20       nGPCR-x expressed from a COS cell culture can be purified by concentrating the cell-growth media to about 10 mg of protein/ml, and purifying the protein by, for example, chromatography. Purified nGPCR-x is concentrated to 0.5 mg/ml in an Amicon concentrator fitted with a YM-10 membrane and stored at -80C.

#### *Expression of nGPCR-x in Insect Cells*

25       For expression of nGPCR-x in a baculovirus system, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60 can be amplified by PCR. The forward primer is determined by routine procedures and preferably contains a 5' extension which adds the *NdeI* cloning site, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60. The reverse primer is also determined by routine procedures and preferably contains a 5'  
30       extension which introduces the *KpnI* cloning site, followed by nucleotides which correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60.

      The PCR product is gel purified, digested with *NdeI* and *KpnI*, and cloned into the corresponding sites of vector pACHTL-A (Pharmingen, San Diego, CA). The pACHTL



expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV), and a 6XHis tag upstream from the multiple cloning site. A protein kinase site for phosphorylation and a thrombin site for excision of the recombinant protein precede the multiple cloning site is also present. Of course, many other baculovirus vectors could be used in place of pAcHTL-A, such as pAc373, pVL941 and pAcIM1. Other suitable vectors for the expression of GPCR polypeptides can be used, provided that the vector construct includes appropriately located signals for transcription, translation, and trafficking, such as an in-frame AUG and a signal peptide, as required. Such vectors are described in Luckow *et al.*, Virology 170:31-39, among others. The virus is grown and isolated using standard baculovirus expression methods, such as those described in Summers *et al.* (A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Texas Agricultural Experimental Station Bulletin No. 1555 (1987)).

In a preferred embodiment, pAcHLT-A containing nGPCR-x gene is introduced into baculovirus using the "BaculoGold™" transfection kit (Pharmingen, San Diego, CA) using methods established by the manufacturer. Individual virus isolates are analyzed for protein production by radiolabeling infected cells with <sup>35</sup>S-methionine at 24 hours post infection. Infected cells are harvested at 48 hours post infection, and the labeled proteins are visualized by SDS-PAGE. Viruses exhibiting high expression levels can be isolated and used for scaled up expression.

For expression of a nGPCR-x polypeptide in a Sf9 cells, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60 can be amplified by PCR using the primers and methods described above for baculovirus expression. The nGPCR-x cDNA is cloned into vector pAcHLT-A (Pharmingen) for expression in Sf9 insect. The insert is cloned into the *NdeI* and *KpnI* sites, after elimination of an internal *NdeI* site (using the same primers described above for expression in baculovirus). DNA is purified with Qiagen chromatography columns and expressed in Sf9 cells. Preliminary Western blot experiments from non-purified plaques are tested for the presence of the recombinant protein of the expected size which reacted with the GPCR-specific antibody. These results are confirmed after further purification and expression optimization in HiG5 cells.

#### **Example 9: Interaction Trap/Two-Hybrid System**

In order to assay for nGPCR-x-interacting proteins, the interaction trap/two-hybrid library screening method can be used. This assay was first described in Fields *et al.*, *Nature*, 1989, 340, 245, which is incorporated herein by reference in its entirety. A protocol is published in Current Protocols in Molecular Biology 1999, John Wiley & Sons, NY, and Ausubel, F. M. *et*

*al.* 1992, Short protocols in molecular biology, Fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety. Kits are available from Clontech, Palo Alto, CA (Matchmaker Two-Hybrid System 3).

A fusion of the nucleotide sequences encoding all or partial nGPCR-x and the yeast transcription factor GAL4 DNA-binding domain (DNA-BD) is constructed in an appropriate plasmid (*i.e.*, pGBKT7) using standard subcloning techniques. Similarly, a GAL4 active domain (AD) fusion library is constructed in a second plasmid (*i.e.*, pGADT7) from cDNA of potential GPCR-binding proteins (for protocols on forming cDNA libraries, see Sambrook *et al.* 1989, Molecular cloning: a laboratory manual, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY), which is incorporated herein by reference in its entirety. The DNA-BD/nGPCR-x fusion construct is verified by sequencing, and tested for autonomous reporter gene activation and cell toxicity, both of which would prevent a successful two-hybrid analysis. Similar controls are performed with the AD/library fusion construct to ensure expression in host cells and lack of transcriptional activity. Yeast cells are transformed (*ca.* 105 transformants/mg DNA) with both the nGPCR-x and library fusion plasmids according to standard procedures (Ausubel *et al.*, 1992, Short protocols in molecular biology, fourth edition, Greene and Wiley-interscience, NY, which is incorporated herein by reference in its entirety). *In vivo* binding of DNA-BD/nGPCR-x with AD/library proteins results in transcription of specific yeast plasmid reporter genes (*i.e.*, lacZ, HIS3, ADE2, LEU2). Yeast cells are plated on nutrient-deficient media to screen for expression of reporter genes. Colonies are dually assayed for  $\beta$ -galactosidase activity upon growth in Xgal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside) supplemented media (filter assay for  $\beta$ -galactosidase activity is described in Breeden *et al.*, Cold Spring Harb. Symp. Quant. Biol., 1985, 50, 643, which is incorporated herein by reference in its entirety). Positive AD-library plasmids are rescued from transformants and reintroduced into the original yeast strain as well as other strains containing unrelated DNA-BD fusion proteins to confirm specific nGPCR-x/library protein interactions. Insert DNA is sequenced to verify the presence of an open reading frame fused to GAL4 AD and to determine the identity of the nGPCR-x-binding protein.

#### **Example 10: Mobility Shift DNA-Binding Assay Using Gel Electrophoresis**

A gel electrophoresis mobility shift assay can rapidly detect specific protein-DNA interactions. Protocols are widely available in such manuals as Sambrook *et al.* 1989, *Molecular cloning: a laboratory manual*, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY and Ausubel, F. M. *et al.*, 1992, *Short Protocols in Molecular Biology*, fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety.

Probe DNA(<300 bp) is obtained from synthetic oligonucleotides, restriction endonuclease fragments, or PCR fragments and end-labeled with  $^{32}\text{P}$ . An aliquot of purified nGPCR-x (ca. 15  $\mu\text{g}$ ) or crude nGPCR-x extract (ca. 15 ng) is incubated at constant temperature (in the range 22-37C) for at least 30 minutes in 10-15  $\mu\text{l}$  of buffer (*i.e.* TAE or TBE, pH 8.0-8.5) containing radiolabeled probe DNA, nonspecific carrier DNA (ca. 1  $\mu\text{g}$ ), BSA (300  $\mu\text{g}/\text{ml}$ ), and 10% (v/v) glycerol. The reaction mixture is then loaded onto a polyacrylamide gel and run at 30-35 mA until good separation of free probe DNA from protein-DNA complexes occurs. The gel is then dried and bands corresponding to free DNA and protein-DNA complexes are detected by autoradiography.

#### Example 11: Antibodies to nGPCR-x

Standard techniques are employed to generate polyclonal or monoclonal antibodies to the nGPCR-x receptor, and to generate useful antigen-binding fragments thereof or variants thereof, including "humanized" variants. Such protocols can be found, for example, in Sambrook *et al.* (1989) and Harlow *et al.* (Eds.), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988). In one embodiment, recombinant nGPCR-x polypeptides (or cells or cell membranes containing such polypeptides) are used as antigen to generate the antibodies. In another embodiment, one or more peptides having amino acid sequences corresponding to an immunogenic portion of nGPCR-x (*e.g.*, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more amino acids) are used as antigen. Peptides corresponding to extracellular portions of nGPCR-x, especially hydrophilic extracellular portions, are preferred. The antigen may be mixed with an adjuvant or linked to a hapten to increase antibody production.

##### *Polyclonal or Monoclonal antibodies*

As one exemplary protocol, recombinant nGPCR-x or a synthetic fragment thereof is used to immunize a mouse for generation of monoclonal antibodies (or larger mammal, such as a rabbit, for polyclonal antibodies). To increase antigenicity, peptides are conjugated to Keyhole Lymphet Hemocyanin (Pierce), according to the manufacturer's recommendations. For an initial injection, the antigen is emulsified with Freund's Complete Adjuvant and injected subcutaneously. At intervals of two to three weeks, additional aliquots of nGPCR-x antigen are emulsified with Freund's Incomplete Adjuvant and injected subcutaneously. Prior to the final booster injection, a serum sample is taken from the immunized mice and assayed by western blot to confirm the presence of antibodies that immunoreact with nGPCR-x. Serum from the immunized animals may be used as polyclonal antisera or used to isolate polyclonal antibodies

that recognize nGPCR-x. Alternatively, the mice are sacrificed and their spleen removed for generation of monoclonal antibodies.

To generate monoclonal antibodies, the spleens are placed in 10 ml serum-free RPMI 1640, and single cell suspensions are formed by grinding the spleens in serum-free RPMI 1640, supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units/ml penicillin, and 100 µg/ml streptomycin (RPMI) (Gibco, Canada). The cell suspensions are filtered and washed by centrifugation and resuspended in serum-free RPMI. Thymocytes taken from three naive Balb/c mice are prepared in a similar manner and used as a Feeder Layer. NS-1 myeloma cells, kept in log phase in RPMI with 10% fetal bovine serum (FBS) (Hyclone Laboratories, Inc., Logan, Utah) for three days prior to fusion, are centrifuged and washed as well.

To produce hybridoma fusions, spleen cells from the immunized mice are combined with NS-1 cells and centrifuged, and the supernatant is aspirated. The cell pellet is dislodged by tapping the tube, and 2 ml of 37C PEG 1500 (50% in 75 mM HEPES, pH 8.0) (Boehringer-Mannheim) is stirred into the pellet, followed by the addition of serum-free RPMI. Thereafter, the cells are centrifuged, resuspended in RPMI containing 15% FBS, 100 µM sodium hypoxanthine, 0.4 µM aminopterin, 16 µM thymidine (HAT) (Gibco), 25 units/ml IL-6 (Boehringer-Mannheim) and  $1.5 \times 10^6$  thymocytes/ml, and plated into 10 Corning flat-bottom 96-well tissue culture plates (Corning, Corning New York).

On days 2, 4, and 6 after the fusion, 100 µl of medium is removed from the wells of the fusion plates and replaced with fresh medium. On day 8, the fusions are screened by ELISA, testing for the presence of mouse IgG that binds to nGPCR-x. Selected fusion wells are further cloned by dilution until monoclonal cultures producing anti-nGPCR-x antibodies are obtained.

#### *Humanization of anti-nGPCR-x monoclonal antibodies*

The expression pattern of nGPCR-x as reported herein and the proven track record of GPCRs as targets for therapeutic intervention suggest therapeutic indications for nGPCR-x inhibitors (antagonists). nGPCR-x-neutralizing antibodies comprise one class of therapeutics useful as nGPCR-x antagonists. Following are protocols to improve the utility of anti-nGPCR-x monoclonal antibodies as therapeutics in humans by "humanizing" the monoclonal antibodies to improve their serum half-life and render them less immunogenic in human hosts (*i.e.*, to prevent human antibody response to non-human anti-nGPCR-x antibodies).

The principles of humanization have been described in the literature and are facilitated by the modular arrangement of antibody proteins. To minimize the possibility of binding complement, a humanized antibody of the IgG4 isotype is preferred.

For example, a level of humanization is achieved by generating chimeric antibodies comprising the variable domains of non-human antibody proteins of interest with the constant domains of human antibody molecules. (See, *e.g.*, Morrison *et al.*, *Adv. Immunol.*, 44:65-92 (1989)). The variable domains of nGPCR-x-neutralizing anti-nGPCR-x antibodies are cloned  
5 from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from the hybridoma of interest. The V region gene fragments are linked to exons encoding human antibody constant domains, and the resultant construct is expressed in suitable mammalian host cells (*e.g.*, myeloma or CHO cells).

To achieve an even greater level of humanization, only those portions of the variable  
10 region gene fragments that encode antigen-binding complementarity determining regions ("CDR") of the non-human monoclonal antibody genes are cloned into human antibody sequences. (See, *e.g.*, Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-327 (1988); Verhoeyen *et al.*, *Science* 239:1534-36 (1988); and Tempest *et al.*, *Bio/Technology* 9: 266-71 (1991)). If necessary, the  $\beta$ -sheet framework of the human antibody  
15 surrounding the CDR3 regions also is modified to more closely mirror the three dimensional structure of the antigen-binding domain of the original monoclonal antibody. (See Kettleborough *et al.*, *Protein Engin.*, 4:773-783 (1991); and Foote *et al.*, *J. Mol. Biol.*, 224:487-499 (1992)).

In an alternative approach, the surface of a non-human monoclonal antibody of interest  
20 is humanized by altering selected surface residues of the non-human antibody, *e.g.*, by site-directed mutagenesis, while retaining all of the interior and contacting residues of the non-human antibody. See Padlan, *Molecular Immunol.*, 28(4/5):489-98 (1991).

The foregoing approaches are employed using nGPCR-x-neutralizing anti-nGPCR-x monoclonal antibodies and the hybridomas that produce them to generate humanized nGPCR-x  
25 x-neutralizing antibodies useful as therapeutics to treat or palliate conditions wherein nGPCR-x expression or ligand-mediated nGPCR-x signaling is detrimental.

#### *Human nGPCR-x-Neutralizing Antibodies from Phage Display*

Human nGPCR-x-neutralizing antibodies are generated by phage display techniques such as those described in Aujame *et al.*, *Human Antibodies* 8(4):155-168 (1997);  
30 Hoogenboom, *TIBTECH* 15:62-70 (1997); and Rader *et al.*, *Curr. Opin. Biotechnol.* 8:503-508 (1997), each of which is incorporated herein by reference in its entirety. For example, antibody variable regions in the form of Fab fragments or linked single chain Fv fragments are fused to the amino terminus of filamentous phage minor coat protein pIII. Expression of the fusion protein and incorporation thereof into the mature phage coat results in phage particles that

present an antibody on their surface and contain the genetic material encoding the antibody. A phage library comprising such constructs is expressed in bacteria, and the library is screened for nGPCR-x-specific phage-antibodies using labeled or immobilized nGPCR-x as antigen-probe.

*Human nGPCR-x-neutralizing antibodies from transgenic mice*

- 5 Human nGPCR-x-neutralizing antibodies are generated in transgenic mice essentially as described in Bruggemann *et al.*, Immunol. Today 17(8):391-97 (1996) and Bruggemann *et al.*, Curr. Opin. Biotechnol. 8:455-58 (1997). Transgenic mice carrying human V-gene segments in germline configuration and that express these transgenes in their lymphoid tissue are immunized with a nGPCR-x composition using conventional immunization protocols. Hybridomas are  
10 generated using B cells from the immunized mice using conventional protocols and screened to identify hybridomas secreting anti-nGPCR-x human antibodies (e.g., as described above).

**Example 12: Assays to Identify Modulators of nGPCR-x Activity**

- Set forth below are several nonlimiting assays for identifying modulators (agonists and antagonists) of nGPCR-x activity. Among the modulators that can be identified by these assays  
15 are natural ligand compounds of the receptor; synthetic analogs and derivatives of natural ligands; antibodies, antibody fragments, and/or antibody-like compounds derived from natural antibodies or from antibody-like combinatorial libraries; and/or synthetic compounds identified by high-throughput screening of libraries; and the like. All modulators that bind nGPCR-x are useful for identifying nGPCR-x in tissue samples (e.g., for diagnostic purposes, pathological  
20 purposes, and the like). Agonist and antagonist modulators are useful for up-regulating and down-regulating nGPCR-x activity, respectively, to treat disease states characterized by abnormal levels of nGPCR-x activity. The assays may be performed using single putative modulators, and/or may be performed using a known agonist in combination with candidate antagonists (or visa versa).

25 A. *cAMP Assays*

- In one type of assay, levels of cyclic adenosine monophosphate (cAMP) are measured in nGPCR-x-transfected cells that have been exposed to candidate modulator compounds. Protocols for cAMP assays have been described in the literature. (See, e.g., Sutherland *et al.*, Circulation 37: 279 (1968); Frandsen *et al.*, Life Sciences 18: 529-541 (1976); Dooley *et al.*,  
30 Journal of Pharmacology and Experimental Therapeutics 283 (2): 735-41 (1997); and George *et al.*, Journal of Biomolecular Screening 2 (4): 235-40 (1997)). An exemplary protocol for such an assay, using an Adenylyl Cyclase Activation FlashPlate® Assay from NEN™ Life Science Products, is set forth below.

Briefly, the nGPCR-x coding sequence (*e.g.*, a cDNA or intronless genomic DNA) is subcloned into a commercial expression vector, such as pzeoSV2 (Invitrogen), and transiently transfected into Chinese Hamster Ovary (CHO) cells using known methods, such as the transfection protocol provided by Boehringer-Mannheim when supplying the FuGENE 6 transfection reagent. Transfected CHO cells are seeded into 96-well microplates from the FlashPlate® assay kit, which are coated with solid scintillant to which antisera to cAMP has been bound. For a control, some wells are seeded with wild type (untransfected) CHO cells. Other wells in the plate receive various amounts of a cAMP standard solution for use in creating a standard curve.

One or more test compounds (*i.e.*, candidate modulators) are added to the cells in each well, with water and/or compound-free medium/diluent serving as a control or controls. After treatment, cAMP is allowed to accumulate in the cells for exactly 15 minutes at room temperature. The assay is terminated by the addition of lysis buffer containing [<sup>125</sup>I]-labeled cAMP, and the plate is counted using a Packard Topcount™ 96-well microplate scintillation counter. Unlabeled cAMP from the lysed cells (or from standards) and fixed amounts of [<sup>125</sup>I]-cAMP compete for antibody bound to the plate. A standard curve is constructed, and cAMP values for the unknowns are obtained by interpolation. Changes in intracellular cAMP levels of cells in response to exposure to a test compound are indicative of nGPCR-x modulating activity. Modulators that act as agonists of receptors which couple to the G<sub>s</sub> subtype of G proteins will stimulate production of cAMP, leading to a measurable 3-10 fold increase in cAMP levels. Agonists of receptors which couple to the G<sub>i/o</sub> subtype of G proteins will inhibit forskolin-stimulated cAMP production, leading to a measurable decrease in cAMP levels of 50-100%. Modulators that act as inverse agonists will reverse these effects at receptors that are either constitutively active or activated by known agonists.

#### nGPCR-51

Modulators that act as agonists at receptors which couple to the G<sub>s</sub> subtype of G proteins will activate adenylyl cyclase leading to a 3-10 fold increase in cyclic adenosine monophosphate (cAMP). Compounds to be tested for the ability to activate nGPCR-51 were assayed for cAMP using an Adenylyl Cyclase Activation FlashPlate® Assay from NEN™ Life Science Products.

Briefly, nGPCR-51 cDNA is subcloned into the commercial expression vector pCMVSPORT (Gibco/Life Technologies) and transiently transfected into CHO or COS 7 cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert. 24 hours post transfection the cells are harvested by dislodging from the culture flask using Versene (Gibco/BRL). The cells are counted and prepared as a suspension in

a buffer included in the assay kit that contains the phosphodiesterase inhibitor isobutylmethylxanthine. The assay is conducted in a special 96 well microplate included in the kit which is coated with solid scintillant to which antisera to cAMP has been bound. Dilutions of test compounds to be tested for activation of nGPCR-51 are added to assay wells. Several wells  
5 on the plate receive various amounts of cAMP standard solution. After the addition of cells transiently expressing nGPCR-51, cAMP is allowed to accumulate for exactly 15 minutes at room temperature. The assay is terminated by the addition of lysis buffer containing [ $^{125}$ I] cAMP, and the plate is covered and allowed to incubate at room temperature for 2-24 hours. The plate is then counted using a Packard Topcount™ 96-well microplate scintillation counter.  
10 Unlabelled cAMP from cells (or standards) competes with fixed amounts of [ $^{125}$ I] cAMP for antibody bound to the plate. A standard curve is constructed and cAMP values for the unknowns are obtained by interpolation. Data were analyzed using GraphPad Prism (San Diego, CA). There was no increase in cAMP observed after stimulation with the test compounds at concentrations up to 10  $\mu$ M.

15 *B. Aequorin Assays*

In another assay, cells (e.g., CHO cells) are transiently co-transfected with both a nGPCR-x expression construct and a construct that encodes the photoprotein apoaequorin. In the presence of the cofactor coelenterazine, apoaequorin will emit a measurable luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium. (See generally, Cobbold,  
20 et al. "Aequorin measurements of cytoplasmic free calcium," In: McCormack J.G. and Cobbold P.H., eds., Cellular Calcium: A Practical Approach. Oxford:IRL Press (1991); Stables et al., Analytical Biochemistry 252: 115-26 (1997); and Haugland, Handbook of Fluorescent Probes and Research Chemicals, Sixth edition. Eugene OR: Molecular Probes (1996).)

In one exemplary assay, nGPCR-x is subcloned into the commercial expression vector pzeoSV2 (Invitrogen) and transiently co-transfected along with a construct that encodes the  
25 photoprotein apoaequorin (Molecular Probes, Eugene, OR) into CHO cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert.

The cells are cultured for 24 hours at 37°C in MEM (Gibco/BRL, Gaithersburg, MD)  
30 supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin, at which time the medium is changed to serum-free MEM containing 5  $\mu$ M coelenterazine (Molecular Probes, Eugene, OR). Culturing is then continued for two additional hours at 37°C. Subsequently, cells are detached from the plate using VERSEN (Gibco/BRL), washed, and resuspended at 200,000 cells/ml in serum-free MEM.



Dilutions of candidate nGPCR-x modulator compounds are prepared in serum-free MEM and dispensed into wells of an opaque 96-well assay plate at 50  $\mu$ l/well. Plates are then loaded onto an MLX microtiter plate luminometer (Dynex Technologies, Inc., Chantilly, VA). The instrument is programmed to dispense 50  $\mu$ l cell suspensions into each well, one well at a time, and immediately read luminescence for 15 seconds. Dose-response curves for the candidate modulators are constructed using the area under the curve for each light signal peak. Data are analyzed with SlideWrite, using the equation for a one-site ligand, and EC<sub>50</sub> values are obtained. Changes in luminescence caused by the compounds are considered indicative of modulatory activity. Modulators that act as agonists at receptors which couple to the G<sub>q</sub> subtype of G proteins give an increase in luminescence of up to 100 fold. Modulators that act as inverse agonists will reverse this effect at receptors that are either constitutively active or activated by known agonists.

#### *nGPCR-51*

Agonist activation of receptors that couple to the G<sub>q</sub> subtype of G proteins will lead to the release of intracellular calcium. The photoprotein aequorin emits a characteristic luminescence in the presence of calcium and may be expressed in cells along with the receptor of interest in order to report agonist signalling. Peptide A was tested for the ability to activate nGPCR-51 using an assay for aequorin. Peptide A, a cyclic peptide via a disulfide bond between the Cys residues, has the following amino acid sequence: Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys Trp Gln Val (SEQ ID NO:183). Preferably, Peptide A is used in a solution containing 0.17M acetic acid, or the like, and a stabilizer such as a 0.2% BSA:acetonitrile 90:10. Briefly, nGPCR-51 cDNA is subcloned into the commercial expression vector pCMVSPORT (Gibco/Life Technologies) and transiently transfected along with the expression construct mtAequorin (Molecular Probes, Eugene, OR) into COS 7 cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection protocol provided in the product insert. 24 hours post transfection the cells are harvested by dislodging from the culture flask using Versene (Gibco/BRL) and prepared as a suspension in assay buffer (Dulbecco's Modified Eagle's Medium with high glucose, pyridoxine HCl, L-glutamine, sodium pyruvate, and 0.1% fetal bovine serum (Gibco/BRL)) and containing the cofactor coelenterazine (Molecular Probes). The cell suspension is incubated for 4 hours at room temperature with gentle stirring. After the coelenterazine loading incubation, the cells are counted and diluted to 1 x 10<sup>6</sup> cells/ml in assay buffer. Dilutions of test compound are prepared in assay buffer and pipetted into wells of an opaque 96-well assay plate, 50  $\mu$ l/well. Plates are loaded onto an MLX microtiter plate luminometer (Dynex Technologies, Inc., Chantilly, VA). The instrument is

programmed to dispense 50  $\mu$ l cell suspension into each well, one well at a time, and immediately read luminescence for 20 seconds. Dose response curves are constructed using the area under the curve for each light signal peak. Data are analyzed with SlideWrite, using the equation  $y = a^0 * a^1 / (x + a^1) + a^2$  where  $a^0$  is the maximum response minus the baseline;  $a^1$  is the  $EC_{50}$ ; and  $a^2$  is the maximum response.

Peptide A stimulation of nGPCR-51 expressing COS 7 cells resulted in a dose dependent increase in aequorin luminescence with an  $EC_{50}$  of approximately 40 nM and a maximum response of 52-fold over baseline.

### C. Luciferase Reporter Gene Assay

The photoprotein luciferase provides another useful tool for assaying for modulators of nGPCR-x activity. Cells (e.g., CHO cells or COS 7 cells) are transiently co-transfected with both a nGPCR-x expression construct (e.g., nGPCR-x in pzeoSV2) and a reporter construct which includes a gene for the luciferase protein downstream from a transcription factor binding site, such as the cAMP-response element (CRE), AP-1, or NF-kappa B. Agonist binding to receptors coupled to the  $G_s$  subtype of G proteins leads to increases in cAMP, thereby activating the CRE transcription factor and resulting in expression of the luciferase gene. Agonist binding to receptors coupled to the  $G_q$  subtype of G protein leads to production of diacylglycerol that activates protein kinase C, which activates the AP-1 or NF-kappa B transcription factors, in turn resulting in expression of the luciferase gene. Expression levels of luciferase reflect the activation status of the signaling events. (See generally, George *et al.*, *Journal of Biomolecular Screening*, 2(4): 235-240 (1997); and Stratowa *et al.*, *Current Opinion in Biotechnology* 6: 574-581 (1995)). Luciferase activity may be quantitatively measured using, e.g., luciferase assay reagents that are commercially available from Promega (Madison, WI).

In one exemplary assay, CHO cells are plated in 24-well culture dishes at a density of 100,000 cells/well one day prior to transfection and cultured at 37C in MEM (Gibco/BRL) supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin. Cells are transiently co-transfected with both a nGPCR-x expression construct and a reporter construct containing the luciferase gene. The reporter plasmids CRE-luciferase, AP-1-luciferase and NF-kappaB-luciferase may be purchased from Stratagene (LaJolla, CA). Transfections are performed using the FuGENE 6 transfection reagent (Boehringer-Mannheim) according to the supplier's instructions. Cells transfected with the reporter construct alone are used as a control. Twenty-four hours after transfection, cells are washed once with PBS pre-warmed to 37C. Serum-free MEM is then added to the cells either alone (control) or with one or more candidate modulators and the cells are incubated at 37C for five hours. Thereafter,

cells are washed once with ice-cold PBS and lysed by the addition of 100  $\mu$ l of lysis buffer per well from the luciferase assay kit supplied by Promega. After incubation for 15 minutes at room temperature, 15  $\mu$ l of the lysate is mixed with 50  $\mu$ l of substrate solution (Promega) in an opaque-white, 96-well plate, and the luminescence is read immediately on a Wallace model  
5 1450 MicroBeta scintillation and luminescence counter (Wallace Instruments, Gaithersburg, MD).

Differences in luminescence in the presence versus the absence of a candidate modulator compound are indicative of modulatory activity. Receptors that are either constitutively active or activated by agonists typically give a 3 to 20-fold stimulation of luminescence compared to cells  
10 transfected with the reporter gene alone. Modulators that act as inverse agonists will reverse this effect.

#### *nGPCR-51*

nGPCR-51 was transfected in N7 cells, stable HOS (human osteosarcoma) cells expressing NFkB-Luciferase reporter gene. Briefly, LipofectAMINE, PLUS reagent, and 4  $\mu$ g  
15 DNA were added to cells in Opti-MEM. After 4 hours, serum-containing medium was added to each plate. Twenty-four hour after transfection the cells were trypsinized and seeded in COSTAR white 96-well plate at a cell density of about 20,000 cells per well. Another 24 hours later the culture medium was replaced with fresh growth medium and Peptide A was added. After 5-hr incubation with Peptide A, the medium was aspirated and 100  $\mu$ L diluted Steady Glo  
20 was added. The plates were gently shake at room temperature for 30 minutes and luminescence determined by TopCount at the SPC mode.

The dose-response effect of Peptide A on luciferase expression in Seq51-transfected N7 cells was tested in 2 separate experiments. The EC<sub>50</sub> value of Peptide A for the 2 experiments was 1.6 nM and 1.7 nM, respectively. Salmon Peptide B (Asp Thr Met Arg Cys Met Val Gly  
25 Arg Val Tyr Arg Pro Cys Trp Glu Val; SEQ ID NO:184), a variant of Peptide A, and another variant of Peptide A were also tested in the second experiment. The EC<sub>50</sub> value for salmon Peptide B was approximately 25 nM while variant of Peptide A had no effect on these Seq51-transfected cells at concentration up to 1  $\mu$ M.

Peptide A, a cyclic heptadecapeptide, is a prominent hypothalamic neuropeptide  
30 expressed in cells which are part of an extensive network of neurons projecting to multiple parts of the brain. Despite the large numbers of hypothalamic Peptide A-expressing cells, the Peptide A cell groups are not part of the major hypothalamic nuclei (paraventricular, supraoptic, dorsomedial, ventromedial, arcuate, lateral tuberal or mammillary) (Knigge *et al.*, *J. Peptides*, 1996, 17, 1063-1073), but they have terminal projections to most of these hypothalamic nuclei.

Additionally, Peptide A projections from the hypothalamus to the rest of the brain are extensive, including to the cerebellar cortex, subcortical nuclei, limbic areas, thalamus, brain stem and spinal cord.

Peptide A projections are included in the circuitry involved in processing visual and auditory information, suggesting a general role in arousal and sensorimotor integration (Hervieu *et al.*, *Eur. J. Neurosci.*, 2000, 12, 1194-1216). Intracerebroventricular injection of Peptide A causes a decrease in the conditioning amplitude of the second of a pair of auditory evoked potentials recorded from the hippocampus (hippocampal auditory gating paradigm) and reverses the increases in the conditioning amplitude induced by alpha MSH (Miller *et al.*, *Peptides*, 1993, 14, 431-440). These data would suggest that Peptide A inhibits the normal auditory gating process. Schizophrenics also present with a loss of auditory gating which may contribute to an inability to filter inappropriate information. Dysregulation of Peptide A may contribute to this phenotype in humans and antagonists active at the Peptide A receptors may effectively treat these deficits in schizophrenia.

Major portions of the Peptide A cell group are in areas considered to be part of the extrapyramidal motor circuits. It has been suggested that this location would afford the Peptide A neuronal system the capability of coordinating hypothalamic visceral activity with appropriate motor activity (Knigge *et al.*, 1996).

Peptide A localization in the hippocampal formation, the amygdaloid regions, the bed nucleus of the stria terminalis all suggest a role in cognition and learning and in support of this Peptide A has been shown to alter passive avoidance performance (McBride *et al.*, *Peptides*, 1994, 15, 757-759).

Central injections of Peptide A in rats leads to a reduction in peak circadian ACTH levels and inhibit stress-induced ACTH secretion (Bluet Pajot *et al.*, *J. Neuroendocrinol.*, 1995, 7, 297-303; and Ludwig *et al.*, *Am. J. Physiology*, 1998, 274, E627-E633). However, injections of Peptide A into the medial preoptic area of the hypothalamus of rats result in decreases in time spent in open arms in the elevated plus maze assay indicative of increased anxiety. Injections of the peptide also stimulated female sexual receptivity (Gonzalez *et al.*, *Peptides*, 1996, 17, 171-177).

The lateral hypothalamus has a prominent role in feeding-related behaviors and in fact Peptide A has been shown to be involved in energy balance and food intake. Ob/ob mice express more Peptide A mRNA compared with lean littermates (Qu *et al.*, *Nature*, 1996, 380, 243-247). Additionally, fasting increases hypothalamic Peptide A mRNA levels (Qu *et al.*, 1996).

Intracerebroventricular injection of Peptide A leads to an increase in food intake in rats (Qu *et al.*, *Endocrinology*, 1997, 138, 351-355).

Targeted deletion of the gene which encodes the preprohormone for Peptide A resulted in mice which have reduced body weight and are lean due to reduced feeding. Additionally  
5 these mice have an increased metabolic rate in the presence of decreased leptin levels and POMC levels (Shimada *et al.*, *Nature*, 1998, 396, 670-673). These mice also have a loss of expression of the NEI peptide as well as Peptide A. Nevertheless, the prominent phenotypic feature of these mice is their reduced size. Antagonists of Peptide A receptors may be effective in the treatment of obesity.

10 The SLC-1 gene product was recently identified as a G-protein coupled receptor which is potently activated by Peptide A (Chambers *et al.*, *Nature*, 1999, 400, 261-265; and Saito *et al.*, *Nature*, 1999, 400, 265-269). nGPCR-51 has been identified as a gene encoding a second Peptide A receptor which is also potently activated by Peptide A. Like that of the SLC-1 gene, expression of nGPCR-51 is extensive throughout the brain. The multiple actions of Peptide A  
15 could be differentially mediated by each of the two Peptide A receptors. Selective agonism or antagonism at nGPCR-51 could result in a therapeutic effective in treating schizophrenia, attention disorders, anxiety, depression, obesity, and the like.

#### D. Intracellular calcium measurement using FLIPR

Changes in intracellular calcium levels are another recognized indicator of G protein-coupled receptor activity, and such assays can be employed to screen for modulators of nGPCR-  
20 x activity. For example, CHO cells stably transfected with a nGPCR-x expression vector are plated at a density of  $4 \times 10^4$  cells/well in Packard black-walled, 96-well plates specially designed to discriminate fluorescence signals emanating from the various wells on the plate. The cells are incubated for 60 minutes at 37C in modified Dulbecco's PBS (D-PBS) containing 36  
25 mg/L pyruvate and 1 g/L glucose with the addition of 1% fetal bovine serum and one of four calcium indicator dyes (Fluo-3<sup>TM</sup> AM, Fluo-4<sup>TM</sup> AM, Calcium Green<sup>TM</sup>-1 AM, or Oregon Green<sup>TM</sup> 488 BAPTA-1 AM), each at a concentration of 4  $\mu$ M. Plates are washed once with modified D-PBS without 1% fetal bovine serum and incubated for 10 minutes at 37C to remove residual dye from the cellular membrane. In addition, a series of washes with modified D-PBS  
30 without 1% fetal bovine serum is performed immediately prior to activation of the calcium response.

A calcium response is initiated by the addition of one or more candidate receptor agonist compounds, calcium ionophore A23187 (10  $\mu$ M; positive control), or ATP (4  $\mu$ M; positive control). Fluorescence is measured by Molecular Device's FLIPR with an argon laser (excitation

at 488 nm). (See, e.g., Kuntzweiler et al., Drug Development Research, 44(1):14-20 (1998)).

The F-stop for the detector camera was set at 2.5 and the length of exposure was

0.4 milliseconds. Basal fluorescence of cells was measured for 20 seconds prior to addition of

candidate agonist, ATP, or A23187, and the basal fluorescence level was subtracted from the

5 response signal. The calcium signal is measured for approximately 200 seconds, taking readings

every two seconds. Calcium ionophore A23187 and ATP increase the calcium signal 200%

above baseline levels. In general, activated GPCRs increase the calcium signal approximately

10-15% above baseline signal.

#### *nGPCR-51*

10 HEK293 cells were transiently transfected with an expression vector for nGPCR-51 and empty vector using Lipofectamine plus (Gibco) according to the manufacturer's instructions.

The next day, the cells were seeded into 96-well plates at 25,000 cells per well. The following

day, cells were loaded with 1  $\mu$ M Fluo-4-acetoxymethyl fluorescent indicator dye (Molecular Probes) in MEM (minimal essential media) containing 0.1% bovine serum albumin, 0.04%

15 pluronic acid and 2.5 mM probenecid for 30 minutes at 37C. The cells were washed with pre-

warmed (37C) assay buffer (Hanks buffer containing 15 mM HEPES, 2.5 mM probenecid and 0.1% bovine serum albumin). Assay buffer (100  $\mu$ l) was added to each well and plates were

incubated at 37C for 15 minutes. Various concentrations (0.03 pM-10 nM) of human Peptide A or salmon Peptide B were added and fluorescence produced by fluo-4 (a calcium sensitive dye)

20 was measured every second for 150 seconds on a fluorometric imaging plate reader (FLIPR1;

Molecular Devices). Human Peptide A and salmon Peptide B-induced calcium mobilization in

nGPCR-51-transfected cells with EC<sub>50</sub> values of 3 nM and 6 nM, respectively. Human Peptide

A and salmon Peptide B had no effect on vector-transfected cells.

#### *E. Mitogenesis Assay*

25 In a mitogenesis assay, the ability of candidate modulators to induce or inhibit nGPCR-x-mediated cell division is determined. (See, e.g., Lajiness et al., Journal of Pharmacology and Experimental Therapeutics 267(3): 1573-1581 (1993)). For example, CHO cells stably

expressing nGPCR-x are seeded into 96-well plates at a density of 5000 cells/well and grown at 37°C in MEM with 10% fetal calf serum for 48 hours, at which time the cells are rinsed twice

30 with serum-free MEM. After rinsing, 80  $\mu$ l of fresh MEM, or MEM containing a known

mitogen, is added along with 20  $\mu$ l MEM containing varying concentrations of one or more

candidate modulators or test compounds diluted in serum-free medium. As controls, some wells on each plate receive serum-free medium alone, and some receive medium containing 10% fetal

bovine serum. Untransfected cells or cells transfected with vector alone also may serve as controls.

After culture for 16-18 hours, 1  $\mu$ Ci of [ $^3$ H]-thymidine (2 Ci/mmol) is added to the wells and cells are incubated for an additional 2 hours at 37C. The cells are trypsinized and collected on filter mats with a cell harvester (Tomtec); the filters are then counted in a Betaplate counter. The incorporation of [ $^3$ H]-thymidine in serum-free test wells is compared to the results achieved in cells stimulated with serum (positive control). Use of multiple concentrations of test compounds permits creation and analysis of dose-response curves using the non-linear, least squares fit equation:  $A = B \times [C / (D + C)] + G$  where A is the percent of serum stimulation; B is the maximal effect minus baseline; C is the  $EC_{50}$ ; D is the concentration of the compound; and G is the maximal effect. Parameters B, C and G are determined by Simplex optimization.

Agonists that bind to the receptor are expected to increase [ $^3$ H]-thymidine incorporation into cells, showing up to 80% of the response to serum. Antagonists that bind to the receptor will inhibit the stimulation seen with a known agonist by up to 100%.

#### F. [ $^{35}$ S]GTP $\gamma$ S Binding Assay

Because G protein-coupled receptors signal through intracellular G proteins whose activity involves GTP binding and hydrolysis to yield bound GDP, measurement of binding of the non-hydrolyzable GTP analog [ $^{35}$ S]GTP $\gamma$ S in the presence and absence of candidate modulators provides another assay for modulator activity. (See, e.g., Kowal et al.,

Neuropharmacology 37:179-187 (1998).)

In one exemplary assay, cells stably transfected with a nGPCR-x expression vector are grown in 10 cm tissue culture dishes to subconfluence, rinsed once with 5 ml of ice-cold  $Ca^{2+}$ /Mg $^{2+}$ -free phosphate-buffered saline, and scraped into 5 ml of the same buffer. Cells are pelleted by centrifugation (500 x g, 5 minutes), resuspended in TEE buffer (25 mM Tris, pH 7.5, 5 mM EDTA, 5 mM EGTA), and frozen in liquid nitrogen. After thawing, the cells are homogenized using a Dounce homogenizer (one ml TEE per plate of cells), and centrifuged at 1,000 x g for 5 minutes to remove nuclei and unbroken cells.

The homogenate supernatant is centrifuged at 20,000 x g for 20 minutes to isolate the membrane fraction, and the membrane pellet is washed once with TEE and resuspended in binding buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 10 mM MgCl $_2$ , 1 mM EDTA). The resuspended membranes can be frozen in liquid nitrogen and stored at -70C until use.

Aliquots of cell membranes prepared as described above and stored at -70C are thawed, homogenized, and diluted into buffer containing 20 mM HEPES, 10 mM MgCl $_2$ , 1 mM EDTA, 120 mM NaCl, 10  $\mu$ M GDP, and 0.2 mM ascorbate, at a concentration of 10-50  $\mu$ g/ml. In a final

volume of 90  $\mu$ l, homogenates are incubated with varying concentrations of candidate modulator compounds or 100  $\mu$ M GTP for 30 minutes at 30C and then placed on ice. To each sample, 10  $\mu$ l guanosine 5'-O-(3[ $^{35}$ S]thio) triphosphate (NEN, 1200 Ci/mmol; [ $^{35}$ S]-GTP $\gamma$ S), was added to a final concentration of 100-200 pM. Samples are incubated at 30C for an additional 30 minutes, 1  
5 ml of 10mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>, at 4C is added and the reaction is stopped by filtration.

Samples are filtered over Whatman GF/B filters and the filters are washed with 20 ml ice-cold 10 mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>. Filters are counted by liquid scintillation spectroscopy. Nonspecific binding of [ $^{35}$ S]-GTP $\gamma$ S is measured in the presence of 100  $\mu$ M GTP  
10 and subtracted from the total. Compounds are selected that modulate the amount of [ $^{35}$ S]-GTP $\gamma$ S binding in the cells, compared to untransfected control cells. Activation of receptors by agonists gives up to a five-fold increase in [ $^{35}$ S]GTP $\gamma$ S binding. This response is blocked by antagonists.

G. *MAP Kinase Activity Assay*

Evaluation of MAP kinase activity in cells expressing a GPCR provides another assay to  
15 identify modulators of GPCR activity. (See, e.g., Lajiness et al., Journal of Pharmacology and Experimental Therapeutics 267(3):1573-1581 (1993) and Boulton et al., Cell 65:663-675 (1991).)

In one embodiment, CHO cells stably transfected with nGPCR-x are seeded into 6-well plates at a density of 70,000 cells/well 48 hours prior to the assay. During this 48-hour period,  
20 the cells are cultured at 37C in MEM medium supplemented with 10% fetal bovine serum, 2mM glutamine, 10 U/ml penicillin and 10 $\mu$ g/ml streptomycin. The cells are serum-starved for 1-2 hours prior to the addition of stimulants.

For the assay, the cells are treated with medium alone or medium containing either a candidate agonist or 200 nM Phorbol ester- myristoyl acetate (i.e., PMA, a positive control), and  
25 the cells are incubated at 37C for varying times. To stop the reaction, the plates are placed on ice, the medium is aspirated, and the cells are rinsed with 1 ml of ice-cold PBS containing 1mM EDTA. Thereafter, 200 $\mu$ l of cell lysis buffer (12.5 mM MOPS, pH 7.3, 12.5 mM glycerophosphate, 7.5mM MgCl<sub>2</sub>, 0.5mM EGTA, 0.5 mM sodium vanadate, 1mM benzamidine, 1mM dithiothreitol, 10  $\mu$ g/ml leupeptin, 10  $\mu$ g/ml aprotinin, 2 $\mu$ g/ml pepstatin A, and 1 $\mu$ M  
30 okadaic acid) is added to the cells. The cells are scraped from the plates and homogenized by 10 passages through a 23 3/4 G needle, and the cytosol fraction is prepared by centrifugation at 20,000 x g for 15 minutes.

Aliquots (5-10  $\mu$ l containing 1-5  $\mu$ g protein) of cytosol are mixed with 1 mM MAPK Substrate Peptide (APRTPGGRR; SEQ ID NO:182), Upstate Biotechnology, Inc., N.Y.) and



50  $\mu$ M [ $\gamma$ - $^{32}$ P]ATP (NEN, 3000 Ci/mmol), diluted to a final specific activity of ~2000 cpm/pmol, in a total volume of 25  $\mu$ l. The samples are incubated for 5 minutes at 30C, and reactions are stopped by spotting 20  $\mu$ l on 2 cm<sup>2</sup> squares of Whatman P81 phosphocellulose paper. The filter squares are washed in 4 changes of 1% H<sub>3</sub>PO<sub>4</sub>, and the squares are subjected to liquid

scintillation spectroscopy to quantitate bound label. Equivalent cytosolic extracts are incubated without MAPK substrate peptide, and the bound label from these samples are subtracted from the matched samples with the substrate peptide. The cytosolic extract from each well is used as a separate point. Protein concentrations are determined by a dye binding protein assay (Bio-Rad Laboratories). Agonist activation of the receptor is expected to result in up to a five-fold increase in MAPK enzyme activity. This increase is blocked by antagonists.

#### H. [ $^3$ H]Arachidonic Acid Release

The activation of GPCRs also has been observed to potentiate arachidonic acid release in cells, providing yet another useful assay for modulators of GPCR activity. (See, e.g., Kanterman et al., Molecular Pharmacology 39:364-369 (1991).) For example, CHO cells that are stably transfected with a nGPCR-x expression vector are plated in 24-well plates at a density of 15,000 cells/well and grown in MEM medium supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin for 48 hours at 37C before use. Cells of each well are labeled by incubation with [ $^3$ H]-arachidonic acid (Amersham Corp., 210 Ci/mmol) at 0.5  $\mu$ Ci/ml in 1 ml MEM supplemented with 10mM HEPES, pH 7.5, and 0.5% fatty-acid-free bovine serum albumin for 2 hours at 37C. The cells are then washed twice with 1 ml of the same buffer.

Candidate modulator compounds are added in 1 ml of the same buffer, either alone or with 10  $\mu$ M ATP and the cells are incubated at 37C for 30 minutes. Buffer alone and mock-transfected cells are used as controls. Samples (0.5 ml) from each well are counted by liquid scintillation spectroscopy. Agonists which activate the receptor will lead to potentiation of the ATP-stimulated release of [ $^3$ H]-arachidonic acid. This potentiation is blocked by antagonists.

#### I. Extracellular Acidification Rate

In yet another assay, the effects of candidate modulators of nGPCR-x activity are assayed by monitoring extracellular changes in pH induced by the test compounds. (See, e.g., Dunlop et al., Journal of Pharmacological and Toxicological Methods 40(1):47-55 (1998).) In one embodiment, CHO cells transfected with a nGPCR-x expression vector are seeded into 12 mm capsule cups (Molecular Devices Corp.) at  $4 \times 10^5$  cells/cup in MEM supplemented with 10% fetal bovine serum, 2 mM L-glutamine, 10 U/ml penicillin, and 10  $\mu$ g/ml streptomycin. The cells are incubated in this medium at 37C in 5% CO<sub>2</sub> for 24 hours.

Extracellular acidification rates are measured using a Cytosensor microphysiometer (Molecular Devices Corp.). The capsule cups are loaded into the sensor chambers of the microphysiometer and the chambers are perfused with running buffer (bicarbonate-free MEM supplemented with 4 mM L-glutamine, 10 units/ml penicillin, 10 µg/ml streptomycin, 26 mM NaCl) at a flow rate of 100 µl/minute. Candidate agonists or other agents are diluted into the running buffer and perfused through a second fluid path. During each 60-second pump cycle, the pump is run for 38 seconds and is off for the remaining 22 seconds. The pH of the running buffer in the sensor chamber is recorded during the cycle from 43-58 seconds, and the pump is re-started at 60 seconds to start the next cycle. The rate of acidification of the running buffer during the recording time is calculated by the Cytosoft program. Changes in the rate of acidification are calculated by subtracting the baseline value (the average of 4 rate measurements immediately before addition of a modulator candidate) from the highest rate measurement obtained after addition of a modulator candidate. The selected instrument detects 61 mV/pH unit. Modulators that act as agonists of the receptor result in an increase in the rate of extracellular acidification compared to the rate in the absence of agonist. This response is blocked by modulators which act as antagonists of the receptor.

*J. Radioligand Binding Assay*

HEK 293 or COS7 cells transiently expressing nGPCR-51, or CHO K-1 cells stable expressing nGPCR-51, were grown to sub-confluence (2 days post-transfection for transients), harvested from flasks in Dulbecco's PBS and pelleted by low speed centrifugation (2500 rpm) for 10 minutes. Cell pellets were homogenized in 10 ml tissue buffer (50 mM Hepes, 10 mM MgCl<sub>2</sub>, 2 mM EGTA, 1 µg/ml aprotinin, 1 µg/ml leupeptin hemisulfate, 1 µg/ml pepstatin A, pH 7.0) using a dounce, 10 strokes. Homogenate was centrifuged at 47,000 x g for 15 minutes. Membrane pellet was resuspended in 1 ml tissue buffer using the dounce, 10 strokes. An aliquot of the membrane preparation was used to determine protein concentration. For measurement of saturation binding, 10 µg of cell membranes were incubated with various concentrations of [<sup>125</sup>I] Peptide A (iodinated by routine procedures via the Tyr residue) in 300 µl binding assay buffer (tissue buffer plus 0.15 mM Bacitracin and 0.1% ovalbumin) for 90 minutes at room temperature in 96-well plates. Non-specific binding was defined by the inclusion of 1 µM Peptide A. After the binding incubation, plates were harvested onto GF/C filters presoaked in 0.3% non-fat dry milk. Filters were dried, and 30 µl scintillant was added to each well. Filter plates were then counted in a Packard Topcount™ 96-well microplate scintillation counter. Data were analyzed using GraphPad Prism (San Diego, CA) and K<sub>d</sub> and B<sub>max</sub> values were calculated.

In a saturation binding experiment using COS 7 cells transiently expressing nGPCR-51 the K<sub>d</sub> for [<sup>125</sup>I] Peptide A was determined to be 0.254 nM, with a B<sub>max</sub> of 75 fmol/mg. HEK 293 cells transiently expressing nGPCR-51 were shown to have an estimated B<sub>max</sub> of 108 fmol/mg. Untransfected cells have no specific radioligand binding for [<sup>125</sup>I] Peptide A.

5 **Example 13: Using nGPCR-x Proteins to Isolate Neurotransmitters**

The isolated nGPCR-x proteins, particularly nGPCR-42, 46, 48, 49, 51, 52, 61, 63, or 70 (SEQ ID Numbers 61, 62, 68, 91, 94, 96, 97, 99, 100, and 111-120) can be used to isolate novel or known neurotransmitters (Saito et al., Nature, 400: 265-269, 1999). The cDNAs that encode the isolated nGPCR-x can be cloned into mammalian expression vectors and used to stably or  
10 transiently transfect mammalian cells including CHO, Cos or HEK293 cells. Receptor expression can be determined by Northern blot analysis of transfected cells and identification of an appropriately sized mRNA band (predicted size from the cDNA). Brain regions shown by mRNA analysis to express each of the nGPCR-x proteins could be processed for peptide extraction using any of several protocols ((Reinscheid R.K. et al., Science 270: 243-247, 1996; Sakurai, T., et al., Cell 92; 573-585, 1998; Hinuma, S., et al., Nature 393: 272-276, 1998).  
15 Chromatographic fractions of brain extracts could be tested for ability to activate nGPCR-x proteins by measuring second messenger production such as changes in cAMP production in the presence or absence of forskolin, changes in inositol 3-phosphate levels, changes in intracellular calcium levels or by indirect measures of receptor activation including receptor stimulated mitogenesis, receptor mediated changes in extracellular acidification or receptor mediated  
20 changes in reporter gene activation in response to cAMP or calcium (these methods should all be referenced in other sections of the patent). Receptor activation could also be monitored by co-transfecting cells with a chimeric G<sub>i</sub>/q<sub>13</sub> to force receptor coupling to a calcium stimulating pathway (Conklin et al., Nature 363; 274-276, 1993). Neurotransmitter mediated activation of  
25 receptors could also be monitored by measuring changes in [<sup>35</sup>S]-GTPγS binding in membrane fractions prepared from transfected mammalian cells. This assay could also be performed using baculoviruses containing nGPCR-x proteins infected into SF9 insect cells.

The neurotransmitter which activates nGPCR-x proteins can be purified to homogeneity through successive rounds of purification using nGPCR-x proteins activation as a measurement  
30 of neurotransmitter activity. The composition of the neurotransmitter can be determined by mass spectrometry and Edman degradation if peptidergic. Neurotransmitters isolated in this manner will be bioactive materials which will alter neurotransmission in the central nervous system and will produce behavioral and biochemical changes.

**Example 14: Using nGPCR-x Proteins to Isolate and Purify G Proteins**

cDNAs encoding nGPCR-x proteins are epitope-tagged at the amino terminus end of the cDNA with the cleavable influenza-hemagglutinin signal sequence followed by the FLAG epitope (IBI, New Haven, CT). Additionally, these sequences are tagged at the carboxyl terminus with DNA encoding six histidine residues. (Amino and Carboxyl Terminal Modifications to Facilitate the Production and Purification of a G Protein-Coupled Receptor, B.K. Kobilka, *Analytical Biochemistry*, Vol. 231, No. 1, Oct 1995, pp. 269-271). The resulting sequences are cloned into a baculovirus expression vector such as pVL1392 (Invitrogen). The baculovirus expression vectors are used to infect SF-9 insect cells as described (Guan *et al.*, (1992) *J. Biol. Chem.* 267, 21995-21998). Infected SF-9 cells could be grown in 1000-ml cultures in SF900 II medium (Life Technologies, Inc.) containing 5% fetal calf serum (Gemini, Calabasas, CA) and 0.1 mg/ml gentamicin (Life Technologies, Inc.) for 48 hours at which time the cells could be harvested. Cell membrane preparations could be separated from soluble proteins following cell lysis. nGPCR-x protein purification is carried out as described for purification of the  $\theta$ 2 receptor (Kobilka, *Anal. Biochem.*, 231 (1): 269-271, 1995) including solubilization of the membranes in 0.8-1.0 % n-dodecyl-D-maltoside (DM) (CalBiochem, La Jolla, CA) in buffer containing protease inhibitors followed by Ni-column chromatography using chelating Sepharose™ (Pharmacia, Uppsala, Sweden). The eluate from the Ni-column is further purified on an M1 anti-FLAG antibody column (IBI). Receptor containing fractions are monitored by using receptor specific antibodies following western blot analysis or by SDS-PAGE analysis to look for an appropriate sized protein band (appropriate size would be the predicted molecular weight of the protein). This method of purifying G protein is particularly useful to isolate G proteins that bind to the nGPCR-x proteins in the absence of an activating ligand.

Some of the preferred embodiments of the invention described above are outlined below and include, but are not limited to, the following embodiments. As those skilled in the art will appreciate, numerous changes and modifications may be made to the preferred embodiments of the invention without departing from the spirit of the invention. It is intended that all such variations fall within the scope of the invention. The entire disclosure of each publication cited herein is incorporated herein by reference in its entirety.

**Example 15: Clone Deposit Information**

In accordance with the Budapest Treaty, clones of the present invention have been deposited at the Agricultural Research Culture Collection (NRRL) International Depository

Authority, 1815 N. University Street, Peoria, Illinois 61604, U.S.A. Accession numbers and deposit dates are provided below in Table 6.

**Table 6: Deposit Information**

	Clone	SEQ ID NO:	NRRL Accession Number	Budapest Treaty Deposit Date
5	nGPCR-42	31, 54	B-30246	2000 January 18
	nGPCR-48	36,56	B30263	2000 February 22
	nGPCR-46	34, 55	B-30269	2000 March 10
10	nGPCR-52	40, 58	B-30270	2000 March 10
	nGPCR-70	8, 52	B-30300	2000 June 2
	nGPCR-61	1, 60	B-30301	2000 June 2
	nGPCR-49	37, 59	B-30313	2000 July 6
	nGPCR-51	39, 57	B-30314	2000 July 6
15	nGPCR-63	3, 51, 53	B-30315	2000 July 6

**What is claimed is:**

1. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of: SEQ ID NO:61 to SEQ ID NO:120, said nucleic acid molecule encoding at  
5 least a portion of nGPCR-x.
2. The isolated nucleic acid molecule of claim 1 comprising a sequence that encodes a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:61 to SEQ ID NO:120.  
10
3. The isolated nucleic acid molecule of claim 1 comprising a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:60.
4. The isolated nucleic acid molecule of claim 1 comprising a sequence selected from the  
15 group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60.
5. The isolated nucleic acid molecule of claim 4 comprising a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60.
- 20 6. The isolated nucleic acid molecule of claim 4 wherein said nucleotide sequence is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, and SEQ ID NO:60.  
25
7. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is DNA.
8. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is  
30 RNA.
9. An expression vector comprising a nucleic acid molecule of any one of claims 1 to 5.

10. The expression vector of claim 9 wherein said nucleic acid molecule comprises a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60.

11. The expression vector of claim 9 wherein said nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, and SEQ ID NO:60.

12. The expression vector of claim 9 wherein said vector is a plasmid.

13. The expression vector of claim 9 wherein said vector is a viral particle.

14. The expression vector of claim 13 wherein said vector is selected from the group consisting of adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses.

15. The expression vector of claim 9 wherein said nucleic acid molecule is operably connected to a promoter selected from the group consisting of simian virus 40, mouse mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney virus, cytomegalovirus immediate early promoter, Epstein Barr virus, rous sarcoma virus, human actin, human myosin, human hemoglobin, human muscle creatine, and human metallothionein.

16. A host cell transformed with an expression vector of claim 9.

17. The transformed host cell of claim 16 wherein said cell is a bacterial cell.

18. The transformed host cell of claim 17 wherein said bacterial cell is *E. coli*.

19. The transformed host cell of claim 16 wherein said cell is yeast.

20. The transformed host cell of claim 19 wherein said yeast is *S. cerevisiae*.

21. The transformed host cell of claim 16 wherein said cell is an insect cell.

22. The transformed host cell of claim 21 wherein said insect cell is *S. frugiperda*.

23. The transformed host cell of claim 16 wherein said cell is a mammalian cell.

5 24. The transformed host cell of claim 23 wherein mammalian cell is selected from the group consisting of chinese hamster ovary cells, HeLa cells, African green monkey kidney cells, human 293 cells, and murine 3T3 fibroblasts.

10 25. An isolated nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60, said portion comprising at least 10 nucleotides.

15 26. The nucleic acid molecule of claim 25 wherein said molecule is an antisense oligonucleotide directed to a region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60.

20 27. The nucleic acid molecule of claim 26 wherein said oligonucleotide is directed to a regulatory region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60.

25 28. The nucleic acid molecule of claim 25 wherein said molecule is an antisense oligonucleotide directed to a region of nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, and SEQ ID NO:60.

30 29. A composition comprising a nucleic acid molecule of any one of claims 1 to 5 or 25 and an acceptable carrier or diluent.

30. A composition comprising a recombinant expression vector of claim 9 and an acceptable carrier or diluent.



31. A method of producing a polypeptide that comprises a sequence selected from the group of sequences consisting SEQ ID NO:61 to SEQ ID NO:120, and homologs thereof, said method comprising the steps of:

- a) introducing a recombinant expression vector of claim 10 into a compatible host cell;
- b) growing said host cell under conditions for expression of said polypeptide; and
- c) recovering said polypeptide.

32. The method of claim 31 wherein said host cell is lysed and said polypeptide is recovered from the lysate of said host cell.

33. The method of claim 31 wherein said polypeptide is recovered by purifying the culture medium without lysing said host cell.

34. An isolated polypeptide encoded by a nucleic acid molecule of claim 1.

35. The polypeptide of claim 34 wherein said polypeptide comprises a sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

36. The polypeptide of claim 34 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

37. The polypeptide of claim 34 wherein said sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120 comprises at least one conservative amino acid substitution compared to the sequences in the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

38. The polypeptide of claim 34 wherein said polypeptide comprises an allelic variant of a polypeptide with a sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

39. The polypeptide of claim 34 wherein said polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100,

SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

40. A composition comprising a polypeptide of claim 34 and an acceptable carrier or diluent.

41. An isolated antibody which binds to an epitope on a polypeptide of claim 34.

42. The antibody of claim 41 wherein said antibody is a monoclonal antibody.

43. A composition comprising an antibody of claim 41 and an acceptable carrier or diluent.

44. A method of inducing an immune response in a mammal against a polypeptide of claim 34 comprising administering to said mammal an amount of said polypeptide sufficient to induce said immune response.

45. A method for identifying a compound which binds nGPCR-x comprising the steps of:  
a) contacting nGPCR-x with a compound; and  
b) determining whether said compound binds nGPCR-x.

46. The method of claim 45 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

47. The method of claim 45 wherein binding of said compound to nGPCR-x is determined by a protein binding assay.

48. The method of claim 45 wherein said protein binding assay is selected from the group consisting of a gel-shift assay, Western blot, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, and ELISA.

49. A compound identified by the method of claim 45.

50. A method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x comprising the steps of:

- a) contacting said nucleic acid molecule encoding nGPCR-x with a compound; and
- b) determining whether said compound binds said nucleic acid molecule.

5

51. The method of claim 50 wherein binding is determined by a gel-shift assay.

52. A compound identified by the method of claim 50.

10 53. A method for identifying a compound which modulates the activity of nGPCR-x comprising the steps of:

- a) contacting nGPCR-x with a compound; and
- b) determining whether nGPCR-x activity has been modulated.

15 54. The method of claim 53 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

20

55. The method of claim 53 wherein said activity is neuropeptide binding.

56. The method of claim 53 wherein said activity is neuropeptide signaling.

25 57. A compound identified by the method of claim 53.

58. A method of identifying an animal homolog of nGPCR-x comprising the steps:

- a) comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequence consisting of SEQ ID NO:1 to SEQ ID NO:60, and portions thereof, said portions being at least 10 nucleotides; and
- b) identifying nucleic acid sequences of the animal that are homologous to said sequence selected from the group sequence consisting of SEQ ID NO:1 to SEQ ID NO:60, and portions thereof.

30

59. The method of claim 58 wherein comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60, and portions thereof, said portions being at least 10 nucleotides, is performed by DNA hybridization.

60. The method of claim 58 wherein comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:60, and portions thereof, said portions being at least 10 nucleotides, is performed by computer homology search.

61. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR that is expressed in the brain, wherein the nGPCR comprises an amino acid sequence selected from the group consisting of SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120, and allelic variants thereof, and wherein the nucleic acid corresponds to a gene encoding the nGPCR; and

b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR in the nucleic acid correlates with an increased risk of developing the disorder.

62. The method of claim 61, wherein the nGPCR is nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70, or an allelic variant thereof.

63. The method of claim 61, wherein the nGPCR is nGPCR-51 or nGPCR-52 or an allelic variant thereof.

64. The method of claim 61, wherein the disease is a mental disorder.

65. The method of claim 61, wherein the assaying step comprises at least one procedure selected from the group consisting of:

a) comparing nucleotide sequences from the human subject and reference sequences and determining a difference of either

5           at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-42 allele and an nGPCR-42 reference sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-46 allele and an nGPCR-46 reference  
10          sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-48 allele and an nGPCR-48 reference sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-49 allele and an nGPCR-49 reference  
15          sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-51 allele and an nGPCR-51 reference sequence;

20          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-52 allele and an nGPCR-52 reference sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-61 allele and an nGPCR-61 reference  
25          sequence;

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-63 allele and an nGPCR-63 reference sequence; or

          at least a nucleotide of at least one codon between the nucleotide sequences from the human subject that encodes an nGPCR-70 allele and an nGPCR-70 reference  
30          sequence;

b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences;

c) performing a polynucleotide migration assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and

5 d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

66. The method of claim 65 wherein the assaying step comprises: performing a polymerase chain reaction assay to amplify nucleic acid comprising nGPCR-42, nGPCR-46, nGPCR-48,  
10 nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 coding sequence, and determining nucleotide sequence of the amplified nucleic acid.

67. A method of screening for an nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 hereditary mental disorder genotype in a  
15 human patient, comprising the steps of:

a) providing a biological sample comprising nucleic acid from said patient, said nucleic acid including sequences corresponding to alleles of nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70; and

b) detecting the presence of one or more mutations in the nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 allele;  
20 wherein the presence of a mutation in a nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 allele is indicative of a hereditary mental disorder genotype.

25 68. The method of claim 67 wherein said biological sample is a cell sample.

69. The method of claim 67 wherein said detecting the presence of a mutation comprises sequencing at least a portion of said nucleic acid, said portion comprising at least one codon of said nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61,  
30 nGPCR-63, or nGPCR-70 allele.

70. The method of claim 67 wherein said nucleic acid is DNA.

71. The method of claim 67 wherein said nucleic acid is RNA.

72. A kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association:

a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 gene, the oligonucleotide comprising 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 gene sequence or nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

b) a media packaged with the oligonucleotide, said media containing information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

73. A method of identifying a nGPCR allelic variant that correlates with a mental disorder, comprising the steps of:

a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

b) detecting in the nucleic acid the presence of one or more mutations in an nGPCR that is expressed in the brain, wherein the nGPCR comprises an amino acid sequence selected from the group consisting of SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120, and allelic variants thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding nGPCR;

wherein the one or more mutations detected indicates an allelic variant that correlates with a mental disorder.

74. The method of claim 73 wherein the nGPCR is nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70, or an allelic variant thereof.

75. A purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 allelic variant identified according to claim 74.

5 76. A host cell transformed or transfected with a polynucleotide according to claim 75 or with a vector comprising the polynucleotide.

77. A purified polynucleotide comprising a nucleotide sequence encoding nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or  
10 nGPCR-70 of a human with a mental disorder;  
wherein said polynucleotide hybridizes to the complement of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, and SEQ ID  
15 NO:60 under the following hybridization conditions:

a) hybridization for 16 hours at 42C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate; and

b) washing 2 times for 30 minutes at 60C in a wash solution comprising 0.1x SSC and 1% SDS;

20 wherein the polynucleotide that encodes nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 amino acid sequence of the human differs from SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57,  
25 SEQ ID NO:58, SEQ ID NO:59, and SEQ ID NO:60 by at least one residue.

78. A vector comprising a polynucleotide according to claim 77.

79. A host cell that has been transformed or transfected with a polynucleotide according to  
30 claim 77 and that expresses the nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 protein encoded by the polynucleotide.

80. The host cell of claim 79 that has been co-transfected with a polynucleotide encoding and expressing the nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52,



nGPCR-61, nGPCR-63, or nGPCR-70 amino acid sequence set forth in SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

81. A method for identifying a modulator of biological activity of nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 comprising the steps of:

10 a) contacting a cell according to claim 79 in the presence and in the absence of a putative modulator compound; and

b) measuring nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 biological activity in the cell;

15 wherein decreased or increased nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

82. A method to identify compounds useful for the treatment of a mental disorder, said method comprising the steps of:

20 a) contacting a composition comprising nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 with a compound suspected of binding nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70; and

25 b) detecting binding between nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 and the compound suspected of binding nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70;

30 wherein compounds identified as binding nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 are candidate compounds useful for the treatment of a mental disorder.

83. A method for identifying a compound useful as a modulator of binding between nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or

nGPCR-70 and a binding partner of nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 comprising the steps of:

a) contacting the binding partner and a composition comprising nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 in the presence and in the absence of a putative modulator compound; and

b) detecting binding between the binding partner and nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70;

wherein decreased or increased binding between the binding partner and nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

84. The method of claim 82 or claim 83 wherein the composition comprises a cell expressing nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70 on its surface.

85. The method of claim 84 wherein the composition comprises a cell transformed or transfected with a polynucleotide that encodes nGPCR-42, nGPCR-46, nGPCR-48, nGPCR-49, nGPCR-51, nGPCR-52, nGPCR-61, nGPCR-63, or nGPCR-70.

86. A method of purifying a G protein from a sample containing said G protein comprising the steps of:

a) contacting said sample with a polypeptide of claim 1 for a time sufficient to allow said G protein to form a complex with said polypeptide;

b) isolating said complex from remaining components of said sample;

c) maintaining said complex under conditions which result in dissociation of said G protein from said polypeptide; and

d) isolating said G protein from said polypeptide.

87. The method of claim 86 wherein said sample comprises an amino acid sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

88. The method of claim 86 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:61 to SEQ ID NO:120.

5 89. The method of claim 86 wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:68, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

10

90. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to SEQ ID NO:117.

15 91. The nucleic acid molecule of claim 90 wherein said polypeptide comprises SEQ ID NO:117.

92. An isolated nucleic acid molecule comprising SEQ ID NO:57.

20 93. An isolated polypeptide comprising an amino acid sequence homologous to SEQ ID NO:117.

94. The polypeptide of claim 93 comprising SEQ ID NO:117.

25 95. A method of identifying a compound that binds to nGPCR-51 comprising the steps of:  
a) contacting a composition comprising nGPCR-51, or a polypeptide homologous thereto, and a polypeptide comprising SEQ ID NO:183, or a polypeptide homologous thereto, with said test compound; and  
b) determining whether said test compound binds to nGPCR-51.

30 96. The method of claim 95 wherein said determining whether said test compound binds to nGPCR-51 is by measuring the displacement of said polypeptide comprising SEQ ID NO:183 from a complex between said polypeptide and nGPCR-51.

97. The method of claim 95 wherein said polypeptide comprising SEQ ID NO:183 is radiolabeled.

98. The method of claim 97 wherein step b) is determined by comparatively measuring radioactivity of nGPCR-51 bound to said radiolabeled polypeptide comprising SEQ ID NO:183 with the radioactivity of nGPCR-51 in the presence of the test compound.

99. The method of claim 95 wherein step b) comprises a binding assay selected from the group consisting of filter binding, scintillation proximity assay, gel-shift assay, radiolabeled competition assay, and ELISA.

100. A method for identifying a compound that modulates the activity of nGPCR-51 comprising the steps of :

a) contacting nGPCR-51 with a compound; and

b) determining whether nGPCR-51 activity has been modulated.

101. The method of claim 100 wherein said activity is neuropeptide binding.

102. The method of claim 101 wherein said neuropeptide binding is determined by binding to a polypeptide comprising SEQ ID NO:183.

103. The method of claim 101 wherein said neuropeptide binding is determined by binding to a polypeptide comprising SEQ ID NO:184.

104. The method of claim 100 wherein said activity is neuropeptide signaling.

105. The method of claim 104 wherein said neuropeptide signalling is determined by binding to a polypeptide comprising SEQ ID NO:183.

106. The method of claim 104 wherein said neuropeptide signalling is determined by binding to a polypeptide comprising SEQ ID NO:184.

107. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of nGPCR-51, or an allelic variant thereof; and

5 b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR in the nucleic acid correlates with an increased risk of developing the disorder.

108. The method of claim 107 wherein the disorder is schizophrenia.

109. The method of claim 107 wherein the disorder is an attention disorder.

110. The method of claim 107 wherein the disorder is anxiety.

15 111. The method of claim 107 wherein the disorder is depression.

112. The method of claim 107 wherein the disorder is obesity.

20 113. A kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association:

a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-51, the oligonucleotide comprising 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-51 gene sequence or nGPCR-51 coding sequence, except for one sequence difference selected from the group consisting of a  
25 nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

b) a media packaged with the oligonucleotide, said media containing information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

30 114. A method of identifying a nGPCR allelic variant that correlates with a mental disorder, comprising the steps of:

a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

b) detecting in the nucleic acid the presence of one or more mutations in an nGPCR that is expressed in the brain, wherein the nGPCR comprises SEQ ID NO:117 or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding nGPCR;

5            wherein the one or more mutations detected indicates an allelic variant that correlates with a mental disorder.

115.    A method for identifying a modulator of biological activity of nGPCR-51 comprising the steps of:

10            a) contacting a cell according to claim 79 in the presence and in the absence of a putative modulator compound; and

            b) measuring nGPCR-51 biological activity in the cell;

            wherein decreased or increased nGPCR-51 biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

15            116.    A method of identifying a compound useful for the treatment of a mental disorder, said method comprising the steps of:

            a) contacting a composition comprising nGPCR-51 with a compound suspected of binding nGPCR-51; and

20            b) detecting binding between nGPCR-51 and the compound suspected of binding nGPCR-51;

            wherein a compound identified as binding nGPCR-51 is a candidate compound useful for the treatment of a mental disorder.

25            117.    A method for identifying a compound useful as a modulator of binding between nGPCR-51 and a binding partner of nGPCR-51 comprising the steps of:

            a) contacting the binding partner and a composition comprising nGPCR-51 in the presence and in the absence of a putative modulator compound; and

            b) detecting binding between the binding partner and nGPCR-51;

30            wherein decreased or increased binding between the binding partner and nGPCR-51 in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

118. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to SEQ ID NO:118.

119. The nucleic acid molecule of claim 118 wherein said polypeptide comprises SEQ ID NO:118.

120. An isolated nucleic acid molecule comprising SEQ ID NO:58.

121. An isolated polypeptide comprising an amino acid sequence homologous to SEQ ID NO:118.

122. The polypeptide of claim 93 comprising SEQ ID NO:118.

123. A method of identifying a compound that binds to nGPCR-52 comprising the steps of:

a) contacting a composition comprising nGPCR-52, or a polypeptide homologous thereto, and a polypeptide comprising SEQ ID NO:183, or a polypeptide homologous thereto, with said test compound; and

b) determining whether said test compound binds to nGPCR-52.

124. The method of claim 123 wherein said determining whether said test compound binds to nGPCR-52 is by measuring the displacement of said polypeptide comprising SEQ ID NO:183 from a complex between said polypeptide and nGPCR-52.

125. The method of claim 123 wherein said polypeptide comprising SEQ ID NO:183 is radiolabeled.

126. The method of claim 125 wherein step b) is determined by comparatively measuring radioactivity of nGPCR-52 bound to said radiolabeled polypeptide comprising SEQ ID NO:183 with the radioactivity of nGPCR-52 in the presence of the test compound.

127. The method of claim 123 wherein step b) comprises a binding assay selected from the group consisting of filter binding, scintillation proximity assay, gel-shift assay, radiolabeled competition assay, and ELISA.

128. A method for identifying a compound that modulates the activity of nGPCR-52 comprising the steps of :

- a) contacting nGPCR-52 with a compound; and
- b) determining whether nGPCR-52 activity has been modulated.

5

129. The method of claim 128 wherein said activity is neuropeptide binding.

130. The method of claim 129 wherein said neuropeptide binding is determined by binding to a polypeptide comprising SEQ ID NO:183.

10

131. The method of claim 129 wherein said neuropeptide binding is determined by binding to a polypeptide comprising SEQ ID NO:184.

132. The method of claim 128 wherein said activity is neuropeptide signaling.

15

133. The method of claim 132 wherein said neuropeptide signalling is determined by binding to a polypeptide comprising SEQ ID NO:183.

20

134. The method of claim 132 wherein said neuropeptide signalling is determined by binding to a polypeptide comprising SEQ ID NO:184.

135. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

25 a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of nGPCR-52, or an allelic variant thereof; and

b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR in the nucleic acid correlates with an increased risk of  
30 developing the disorder.

136. The method of claim 135 wherein the disorder is schizophrenia.



137. A kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association:

a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-52, the oligonucleotide comprising 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-52 gene sequence or nGPCR-52 coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

b) a media packaged with the oligonucleotide, said media containing information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

138. A method of identifying a nGPCR allelic variant that correlates with a mental disorder, comprising the steps of:

a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

b) detecting in the nucleic acid the presence of one or more mutations in an nGPCR that is expressed in the brain, wherein the nGPCR comprises SEQ ID NO:118 or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding nGPCR;

wherein the one or more mutations detected indicates an allelic variant that correlates with a mental disorder.

139. A method for identifying a modulator of biological activity of nGPCR-52 comprising the steps of:

a) contacting a cell according to claim 79 in the presence and in the absence of a putative modulator compound; and

b) measuring nGPCR-52 biological activity in the cell;

wherein decreased or increased nGPCR-52 biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

140. A method of identifying a compound useful for the treatment of a mental disorder, said method comprising the steps of:

a) contacting a composition comprising nGPCR-52 with a compound suspected of binding nGPCR-52; and

b) detecting binding between nGPCR-52 and the compound suspected of binding nGPCR-52;

wherein a compound identified as binding nGPCR-52 is a candidate compound useful for the treatment of a mental disorder.

5

141. A method for identifying a compound useful as a modulator of binding between nGPCR-52 and a binding partner of nGPCR-52 comprising the steps of:

a) contacting the binding partner and a composition comprising nGPCR-52 in the presence and in the absence of a putative modulator compound; and

10 b) detecting binding between the binding partner and nGPCR-52;

wherein decreased or increased binding between the binding partner and nGPCR-52 in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

15

## SEQUENCE LISTING

<110> Pharmacia & Upjohn Company  
Vogeli, Gabriel  
Parodi, Luis A.  
Hiebsch, Ronald R.  
Lind, Peter  
Kaytes, Paul S.  
Ruff, Valerie  
Huff, Rita M.  
Wood, Linda S.

<120> Novel G Protein-Coupled Receptors Cross-Reference To Related Applications

<130> 00325.PCT1

<150> 60/184,305

<151> 2000-02-23

<150> 60/184,304

<151> 2000-02-23

<150> 60/184,303

<151> 2000-02-23

<150> 60/184,397

<151> 2000-02-23

<150> 60/184,247

<151> 2000-02-23

<150> 60/188,880

<151> 2000-03-13

<150> 60/217,369

<151> 2000-07-11

<150> 60/217,370

<151> 2000-07-11

<150> 60/218,492

<151> 2000-07-20

<150> 60/186,810

<151> 2000-03-03

<150> 60/188,064

<151> 2000-03-09

<150> 60/186,457

<151> 2000-03-02

<150> 60/213,861

<151> 2000-06-23

<150> 60/194,344

<151> 2000-04-03

<150> 60/218,337

&lt;151&gt; 2000-07-14

&lt;160&gt; 184

&lt;170&gt; PatentIn version 3.0

&lt;210&gt; 1

&lt;211&gt; 545

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

```

cctatgctac aatcggcaga tgggaactag gagccatgat ctctcagatt gcaggtctca      60
ttggaaccac atttattgga ttttcctttt tagtagtact aacatcatac tactcttttg      120
taagccatct gagaaaaata agaacctgta cgtccattat ggagaaagat ttgacttaca      180
gttctgtgaa aagacatctt ttggatcatc agattctact aatagtttgc ttccttcctt      240
atagtatttt taaaccattt ttttatgttc tacaccaaag agataactgt cagcaattga      300
attatttaat agaaacaaaa aacattctca cctgtcttgc ttcggccaga agtagcacag      360
acccatttat atttctttta ttagacaaaa cattcaagaa gacactatat aatctcttta      420
caaagtctaa ttcagcacat atgcaatcat atggttgact tttgaatgga aaaccccaca      480
atattaagaa aagcattcat gtgactttat tagggacact aaactacatc attaacatgt      540
cacag                                             545

```

&lt;210&gt; 2

&lt;211&gt; 1611

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

cagtgagccg agatggtgcc attgcactct agcctggggc aacagagccg actccatctc      60
caaaaaaaaa aggccattct gaggatcaag gcaccactag caacagggag ccccatgggt      120
ctcagaccct ctccccacat ctcttggtcc ctgccccac ctggcgtaca gggaccagcc      180
ccacggaagg ctcttgaggc caggtaacca tggggagggg aggaatgggg acaccttctt      240
cctgagtgtc ttagggaaga gaagcttagg tcaggtggct gaggggtggaa atgagagagg      300
ggtctcctcc tggagggtct caccattccc ttggtcacc acccaactct catctccctt      360
gatgtgggga ggagcagggg gcatggattc ctgagcccca gactcaactg ttgtgggtta      420
caggggcatc aggagagaga gcgagcagaa cacactcctg cagcatcccc tggccccccg      480
cccatgatg gagccagag aagctggaca gcacgtgggg gccgccaacg gcgcccagga      540
ggatgtggcc ttcaacctca tcctcctgtc cctcaccgag gggctcggcc tcgggtgggt      600

```

gctggggaat ggggcagtc tctggctgct cagctccaat gtctacagaa accccttcgc 660  
 catctacctc ctggacgtgg cctgcgcgga tctcatcttc cttggctgcc acatgggtggc 720  
 catcgteccc gacttgctgc aaggccggct ggacttcccg ggcttcgtgc agaccagcct 780  
 ggcaacgctg cgcttcttct gctacatcgt gggcctgagt ctcttgccgg ccgtcagcgt 840  
 ggagcagtgc ctggccgccc tcttcccagc ctggtactcg tgccgccgcc cagccacct 900  
 gaccacctgt gtgtgcgccc tcacctgggc cctctgcctg ctgctgcacc tgctgctcag 960  
 cggcgccctgc acccagttct tcggggagcc cagccgccac ttgtgccgga cgctgtggct 1020  
 ggtggcagcg gtgctgctgg ctctgctgtg ttgcaccatg tgtggggcca gccttatgct 1080  
 gctgctgcgg gtggagcgag gccccagcg gccccaccc cggggcttcc ctgggctcat 1140  
 cctcctcacc gtctcctct tctcttctg cggcctgcc ttcggcatct actggctgtc 1200  
 ccggaacctg ctctggtaca tccccacta cttctaccac ttcagcttcc tcatggccgc 1260  
 cgtgcaactgc ggggccaagc ccgtcgtcta cttctgcctg ggcaagtccc agggccgcag 1320  
 gctgcccctc cggctgggtcc tccagcgagc gctgggagac gaggctgagc tgggggcccgt 1380  
 cagggagacc tcccgcggg gcctggtgga catagcagcc tgagccctgg ggcccccgac 1440  
 cccagctgca gcccccgta ggcaagagg tgacgtgggg aaggtggtgg ggtcagaggc 1500  
 tggggccagc cggacctgga ggaggccttg gtgggtgacc cggatcatgt ctgtcaaagt 1560  
 tgtgaccctt ggtctggagc atgaggctcc cctgggaggc agctggaaag g 1611

<210> 3  
 <211> 930  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 aaaattgctc ttctcctga gcttgtagac aatgattgag ttcaagatga agaagatgga 60  
 gcagggcacc aggtagacgg tgaagcagtg gatccagatg aggacgtgat gcacagagg 120  
 gctgatgtag tcttcagtcc agatgttggg ccaccagtaa taggggatgc tggtcaggaa 180  
 gcaggtgatg taaacactta caatgacttt ccgggtgcgg gctgggtatg agaccgtgtg 240  
 gtacttgagc ggggtggcaga cagcgatata cctgtcaatg gttaacggta cagtaatcca 300  
 tatggagggtg tggatggatg agaattccag cacttctatg atcttgtcgg ggacctgagg 360  
 catctgcatg ttcaagatga aatcttccaa caggaagtcc acaaacta tgaaaaagag 420  
 gaccaagatg tcggcagcag cgagtgccaa gagatagttg taggaggact tctgtcttct 480  
 tgccaccagc tgggagagga tgatcactgt caagatattt gctgtggaga gaagaaaaac 540

tgggttagct ctgaagcaaa gatgacttcg ttggctccta tgggggccct aggcatatgt 600  
 ttatitttgca ctcccatgga agtgaaaatg attgaatcaa tgcttttgag ggacaacctc 660  
 agcattacaa atagcacctc atacaattag tggatactat tttaaagtta tgcttatatt 720  
 ctaacacaac catgagaggt ggtgcctcca ttctcctcat cttagaagtg aaactggggc 780  
 tctgagagcc tcacacagcc atgagaggtg gtgccgcat tctcctcaca caaccatgag 840  
 tgggtgttgc gccattctcc tcacacaaca atgagaggtg gtgccaccat tcttctcaca 900  
 caaccgtgag aggcgatgct gccattatcc 930

<210> 4  
 <211> 533  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 ggcccgtcc cacgctgtgt agtgtacttt catitttcaat aaatcacttc attccttcct 60  
 tgctttgttt gtgcgttttg tccaattctt tgttcaagac gccaagaacc tggacagcct 120  
 ccaccattaa gaatacaaga gcagtttctg tcacatgtac atatgggggt ggggtggatct 180  
 cgctcagcct ttccaggaca cagcgttttg atggaatgct tttctgaaac tcgtggcaga 240  
 atgagtacgg aagagggctg atagccgaat tcagccagtg gagccagaag cgtgtctcat 300  
 agagaaagtg ctggacatgt gaccctggc aggcagctcc gatgatcacg agcagtgtag 360  
 ggagcccagc acaggacaaa tgcacacacc atgacgtcaa gcgacgtgga cgctcatttg 420  
 tcccgggaga cgccgtgctg gggaggctcg gctgtctgct gccattgcag gcgttctcgc 480  
 tgccctgtag cgggtcttct gcgtaggatt atcaagtcca cctgcagagg tga 533

<210> 5  
 <211> 589  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 tacttttacc caaaactatt atgttcatta ttagagtttc ctagaaaaat acctaaggag 60  
 ttaatggttt catatctttt gctctattat gaagaagaaa aatgttttac aaatattttc 120  
 attattggag cattttttgt tgttagtgaa attatcaaaa ctaggattga tttctattct 180  
 gtttactttt gttataatct ttatcctttt ctcttaattt ctgtattttg gatgcctaac 240  
 ttagaataca ttaccaaagt taccttttca ttagtctct caatacaaga tgatttaaaa 300  
 catttatggg tacctttttt aatttttttg ctatgcaaat ttataaaagg gcaaagtctt 360  
 tgtgctctaa taatacctgc tttctcatgt ttacatgtt ctacgattta ttttgttttt 420

ataatgtaat tttcgttttac ctaattgtgc acatagtga taatagatta taatgaagaa 480  
aacttggatt aaaatctatt gttaaaaagg tttttcaggc aataataaat cattggattt 540  
ttctgatgta ttttaaaaag atatgtttat ttttgagcaa ctctgtgtgc 589

<210> 6  
<211> 864  
<212> DNA  
<213> Homo sapiens

<400> 6  
ggctacttgt tatggaaaag ttatgattgg tgtagaactg tacaggttgc ttctggcact 60  
tgaagaaagc cttgagcgtg gcggtgccca tggcgcgga cgccttgctg atgaagcagt 120  
agaggaagaa gttgatggct gtgttcagaa gggctagcat gttggcaatg tcggacatga 180  
tgtgtaccag ccagcgggtc tggatgggcg ccccatagag gtggtaaaga atcatgatga 240  
tgcggggggc ccaaagtgtg gcaaagatgg aggtaatggt gaacaagatg gcggtggtct 300  
tccccgtgga gtagccacgg agacgaaaat tgctcttctt cctgagcttg tacacaatga 360  
ttgagttcaa gatgaagaac atggagcagg gcaccaagta gacggtgaag cagtggatcc 420  
agatgaggac gtgatgcaca gaggtgctga tgtaatcttc agtccagatg ttgggccacc 480  
agtaataggg gatgctggtc aggaagcagg tgatgtaaac acttacaatg actttccggg 540  
tgcgggctgg gtatgagacc gtgtggtact tgagcgggtg gcagacagcg atatacctgt 600  
caatgggttaa cggtagagta atccatatgg aggtgtggat ggatgagaat tccagcactt 660  
ctatgatctt gtcggggacc tgaggcatct gcatgttcaa gatgaaatct tccaacagga 720  
agtccacaaa cactatgaaa aagaggacca agatgtcggc agcagcgagt gccaaagat 780  
agttgtagga ggacttctgt cttcttgcca ccagctggga gaggatgatc actgtcaaga 840  
tatttgctgt ggagagaaga aaac 864

<210> 7  
<211> 823  
<212> DNA  
<213> Homo sapiens

<400> 7  
gacacctgcc aacatgttca ttatcaacct cgcggtcagc gacttcctca tgccttcac 60  
ccaggcccct gtcttcttca ccagtagcct ctataagcag tggctctttg gggagacagg 120  
tagatgctgg ggctcccttt tgctggaggg aggaggaggg ttttgacctg gggatgccct 180  
caatggaggg tggcccaaag gaggtgattt gctgcttctg ggcagagagt gggtagctgc 240

```

cctcagtcct gtgagtaagc aagaagggaa gatgcagtgt tggtcctaag gcctctgccca 300
gccttggcca gatgtggcag gtggaggggg tggagtgcgc tcagtcctgc tcttcctgtg 360
aggtgaaggc cagagcagag tctaccctgt cccagaccc tcctccccag gactcagagc 420
aggggctgtg ccacacaggct gcgagttcta tgccttctgt ggagctctct ttggcatttc 480
ctccatgata accctgacgg ccacgcacct ggaccgctac ctggtaatca cagccccgct 540
ggccaccttt ggtgtggcgt ccaagaggcg tgcggcattt gtcctgctgg gcgtttggct 600
ctatgccctg gcctggagtc tgccaccctt cttcggctgg agtaagtggg ctgctggaac 660
tggaaggggg gcagatgggc tgggaggggc acattcaagg gtaagtaggt gacttgggtc 720
agccagctgg cgggagcagg gtgcccagga gctacctgag cctcaggtga gatggacatt 780
cagggggaca tgactggcag caagggaaac tgacaactgcc cca 823

```

```

<210> 8
<211> 619
<212> DNA
<213> Homo sapiens

```

```

<400> 8
ttctcccctt gacgggtgac taactctgcc tgcgtgtttc ttttgtcacc agcataggca 60
ctgagtgcgg tctgtgcacc cctttgccac ccaccggtgc cggcactgag cctgcaacct 120
gtctcacgcc ctctggctgt tgccatgacg tccacctgca ccaacagcac gcgcgagagt 180
aacagcagcc acacgtgcat gcccctctcc aaaatgccc tccagcctggc ccacggcatc 240
atccgctcaa ccgtgctggt tatcttctc gccgcctctt tcgtcggcaa catagtgtg 300
gcgctagtgt tgcagcgcaa gccgcagctg ctgcaggtga ccaaccgttt tatctttaac 360
ctcctcgtca ccgacctgct gcagatttcg ctcggtggcc cctgggtggt ggccacctct 420
gtgcctctct tctggcccct caacagccac ttctgcacgg ccctgggttag cctcaccac 480
ctgttcgcct tcgccagcgt caacaccatt gtcgtggtgt cagtggatcg ctacttgtcc 540
atcatccacc ctctcttcta ccggtccaag atgaccacgc gccgcgggta cctgtctctc 600
tatggcacct ggattgtgg 619

```

```

<210> 9
<211> 550
<212> DNA
<213> Homo sapiens

```

```

<400> 9
aggacaacga tggtcactga tttggtgacc ttcgacagtc tccgggcgct ggctccggtc 60
gggcgtctc cggctaccgc ggcccctcct ttggtcccc cgcgcggcg gtcggcgatg 120

```



aagcgcacca gcagcaggta gcacaagata atgatgccca gcggcagcac gaagcccagc	180
agcaccttct gcgagtggta gaggcccagc cagaactgcc tgtcgcggcc cagcaacttg	240
tccgggaaac gcaccaggca cagctcctcg cccatcacct tgaccgtggt ggagaaaatg	300
gcactgggca gcgaggccag cgcgggccaaa gccagatcc acacacacag cgccttgcc	360
gagaagcagc agctgtcccc caggctccgg ccgcagcagt cgcgccggcc gtgtcctcgg	420
gtcccgttg ctcttcagag ccgaggccac cgaatggtag cgcgtcacac tcatggcagt	480
gaggaagaac acgctggcgt acatgttcat gggacgtcac catggacacg atcttacaca	540
tggccttgcc	550

<210> 10  
 <211> 1104  
 <212> DNA  
 <213> Homo sapiens

<400> 10	
ctggaaaggg tgctgggtggc gcgcgcgcac gccgcggcgc gactgcctgc ctggtactca	60
acctcttctg cgcggacctg ctcttcatca gcgtatccc tctggtgctg gccgtgcgct	120
ggactgaggc ctggctgctg ggccccgttg cctgccacct gctctttctac gtgatgacct	180
tgagcggcag cgtcaccatc ctcacgctgg ccgcggtcag cctggagcgc atggtgtgca	240
tcgtgcacct gcagcgcggc gtgcggggtc ctgggcggcg ggcgcgggca gtgctgctgg	300
cgctcatctg gggctattcg gcggtcgcc ctctgcctct ctgcgtcttc ttccgagtcg	360
tcccgcaacg gctccccggc gccgaccagg tgagcgcgcc tctgtgtgtg ccgggcaggc	420
gtcctgcgca ggctgggaag cggggccccc acggaagctg ggatgaggat gatcaagaac	480
aacaatagcc atttattgca cttaatcggt gtgccaaatc ttgtgcccac ggctgtgaag	540
tttaatctct taaatctcac tacaacgctg tgcacacgcc ctctaaatg atgtaagtgg	600
agtcccccaa attcttgcaa aatgcaatga ctgttgcgag gttaattaac gagtagttta	660
ggagcgagac ggaacttttg ggggtgcaggg tggccaaaca ctttgtattg aatcatgatt	720
cctcgccagg tgctacaata ctgttattat cacaccatt tcacagatga gaaccagagg	780
cacaccgaag tgtataataa cttgccaga gtattttatc cgtaattcga ggaggagatg	840
ggctccttcc agaagtttac ccgtaattca aggaggagt gggtcctgt ccagggttg	900
gttatggtcc tgctttgaaa gcgcgcggac aggcattgta gaccgggga cccagatgc	960
aatgctgtct ttaggggact tgtgacagaa ttcccttcg gggcttcag ttttttcagc	1020
tgcaaaacgg aaggattaca ctagacctc gaggtgtcct gggcgctga aatgtgcaga	1080

ttacagaggc tggaccgacg agct

1104

<210> 11  
 <211> 591  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 ttaatccctg gaagtccacg aacaatgaat ccatttcatg catcttggtg gaacacctct 60  
 gccgaacttt taaacaaatc ctggaataaa gagtttgctt atcaaactgc cagtgtggta 120  
 gatacagtca tcctcccttc catgattggg attatctggt caacagggct ggttggcaac 180  
 atcctcattg tattcactat aataaggtaa ggaatggctc cttttttttt tttttccttc 240  
 catacttttag gaaactacag tcaaagctcc ctaaagtagt cctttccctt gtagcatttt 300  
 gcttaatgaa atgcaatttt ggaaatattt gcttaagata attaataag attctacaga 360  
 tatttttcgtc atgcattagg taacatctca gttgcaaact tcaacatgct aagacctagg 420  
 ccaatgctta ctgctgggtc agtgagtttt tagggaaatg actctcactc tcagtcttag 480  
 ctgcatatta gaatcatctg gggagcttta aaaactcctg atatgcagtt tcaccccaga 540  
 cccattaact cagaatctct aagggtaggg cccgggtaag atttaaaact g 591

<210> 12  
 <211> 378  
 <212> DNA  
 <213> Homo sapiens

<400> 12  
 gggtgacagg aactctggaa ggtgatagtt ttccagggtga agaagggaca gtaggtttcc 60  
 ctgctttact gctatctccc tatggccttc ggtcaggcct gaactgtgat gagagtctct 120  
 gccctgctc tttgggattc ctcaaataac ctcccttcat tgttccctga attatctaga 180  
 tcaatacaaaa ccaatccctc ctctgcttaa aacttttaat tgcttacagg ggaaagtgoa 240  
 gactccttag ctggcgacac agcctttttgc acctggcaca catagagctt tctagggcta 300  
 tctcctgcca ttctctagcc tgtctccttc tgctctgggt ctccagaact gcccatgaat 360  
 tccctcacat gcctttct 378

<210> 13  
 <211> 503  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 cctatgcttt cctatagctc atggaccaag ccatttctcc aacatcataa tttttcctta 60

```

tcatcccttt ccttgcaatta ttctccagat gaaaaaagtt caaatgcctc ctccctattt 120
tcagctgaga aggtgagatg cccgtgcctg gatttccagg ctctgcctg cttggcagac 180
cactgtgacc gtgcacgact aagtcactc ttcccagcca gcatcccctg ctctccccag 240
gccaacttct ttgtgttcta ttcatctcta tctccttgcc ggaactactc agaccctct 300
gtctttctgc ctcttgccca atcttccagg ctctagtttg ctggctttct gcctccaaaa 360
atgattttta gcatttgtga gtctttctct ccacagaact ccagactctt tgaaaatgcc 420
gtagtctgta aattacttcc aatatttaag aagtgcctat catgcttccc tgcgggttca 480
tgctctttgg tattgatattt cct 503

```

```

<210> 14
<211> 671
<212> DNA
<213> Homo sapiens

```

```

<400> 14
caaatgaggt ggatgcactt agaaggaagg gtagaacaaa tattcggtga tgatatacaa 60
tgtgtcagat acagagatac acagactagg gctgtgggaa tctcagattt taatttgtca 120
acagtgtgtt ttgatttttt tgttttgtat atttgctcc ccaattaatt tcacaggctt 180
aatcatcttt acaagacatt attttaaga gaaagcgaac ttactgaagt tttatgcttc 240
cctgattgtg atgagctggt tgattctagc tctagtttca atgttctgga aaatactgaa 300
gtacttccac ctggtggcac ttagtgaaca ttgcagaacc gagtaatagg ttattccgtt 360
gggtttctcg aacaaatctg agttatagct aggaactctc aactaacaat ttatgagaac 420
cttctgctac acatgtgaat ttacatattt attcctttgc agttgaagga tgggatgtat 480
gcaaaggaga atgaactcta ctacagtaat ggaaagaagt gatagaagat gaaactccaa 540
aatgcctatc tgctacttga gtggaactca ctagaatata tcaaatgcag agagaaacat 600
gtgactaggc tctagtcaaa caattgtcat caaatacttg ctgaatatat aacacatttt 660
aggagctgt g 671

```

```

<210> 15
<211> 383
<212> DNA
<213> Homo sapiens

```

```

<400> 15
aaaaaatac attcctgtat ggaagactaa atattaaaga aaaaaacagt tatccatag 60
ttgtttcata gctttaagta tcctagtcta aaatccaaag ggactttttt ggggactcaa 120

```

aaaaaaaaaat tctaaagttg atccagaaga gtaaagagac taatatagtc aaaataatat 180  
 ttgaaaaggt taaataatta tgggaaaaca tgcttagcaa aaatataaac aaacaaaact 240  
 tcaaacagac aaaaacccta tgaaaaacca aaaacctatt taaaaactaa attctaaatg 300  
 agaactaaca tacaaatcaa gaatcagcat agatagctaa gaagcaaatt atagcatata 360  
 aactaattca atttattata aag 383

<210> 16  
 <211> 742  
 <212> DNA  
 <213> Homo sapiens

<400> 16  
 ctaggctgct cctgtgtggt tataatgaat ctcatcagct acaatcattt cccaaaaagg 60  
 acagaaaagt aatctgtctt ataaagatgg ttttcaaagc caactacaga tatcatagaa 120  
 aaaaaaaaaa aaacagtggc agataagacc ttcatcttt ttcttatccc atgggctttt 180  
 ctgcctccag atttccttgc acatcaaggg ggttctgtgc aaatcagtgg gcctggccct 240  
 ctgcccaccc tgggctaagc tgtggttgcc atgggtacga agacaagaat gacccttggg 300  
 tttatagagc cctgggtccc ttgctcacag gccttttcag ttgatatttc ttcatcttc 360  
 ttcagaaccc catgtggcag gtgcaacaag gagttcacca atgctccggc caaagtgagt 420  
 aatgagggac acatgcacga ccaggcagag cagccctgaa gaagtgggtt gtgaggggtg 480  
 ggaggaaggg ctgcttcca cactctgcac ttctgggtcc taaggcacta atcacacctg 540  
 gctcaggcat cattctcagc ctcatgctac ttttgtccag ggacagaaca agaggtgcca 600  
 gcactcaagg gaatcctgca ccacaacagg caggggacca tcatgactgc tgaaataggc 660  
 agcaggagtc cgaactatgg cataaaaatg tcacagcaac accacagggg ggaacccctg 720  
 gggggaagta ggggcatggt ga 742

<210> 17  
 <211> 228  
 <212> PRT  
 <213> Homo sapiens

<400> 17

Ala Thr Thr Ala Thr Cys Thr Cys Cys Ala Thr Thr Ala Thr Gly Cys  
 1 5 10 15  
 Thr Cys Thr Ala Thr Cys Ala Gly Thr Thr Thr Cys Thr Thr Thr Ala  
 20 25 30  
 Thr Ala Gly Ala Ala Thr Ala Thr Gly Ala Thr Thr Ala Cys Thr Ala  
 35 40 45

Cys Thr Thr Ala Gly Thr Gly Ala Thr Ala Ala Ala Gly Cys Thr Thr  
 50 55 60  
 Cys Cys Thr Thr Thr Gly Cys Thr Ala Ala Gly Ala Thr Thr Thr Cys  
 65 70 75 80  
 Ala Gly Cys Cys Thr Ala Cys Gly Ala Ala Cys Cys Ala Thr Gly Ala  
 85 90 95  
 Thr Cys Cys Ala Ala Ala Cys Cys Cys Thr Ala Cys Thr Thr Cys Thr  
 100 105 110  
 Ala Ala Ala Ala Cys Ala Thr Ala Thr Ala Ala Ala Cys Ala Thr Gly  
 115 120 125  
 Cys Thr Thr Thr Ala Cys Ala Ala Gly Thr Ala Thr Cys Cys Thr Ala  
 130 135 140  
 Thr Ala Thr Ala Thr Gly Gly Ala Ala Ala Ala Gly Thr Cys Cys Thr  
 145 150 155 160  
 Thr Gly Gly Ala Ala Thr Thr Ala Thr Thr Thr Gly Gly Gly Thr Ala  
 165 170 175  
 Ala Thr Thr Ala Ala Cys Cys Thr Gly Gly Thr Ala Thr Thr Ala Cys  
 180 185 190  
 Ala Thr Gly Thr Gly Thr Thr Ala Thr Thr Thr Cys Cys Thr Gly Ala  
 195 200 205  
 Thr Cys Thr Ala Cys Thr Ala Thr Thr Gly Cys Ala Ala Ala Thr Thr  
 210 215 220  
 Gly Ala Cys Ala  
 225

<210> 18  
 <211> 751  
 <212> DNA  
 <213> Homo sapiens

<400> 18  
 tcttgattta cacaaaaaac cataactaac taaaatagaa aacttttaat tatctgtatt 60  
 gttgtataaa atgttttata tattaggaat ctaaaaattt gttttttgct tttatcctct 120  
 gtcataaggaa gaagacatca tgtctctggt tactgtaacc attaataaag gctaataaca 180  
 gacagtacat gatgatactt ttaactaggg caaacaaaag taatatttta acaatgaggt 240  
 ttggctcttg ctatctatac ctcatgtcta attttcccta caatgtaaat gtcattcctc 300  
 ctctctaccc atttgtaagg gtctcagttt tctgctcttg catgacttat tttaaagggt 360  
 cacaataagg ccaggtaatt catattttta aatttccatt tagaataatt acatctaaaa 420  
 attcacaaga aagacaattt caatataaaa taataaatta ctaatattgg aatttcaagc 480

```

attagtcatg gcaaaaaaga gataatttgt agcagaatat tttaatggca actttcttat    540
tctatcactt attgtgttct atttgttatg accaaagaaa ttactctata tccactacaa    600
ttcataaaac aggcatggaa gaagtctttt ttttcttggg gctcatgtct aagaagatga    660
acctcagaag tatgtcattt ttcaatacta tgttctgaac agacagcaca cattatTTTT    720
gaatggacac caaatctcaa acatatatag a                                     751

```

```

<210> 19
<211> 657
<212> DNA
<213> Homo sapiens

```

```

<400> 19
gtcctaataa tcttcaatga attttaacta attttcaggg ataacaagc acttcagatt    60
gaagtcaatt catgtatact atttactcag gaacttttaa tttttcctat cacatacatg    120
ctgttgctgg gtttcttatt tgttaaaaga tatttcattc cctactgtgt ttaccctttg    180
ttagcaaagt tggtttagag tgacatagcc tgatgaaacc cataaaacag ccataaattg    240
ctcttatatg ggataaaaca atatttgaac actatatttc ttaaaaatat aatcttatat    300
tgggtgggta gaagtgatct tcacatcgtg tgtgtgtgtg tgtgtgtgtg tgtaatatat    360
aatatgaaag acttttaaaa gtaactttta aaatacatat ttttatatac atattttcat    420
acatatttac atacattttc tgttttcaat tcatctaggt ttacattaga ctatgtcctt    480
agtttgagtg ttaaaactat aaaaagagaa taaagttaca gcagaattaa ttgccaggga    540
tatgacagtt cgagcactac agtaaaaaat gagacacagg ttataaaaa catttaaatt    600
ctgaattttt gctttcttag gtttctctgt cagtaataga taattgttag tatgaaa      657

```

```

<210> 20
<211> 509
<212> DNA
<213> Homo sapiens

```

```

<400> 20
tttctaaatc tttgttacat tcttctttac tctgaggggt tttagtcac tgctagtaaa    60
ctagttagtc ctttccgtca gctgggtaaa aagctataaa tggttttcaa ctctatgcag    120
ctcccttggg ttgctcattt gatctgagcc aggtttgaag ttgggccatt gagaatgatc    180
tgtcctcatg atgtaggggg tgactccagc caattcttaa tattgctgga attaattaac    240
tttgattcct ctaactgaaa gagggcattt tactctaccg tgtattatta aataccttat    300
aacttgattt ttttgatttt gttttcttgc tcagtgahta cttaaaatat ccctgccgat    360
atagtaattt gttccctggt taatatggct ttttcctttg tttactctgg actcaagtct    420

```

ttgtagettg tttttctaag tcagctcaaa agttgctggc cattggattt ggcagccttt 480  
tcaagcttta cccaccagaa tcctacagg 509

<210> 21  
<211> 582  
<212> DNA  
<213> Homo sapiens

<400> 21  
ccccgccag tctccacatc ttgtaagtgg ggagataaaa cctatctaag ggggatatta 60  
tgaggagcag gagagaaggc atgtcattct cctaattaat tggcattcac taaagagtta 120  
tgtgattatt aaatacataa taaaatataa aaatataagg tgctctagtt atataatcat 180  
gaggtctgag gatggtgatg actgcacttc agtggactac cttatgaggg actgactgtc 240  
aagaagctct gtgcatgtgt gcttgtgtgt gtgcatctgt gtattatata tatacatata 300  
cggatatata ataggcactt ttattatata agaaaatgat gtcacagtaa aaatgcataa 360  
atattattga gcatgtttgt ataaacgagt gaacaaagag acatttggga tgcaagaoca 420  
gtggtataat ctgcccacac acaagccccg ttttactgtt gctaattgcac aaagagaggg 480  
ctggcagaca tggcctgcat gctctgtaca cctgcccata tgcaaccacac tcccccatgg 540  
tcttgcctgt gttgctgtgt tgggctcctc ccgcctctca ca 582

<210> 22  
<211> 307  
<212> DNA  
<213> Homo sapiens

<400> 22  
aggcagagat tatgtgagac tttgcatgat aggggtctgt gcattcttgc gcattttotta 60  
gctctgtgtg agcaatatatt aagtagtcat tacataaaat ctattgaacc tatacaaaga 120  
agtgaacaca ctgtgatgag ttggctccta ccaagtaagg cctggggggtt ggtcctttat 180  
ggttccccag cacacatctt aaaatagttc ctttgtggat aaactctctt tgaattattc 240  
tattttgagt gtgccatctg attcttgtta ggactctgac tttaaaaagg tagagcattg 300  
ctctcat 307

<210> 23  
<211> 633  
<212> DNA  
<213> Homo sapiens

<400> 23  
caaatggatt aatttacata tggatgaaca tgatctgttg ctctccaggt cccaaaggat 60

```

acataagaag aaaaatttag tgatgttact ggatgatgtc ttttaagaca acacaataca 120
atatctgagt atgtaccctt acgacataga gaagggattt tcaaaatatt ttaacttaaa 180
tagattcact aaaagaaatc accttccaac cactgttctt tgtctctggt caattagggt 240
cataatattg ttttcattgt attacaaaag gtaagaatgt aactgtata aatgaataaa 300
taatatagat tactagataa gcagataaat aaatacaaaa gcacaaaaat acaaaagcaa 360
atgaccactc aactcacttc caccacttc aaggaaaaca cagttccttt atcatagtta 420
ctattagaag tctttctgtt cttcattttc ctacaagcat gcagaaatat atatgtggac 480
acatttgtgc cagaattcct tttttctga cactcacttt tttcctccta ctccacaata 540
tgtcaggaca attttctaca agatatagca aatggagtaa catagaatag agcaaaacat 600
gaaaacctca aactcattag tggatgatgt ttt 633

```

```

<210> 24
<211> 674
<212> DNA
<213> Homo sapiens

```

```

<400> 24
gggaaggctc ttctaagaaa ccacgccac acacaaatta gtaaattoga caaaacagga 60
aacaacaaaa aatttttgtt taaaagtaac actaaatcaa aatgaaaaga taaaaggcaa 120
acacaaaatt gacaaggatg cttgcaactt gtgtaactga taaagggcta ttttcctgaa 180
tataatttctc tattctataa ttaotttctt taatatattt taaaaacttc tgtaagtaag 240
tatgaaatag acaaatgata agaacagtta tcagaaaatg aaatacaaat gattcaaaca 300
tgaaaagatg ctcgagctca gtctataata atagaccctt aaaagtataa tgaaatatct 360
ttttttaaac ccctatctga ttagcaaaga ccagcaagtt ggaaaaacag agggcttttt 420
caaagatcaa gcttgtgagc caaactaaag gattctgtgc tttcaagaat tacactgttt 480
agagtttgga ctttttgaaa aaaatgtaca tgtctatgaa ataaattggc ctttttaaaa 540
agagtttgca gcaatgctga aagtagtggc caaagcatg atcaagagat aggatattca 600
tctaattctc aaccatctcc ctacaagcta tttatcaatt acaaagtga aaataactaac 660
tgcacagtgg tgaa 674

```

```

<210> 25
<211> 600
<212> DNA
<213> Homo sapiens

```

```

<400> 25

```



```

acaaaagtca ctctctgct caaatcttt gcacggtttc ctgctgtact tggaatacaa      60
ccccaagtcc tttcctaggg cactagcctt gtattggcct tgatcactgc tcctgcccac    120
ctcataccac ttgtcatttg gccattatag tccaacaaca tccactcatc cgctttctcg    180
caggccctgg aatattctaa gcactttccc atctcagggg gttgtgaatg ctgtttcctc    240
tacctgctct catcctccca ctctttatca actaagttct cacatacaca tgtttgatct    300
caagctgtgt gtgtttactt gtttgttatt tgaagcacta tgctgagcca tagtggaatc    360
tgaggtaaaa ggaaaaataa gtaatattga ctctgagttt atgtaatatt ttgatttttt    420
cccatgaagg aattttgcat taatttttat tttaaaaata ttgtattaaa atttttattta    480
tcttgattac tgagtatttt ggtgtcccct taaaattctc accaaaggac aagtgcctcg    540
ctcacctcac cttttttcca gtactgggtc ttggtctgtc ccaccaaact gtaagtaaac    600

```

<210> 26  
 <211> 709  
 <212> DNA  
 <213> Homo sapiens

```

<400> 26
ccgagggtt tccctgtgag aatcagtttc aaatatgagg ttgattttat tttccccggt    60
tgcaattaac tgaaaaaaat atttcaaagc tgtataatta aaggccggca ttttattagg    120
gtttaaaaata gtcgctatga aatttgctcc tgccattggg aattacaacg tacatcttaa    180
tatccccctg cttgccagcc ccgagataac agcgtgtgtg gtgctgcatt tggctttgca    240
gagaaggtaa aatccttata ttatctccat ctgcacacaa caaccagcct ctcttccaat    300
tcaaccctaa tcttcttagt aaataatact gatatttttc accataacaa aatggagctg    360
gtgccagatt ctactgata acacctctcg tttacacacc attctctgtc aaaatcggcc    420
cgatttttct taatctcttt aacttatcaa aaaaaatact ttttacattt ggccatagta    480
agttcttttc taccaaataa taatattata aataaatccc tttgcctaca gctgactcat    540
tgaaaagata ctacgcctcc tcattttctt catgaatgga aattgttctg ctcttcaatt    600
ttcctcccaa gtttgggggt ccacaccaag ggaattgttc ccaggcaggc cctgagccat    660
ctttgtatct tgcagggggg catcttggtc caggatgtgc atgcttctc                709

```

<210> 27  
 <211> 397  
 <212> DNA  
 <213> Homo sapiens

```

<400> 27
agtataaaat ctgggagaaa aatccatcaa attataaaat agaactaata atgctttcat    60

```

gggtaagcaa tgaacaaaac caaaggtaaa ccagcgTTTT aaagattgat atttcttttaa 120  
 gaagagaatg gaaaataaat taaaatatac attttatatt gattatcatg aagtttttct 180  
 agtttcttac ctgaagtttc ctggaggaac ttttaacagca ctcatttgac attaggaaga 240  
 agccgaatgt aaataaatcc taggtggata agcataatat aatgagtaat atgatcttaa 300  
 atggaaaaaa gattatgtaa agttgtttat actcttaaca ttttccaccc actacactgt 360  
 gaagcagatt aaccttttgt caaaggtgta accttct 397

<210> 28  
 <211> 537  
 <212> DNA  
 <213> Homo sapiens

<400> 28  
 ctttgaagga tgaagagaag ttcaccaagt gaacgtggag gaaaagcagt ccaggcagaa 60  
 ggaacagctg ctgaaaaggt caggtgccca aaccagaaag gcatgtgagg gaattcatgg 120  
 gcagttcttg agaaccagag cagaaggaga caggctagag aatggcagga gatagcccta 180  
 gaaagctcta tgtgtgccag gtgcaaaagg cttgtgcgcc agctaaggag tctgcacttt 240  
 cccctgtaag caattaaaag ttttaagcag aggagggatt ggtttgtatt gatctagata 300  
 attcagggaa caatgaaggg aggatatttg aggaatccca aagagcaggg gcagagactc 360  
 tcatcacagt tcaggcctga ccgaaggcca tagggagata gcagtaaagc agggaaacct 420  
 actgtccctt cttcacctgg aaaactatca ccttccagag ttctgtcac cccagtgaa 480  
 ggcaaccact gtctcttctg ttacctaaca cgtggtacaa acagctacca tagcatg 537

<210> 29  
 <211> 578  
 <212> DNA  
 <213> Homo sapiens

<400> 29  
 tataaggtag ttgtgtagct ttaagactaa gacctctaca cctggaaatt ggtatacttt 60  
 ttcaataaag catcagatag taaagttttg tgggccatat gtctctgctg caactactca 120  
 gctctgcat tgtgcaaaag cagccacaga taatatgtaa atgaatgaac ttggctgttt 180  
 tcacaaaaac aggcggctgt taggatttgg cttgctgact ctttaacttg attctcactt 240  
 ctgaaagcat ggtctaggag accagcagca tcagcattac ctgtgaacct gttagagata 300  
 cagaatgttg agccccttct catactgatt gctctagagt ctaaaccatg tttctcaaac 360  
 ttcagcatac ataggactca cctgggggac ttgtgaaagt acagactctg agatagcagg 420

gctgaacacc atactagcct tggggaggca atgttatgac ctggtcagtg ttccttgggg 480  
 agtccgcaat ctattaggag agactgctat ataaacaaat aatttcaata cagactagta 540  
 aatgctataa cggagggtatc ttcagcagag gagcacca 578

<210> 30  
 <211> 589  
 <212> DNA  
 <213> Homo sapiens

<400> 30  
 attttcatgt tgatggcaat catccaagag agagcaggcc atgtcacgca ataatagaac 60  
 aaggagaaat gcacactttt ctttttccat gtctctgctt gcattacatt tgctaatacc 120  
 actttagcca gataaattaa cacagcgcag cacatagtca aaaaaataga agaccagaag 180  
 ttacagaaca actgacatgg tttagaaaag caattaagac catcagtga atcagtctac 240  
 cacaaaaggt aattcatttg ttcacaacgc tgttgaaaag acttatcttc caagacagga 300  
 aatggttctc cactgaaggg tgaagacatt tcaattttca gtcatttggg gaagagttgg 360  
 atctccaacg agtaactttc atgcaaggac aagaatttag tagtgaaata gaggttattc 420  
 gtttttttac cataaaataa ttaataatct tggaggcagt ttcctcatag cagttattat 480  
 ggcagttgtg ttcatttaca ggaaaactga gaaactctaa gatgtttttg ggaaaaaaaa 540  
 gtattttgaa agcttgcgag tgttaacttc cacaatagat atactcttc 589

<210> 31  
 <211> 1750  
 <212> DNA  
 <213> Homo sapiens

<400> 31  
 cgcacagcgc gcaggctctc accagagctc tgggtggccac ctctgtcccg ccatgctgct 60  
 caccgacagt ggccagggcc cacagcacca agaggcttgg gccacaaagt aaagggtcgc 120  
 ggagcctcgc cggccgccat gtggagctgc agctgggttca acggcacagg gctgggtggag 180  
 gagctgcctg cctgccagga cctgcagctg gggctgtcac tgttgtcgt gctgggcctg 240  
 gtggtgggcg tgccagtggg cctgtgctac aacgccctgc tgggtgctggc caacctacac 300  
 agcaaggcca gcatgaccat gccggacgtg tactttgtca acatggcagt ggcaggcctg 360  
 gtgctcagcg ccttggcccc tgtgcacctg ctcggccccc cgagctcccg gtgggcgctg 420  
 tggagtgtgg gcggcgaagt ccacgtggca ctgcagatcc ccttcaatgt gtcctcactg 480  
 gtggccatgt actccaccgc cctgctgagc ctcgaccact acatcgagcg tgcactgccg 540  
 cggacctaca tggccagcgt gtacaacacg cggcacgtgt gcggcttcgt gtgggggtggc 600

gcgctgctga ccagcttctc ctcgctgctc ttctacatct gcagccatgt gtccacccgc 660  
 gcgctagagt gcgccaagat gcagaacgca gaagctgccg acgccacgct ggtgttcatc 720  
 ggctacgtgg tgccagcact ggccaccctc tacgcgctgg tgctactctc ccgcgtccgc 780  
 agggaggaca cgcccctgga cggggacacg ggccggctgg agccctcggc acacaggctg 840  
 ctgggtggcca ccgtgtgcac gcagtttggg ctctggacgc cacactatct gatcctgctg 900  
 gggcacacgg gcatcatctc gcgaggggaag cccgtggacg cacactaact ggggctactg 960  
 cactttgtga aggatttctc caaactcctg gccttctcca gcagotttgt gacaccactt 1020  
 ctctaccgct acatgaacca gagcttcccc agcaagctcc aacggctgat gaaaaagctg 1080  
 ccctgcgggg accggcactg ctccccggac cacatggggg tgcagcaggt gctggcgtag 1140  
 gcggcccagc cctcctgggg agacgtgact ctgggtggacg cagagcactt agttaccctg 1200  
 gacgctcccc acatccttcc agaaggagac gagctgctgg aagagaagca ggaggggtgt 1260  
 ttttcttgaa gtttcctttt tcccacaaat gccactcttg ggccaaggct gtgggtccccg 1320  
 tggctggcat ctggcttgag tctccccgag gcctgtgcgt ctcccaaaca cgcagctcaa 1380  
 ggtccacatc cgcaaaagcc tcctcgctt cagcctctc agcattcagt ttgtcaatga 1440  
 agtgatgaaa gcttagagcc agtatattata ctttgtggtt aaaatacttg attccccctt 1500  
 gtttgtttta caaaaacaga tgtttcctag aaaaatgaca aatagtaaaa tgaacaaaac 1560  
 cctacgaaag aatggcaaca gccagggtgg ccggggcctg ccagtgggog gcgtgtgcta 1620  
 gcaaggcctg ccgggtgtgc cgcagtcacc acagggttct gagaacattt cacagaagtg 1680  
 cctgagacgc ggagacatgg ctggtgttaa atggagctat tcaatagcag tgacgcgctc 1740  
 tcctcagcca 1750

<210> 32  
 <211> 551  
 <212> DNA  
 <213> Homo sapiens

<400> 32  
 ccacaagaca gtgccattta gtggcagccc taggataaag atgatactgt aggccaggga 60  
 gaggtagact tgcttgact tctctgagaa ctggcagaga ccttgttcct gtgatgtatt 120  
 catgtccacc ttctccatgt cccgggaggc tcctccagg agcagagctc cagcagggct 180  
 cccgcttctg cttccccctg aaggaagcaa aatggacagc ataacatcat gacctcagca 240  
 tggcatcagc ctctgcccc acccggcatt caactttgtc ccccaaagtc tagtcagagg 300  
 ccttgtttaa aagagaccaa cccagggaaa atagagtatt caacaagggc caaataatct 360

tccaaagctg ggcgtctaga cttaccaggt ccaaaggggtt ggccttacat ggacttgtat 420  
 gctgtctact ctaggctggg ttgtgataga aatgggtggg ggatgacact gtcattttaa 480  
 cagggtactc agaacctcca gctctgtgtc tctcctgct agaateccatt gcatagagtc 540  
 agaaaatctc c 551

<210> 33  
 <211> 426  
 <212> DNA  
 <213> Homo sapiens

<400> 33  
 catcctaagc tcgggcacac aaagtcccag aggcacacag attcagtggc ttggggatag 60  
 catattccaa ctcttgggcc cctaactcaa aatccccaac aacctcagt cacctgtggc 120  
 tccagtcctg ttttggtttt ccacagatca caaaggcatc aaggaagagg ctcacggtaa 180  
 gcctggccta ctcgagagc caccagatcc gcgtgtccca gcaggacttc cggctcttcc 240  
 gcacctctt ccttctaatt gtctccttct tcatcatgtg gagccccatc atcattacca 300  
 tctcctaatt cctgatccag aacttcaagc aagacctggt catctggccg tccctcttct 360  
 tctgggtggg gggcttcaca ttgctaatt cagccctaaa cccatcctc tacaacatga 420  
 cactgt 426

<210> 34  
 <211> 537  
 <212> DNA  
 <213> Homo sapiens

<400> 34  
 gctgggaagt gtcctgggtg cagccggcgg tagatggcgg agggcacggt gagcagcaag 60  
 gccagtgtcc aggtgtcccc acaggccacc tgcacccgc acgcccgtg aacctatac 120  
 caccaggcag gcccgagagc caggaagcag aggtcggcac tgagagctgc caggagcagg 180  
 acgtggcat acatggtcag caggatgatg gagggcagcg cccgacagcc cactgcacca 240  
 tacggtcagt ggctccacg ggcaatgggc actgccagga tgggcagaga caaacagcac 300  
 agcaaattccg ccacggccag gtggagcaac caggtggcac ccacctccg gcgggccacc 360  
 ttcccagcca cccaggccac catggcattg cccggcacc ccaccaggaa gatggcggca 420  
 tacagtggga gcggggccac gcgcagcggg tcgatggcca ggcaggcgcc atccaggcag 480  
 tccacagggc ggtccgagag gtcgctgtaa tcccatact cgtagctgac agaatcg 537

<210> 35

<211> 536  
 <212> DNA  
 <213> Homo sapiens

<400> 35  
 gggtccactg gccttttcggc aaggggctct gcaggctgac ggcgtttgtg ctctacaccg 60  
 acacctacgg ggggtctacc tcatggcctg tgtgagcgtg gaccattacc cagctgtgggt 120  
 ctgtgcccac tggggcccgt gcctccgcac ggctggccgc gccaggctgg tctgcgtggc 180  
 catctggacc ttggtgctgc tgcagacgat gcccttgctc ttgatgcca tgaccaagcc 240  
 gctggtgggc aagctggcct gcatggagta cagcagcatg gagtcagtcc tcggggctgc 300  
 ccctcatggg cctggtggcc ttgccattg gcttctgtgg gccagtgggg atcatcctgt 360  
 cctgctatat gaagatcacc tggaagctgt gcagcacagc tgggagaacc cagtgaccag 420  
 cgggaaagga caccaccggc ggggcagccc gggaggaccc agtgaccagc aggaaaggac 480  
 gccaccggcg gggcagccca ggaggacca gtgaccagcg ggaaaggaca ccaccg 536

<210> 36  
 <211> 1449  
 <212> DNA  
 <213> Homo sapiens

<400> 36  
 gaggcaggct gcagggaagt aaggaggagg catggcacct tctcatcggg catcacagggt 60  
 ggggttttgc cccacccctg aacgccctct gtggcgctt ccacccacct gtaggcccag 120  
 aaggatgtcg gtctgtacc gtccccagg gaacgagaca ctgctgagct ggaagacttc 180  
 gcgggccaca ggcacagcct tcctgctgct ggcgcgctg ctggggctgc ctggcaaccg 240  
 ctctgtggtg tggagcttgg cgggctggcg gcctgcacgg gggcgaccgc tggcgccac 300  
 gcttgtgctg cacctggcg tggccgacgg cgcggtgctg ctgctcacgc cgctctttgt 360  
 ggcttcctg acccggcagg cctggccgct gggccaggcg ggctgcaagg cgggtgtacta 420  
 cgtgtgcgcg ctacagcatg acgccagcgt gctgctcacc ggctgctca gcctgcagcg 480  
 ctgcctcgca gtcacccgcc ccttcctggc gcctcggtg cgcagcccgg ccctggcccg 540  
 ccgcctgctg ctggcggtct ggctggccgc cctgttgctc gccgtcccgg ccgcgctcta 600  
 ccgccacctg tggagggacc gcgtatgcca gctgtgccac ccgtcgccgg tccacgccgc 660  
 cgccccctg agcctggaga ctctgaccgc ttctgtgctt cctttcgggc tgatgctcgg 720  
 ctgctacagc gtgacgctgg cacggctgcg gggcgccgc tggggctccg ggcggcacgg 780  
 ggcgcgggtg ggccggctgg tgagcgccat cgtgcttgcc ttcggcttgc tctgggcccc 840  
 ctaccacgca gtcaaccttc tgcaggcggg cgcagcgtg gctccaccgg aaggggcctt 900

```

ggcgaagctg ggcggagccg gccaggcggc gcgagcggga actacggcct tggccttctt 960
cagttctagc gtcaaccccg tgctctacgt cttaccgct ggagatctgc tgccccgggc 1020
aggccccgt ttcctcacgc ggctcttcga aggctctggg gaggcccgag ggggcggccg 1080
ctctagggaa gggaccatgg agctccgaac taccctcag ctgaaagtgg tggggcaggg 1140
ccgcggcaat ggagacccg ggggtgggat ggagaaggac ggtccggaat ggacctttga 1200
cagcagaccc tacaacctgc tgcccttccc tgctccttcc cccccccac ccacctcca 1260
gaggtcagtg ttctgggaca tttggggacc cttctttgac tagagtttgg atctggctgg 1320
gtaggattac tatacacttg gggcaggccc aggctcctcc aaactgaggg attatgaggg 1380
tggtgatggt ccctgttaag gactattgtg tgcttgcaag ttggcatgta cccatgtgcc 1440
agcattgct 1449

```

```

<210> 37
<211> 635
<212> DNA
<213> Homo sapiens

```

```

<400> 37
ccagccgtcc aggcgacgcg ggccagcacc aggaaccagg tgacgctggc cacgtggtag 60
cgcgggcaga ccagggggca cgcgctgttg cgcagcaggc cgcaggcggt ggcgggcagc 120
ggcacggccg gcaggggtcg ggtccacacc agggcgcaga ggacggccga ggcgtgtctg 180
ggccggcagc cctggttagca ggcggggaag aggtcggaga ggcagcgctc cacgtgaag 240
gccgccagca gccagagccc caccgcgaac cacaggaagg tgagcacgaa gtagagtgtg 300
tcctgggcgc ccagggcagc ctgagccacg gagaagccca cacggcagga gaggaacagg 360
aagtcggcgg cggccagggt cagcaggtag atggagaagg ggcccttctt gatgcggaag 420
ccgaggttcc agagcaccag cccgttacct accggtcccc cgaggccac gatcagcgtc 480
aggtagaaga ccacactgtc gaaggttctc cagaggccga acagcccaaa catcctggcc 540
ggctcagagg gtgctggcga gggacctgca agatggagac agacatggtc ctcaggagtc 600
ttttgtgccc ccgctggggg cacagggagt ggcac 635

```

```

<210> 38
<211> 564
<212> DNA
<213> Homo sapiens

```

```

<400> 38
atgtctacag aaaccccttc gccatctacc tcctggtacg tggcctgcag caggatctca 60

```

```

tcttccttgg ctgccacatg gtggccatcg tccccgactt gctgcaaggc cggctggact    120
tcccgggctt cgtgcagacc agcctggcaa cgtgcgctt cttctgctac atcgtgggcc    180
tgagtctcct ggcggccgtc agcgtggagc agtgcctggc cgccctcttc ccagcctggg    240
actcgtgccg ccgcccacgc cacctgacca cctgtgtgtg cgccctcacc tgggcccttt    300
gcctgctgct gcacctgctg ctcagcagcg cctgcaccca gttcttcggg gagcccagcc    360
gccacttgct cgggacgctg tggctggtgg cagcgggtgct gctggctctg ctgtgttgca    420
ccatgtgtgg ggccagcctt atgtgtgtgc tgcgggtgga gcgaggcccc cagcggcccc    480
caccgccggg cttccctggg ctcctccttc ctcaccgtcc tcctcttctt cttctgcggc    540
ctgcccttcg gcattctactg gctg                                           564

```

```

<210> 39
<211> 187
<212> PRT
<213> Homo sapiens

```

```

<400> 39

```

```

Val Tyr Arg Asn Pro Phe Ala Ile Tyr Leu Leu Val Arg Gly Leu Gln
1          5          10          15
Gln Asp Leu Ile Phe Leu Gly Cys His Met Val Ala Ile Val Pro Asp
20          25          30
Leu Leu Gln Gly Arg Leu Asp Phe Pro Gly Phe Val Gln Thr Ser Leu
35          40          45
Ala Thr Leu Arg Phe Phe Cys Tyr Ile Val Gly Leu Ser Leu Leu Ala
50          55          60
Ala Val Ser Val Glu Gln Cys Leu Ala Ala Leu Phe Pro Ala Trp Tyr
65          70          75          80
Ser Cys Arg Arg Pro Arg His Leu Thr Thr Cys Val Cys Ala Leu Thr
85          90          95
Trp Ala Leu Cys Leu Leu Leu His Leu Leu Leu Ser Ser Ala Cys Thr
100         105         110
Gln Phe Phe Gly Glu Pro Ser Arg His Leu Cys Arg Thr Leu Trp Leu
115         120         125
Val Ala Ala Val Leu Leu Ala Leu Leu Cys Cys Thr Met Cys Gly Ala
130         135         140
Ser Leu Met Leu Leu Leu Arg Val Glu Arg Gly Pro Gln Arg Pro Pro
145         150         155         160
Pro Arg Gly Phe Pro Gly Leu Ile Leu Pro His Arg Pro Pro Leu Pro
165         170         175

```



Leu Leu Arg Pro Ala Leu Arg His Leu Leu Ala  
 180 185

<210> 40  
 <211> 544  
 <212> PRT  
 <213> Homo sapiens

<400> 40

Cys Ala Ala Thr Thr Thr Thr Cys Thr Ala Thr Thr Gly Cys Cys Thr  
 1 5 10 15  
 Cys Thr Cys Thr Gly Gly Cys Cys Thr Gly Thr Gly Cys Thr Gly Ala  
 20 25 30  
 Cys Thr Thr Cys Thr Thr Gly Gly Thr Ala Gly Gly Thr Gly Thr Gly  
 35 40 45  
 Ala Cys Thr Gly Thr Gly Ala Thr Gly Cys Thr Thr Thr Thr Cys Ala  
 50 55 60  
 Gly Cys Ala Thr Gly Gly Thr Cys Ala Gly Gly Ala Cys Gly Gly Thr  
 65 70 75 80  
 Gly Gly Ala Gly Ala Gly Cys Thr Gly Cys Thr Gly Gly Thr Ala Thr  
 85 90 95  
 Thr Thr Thr Gly Gly Ala Gly Cys Cys Ala Ala Ala Thr Thr Thr Thr  
 100 105 110  
 Gly Thr Ala Cys Thr Cys Thr Thr Cys Ala Cys Ala Gly Thr Thr Gly  
 115 120 125  
 Cys Thr Gly Thr Gly Ala Thr Gly Thr Gly Gly Cys Ala Thr Thr Thr  
 130 135 140  
 Thr Gly Thr Thr Ala Cys Thr Cys Thr Thr Cys Thr Gly Thr Cys Cys  
 145 150 155 160  
 Thr Cys Cys Ala Cys Thr Thr Gly Thr Gly Cys Thr Thr Cys Ala Thr  
 165 170 175  
 Cys Thr Gly Cys Ala Thr Cys Gly Ala Cys Ala Gly Gly Thr Ala Cys  
 180 185 190  
 Ala Thr Thr Gly Thr Gly Gly Thr Thr Ala Cys Thr Gly Ala Thr Cys  
 195 200 205  
 Cys Cys Cys Thr Gly Gly Thr Cys Thr Ala Thr Gly Cys Thr Ala Cys  
 210 215 220  
 Cys Ala Ala Gly Thr Thr Cys Ala Cys Cys Gly Thr Gly Thr Cys Thr  
 225 230 235 240  
 Gly Thr Gly Thr Cys Gly Gly Gly Ala Ala Thr Thr Thr Gly Cys Ala  
 245 250 255  
 Thr Cys Ala Gly Cys Gly Thr Gly Thr Cys Cys Thr Gly Gly Ala Thr  
 23

260						265						270					
Thr	Cys	Thr	Gly	Cys	Cys	Thr	Cys	Thr	Cys	Ala	Cys	Gly	Thr	Ala	Cys		
		275					280					285					
Ala	Gly	Cys	Gly	Gly	Thr	Gly	Cys	Thr	Gly	Thr	Gly	Thr	Thr	Cys	Thr		
	290						295				300						
Ala	Cys	Ala	Cys	Ala	Gly	Gly	Thr	Gly	Thr	Cys	Ala	Ala	Thr	Gly	Ala		
305					310					315					320		
Thr	Gly	Ala	Thr	Gly	Gly	Gly	Cys	Thr	Gly	Gly	Ala	Gly	Gly	Ala	Ala		
				325					330					335			
Thr	Thr	Ala	Gly	Thr	Ala	Ala	Gly	Thr	Gly	Cys	Thr	Cys	Thr	Cys	Ala		
			340					345					350				
Ala	Cys	Thr	Gly	Cys	Gly	Thr	Ala	Gly	Gly	Thr	Gly	Gly	Cys	Thr	Gly		
		355					360					365					
Thr	Cys	Ala	Ala	Ala	Thr	Thr	Ala	Thr	Thr	Gly	Thr	Ala	Ala	Gly	Thr		
	370						375				380						
Cys	Ala	Ala	Gly	Gly	Cys	Thr	Gly	Gly	Gly	Thr	Gly	Thr	Thr	Gly	Ala		
385					390					395					400		
Thr	Ala	Gly	Ala	Thr	Thr	Thr	Cys	Thr	Gly	Thr	Thr	Ala	Thr	Thr			
				405				410					415				
Cys	Thr	Thr	Cys	Ala	Thr	Ala	Cys	Cys	Thr	Ala	Cys	Cys	Cys	Thr	Thr		
			420					425					430				
Gly	Thr	Thr	Ala	Thr	Gly	Ala	Thr	Ala	Ala	Thr	Thr	Cys	Thr	Thr	Thr		
		435					440				445						
Ala	Cys	Ala	Gly	Thr	Ala	Ala	Gly	Ala	Thr	Thr	Thr	Thr	Thr	Cys	Thr		
	450				455					460							
Thr	Ala	Thr	Ala	Gly	Cys	Thr	Ala	Ala	Ala	Cys	Ala	Ala	Cys	Ala	Ala		
465					470				475						480		
Gly	Cys	Thr	Ala	Thr	Ala	Ala	Ala	Ala	Ala	Thr	Thr	Gly	Ala	Ala	Ala		
				485				490					495				
Cys	Thr	Ala	Cys	Thr	Ala	Gly	Thr	Ala	Gly	Cys	Ala	Ala	Ala	Gly	Thr		
			500				505					510					
Ala	Gly	Ala	Ala	Thr	Cys	Ala	Thr	Cys	Cys	Thr	Cys	Ala	Gly	Ala	Gly		
		515					520				525						
Ala	Gly	Thr	Thr	Ala	Thr	Ala	Ala	Ala	Ala	Thr	Cys	Ala	Gly	Ala	Gly		
	530				535						540						

<210> 41  
 <211> 595  
 <212> DNA  
 <213> Homo sapiens

<400> 41

```

attattcatt ccttttagaca ttaaaccattc attgagcacc tgctgtatgc aaagcactgg      60
gcacccacac taaggatgaa aaccaaggat aagtaagaca gcatgtagat tctagctgcc      120
tggttgagag gttacagtca gaaagtcctt atagccattc agtgacatac agatagggat      180
aggagaggaa atgggggtgag cacaggggaag ggataagtat agggtcaggg ctaccaccct      240
tcttttcccc atgaccccat gggggcaa atgtcctgtc tcttcctggg tgggtgatgt      300
tactgcccc tttctccatt ccaattggaa cttctagatt gagcccgaag ctagacttgc      360
agatcacaat tttaagaaag ttggatgttc tgcaacaggt tacaacaggag ctatcattga      420
gtcatctttt cgttcaccca atcattttatt cattcattca ttcaacaagt atttcccaag      480
caccaattcc atcctcaaca aaagactcct aaatcacagg ccagatgaaa gacgtcacta      540
ttcacttggg aaaattcagc tgattatgat cctggtcgga gtgtactcag cgtcc          595

```

<210> 42  
 <211> 651  
 <212> DNA  
 <213> Homo sapiens

```

<400> 42
atggaattag aaaaatgaga aagtaatcaa agattgtaaa aaatactgag atagaattaa      60
aaggaataaa aaatttgtaa atcaaatatg taattctttt taagttgaca cattgtaatt      120
ttatatattt gtggagtatg atttggtaca ttaatttgta tatgtgttgt atgctgatca      180
aatcagggta tgtactaggc catTTTTgca ttggtataaa gaaatacctg aagctgggta      240
atttaaagaa aagaagttta ttgactcat ggttctatag gctgtgaaca acatctgctt      300
ggcttctggg gggagcctcg gaaagctttc aatagtggca gaaggggaag gggaagctgg      360
agtatcacat ggtgtgaaag ggagcaagag agagagagag agacaggagg tcctagactt      420
ttaaacaacc agatcttgtg tgaactatTT catgtgacta agaactcatt catcaccaag      480
cggaagggtg caagtcattc atgagggatc caccctcata atacactacc ttccaccaag      540
tctcacttac aacattggga atgacatttc agcataagat ttggagagga caaatgtcca      600
aactatatca gggtagttag ctcatcattc acctcatgtc tttatcattt c          651

```

<210> 43  
 <211> 668  
 <212> DNA  
 <213> Homo sapiens

```

<400> 43
atgtaaaggg ccttttggaa gaaaactgta aactccactg tggggccatt aaagaagtct      60
tgaatacata gattaaatcc cttgggttga gataataaaa ctctcaatat ttaccatgct      120

```

```

tttgtcattt aaatacaaat tgtggacatt tttaccaggg caacagatga gttacaacca 180
tccttttatc agttgcacag aatcttattg aatgggtata tcctacttat gtcagttctt 240
agtagtgaat attaggttgt ttctaatttt tccatattag aaacaatggt aaaagaactt 300
gctcttattt gtgtttctta tgtctacact agtattcaat acctttctag gagtggaaat 360
tcctggctca aacaaactgc cttccaagaa gatttactaa tatacattcc tactaatgct 420
gttttagaat gcctttttct ccatattcct atgaacactg agtggtctct ttttaacact 480
ttttttcttc aatctactag actccttcta aaatatatac ttttcaaac gcctgcttac 540
ccttcaatat attgtctttt aaaagattat ttgtaggtac tgtttatgtg ttaggttggt 600
ggaaaattaa ttgtgatttt gcaaaaactg caattacttt tgcaccaacc taatattata 660
catgtaga 668

```

```

<210> 44
<211> 604
<212> DNA
<213> Homo sapiens

```

```

<400> 44
tctcttttgg ttcccttttt tcctcattac ctacctcttc tcctttgctg gtccttcaca 60
tcttcctggc ctcttctttg tctgcattta cttgtgtggt cttccgtctt taacacatct 120
acctatcagc tactcccaaa tgtatatatc tattcccgga cctttcctct gaactccaga 180
tttgtatttc caactaccta ctcaatagta ccttcttgga tatttattaa aatttgaatc 240
acgacatgcc taaaattgaa cttcctatct ctgaagcaaa gccctatcct tctattgttt 300
ttcccatccc aataaattgc cactcattct tcagatgct tgggcaacat ttttgcagtc 360
atctttgatt tccttctttc tttgatatcc cacatacatt caccagcata tgctgccctg 420
tctacatttt aaacataacc agcatccaac atttctacaa ttgctactag ccttgtatag 480
ccctatcctc tctcctctgg attactgtat tagcctctga actgggatcc cgcttcaacc 540
ctttccaacc tcctcctccc aaagccaact ctgtgaacag cagctaaagt tgttctttca 600
aaaa 604

```

```

<210> 45
<211> 578
<212> DNA
<213> Homo sapiens

```

```

<400> 45
caaccctggt aaagcatcgc accttggggt atgtacctca gggttatttg acgcactggg 60

```

```

ctaaaatggt gaaggacatc ctgtttccag gtgggggactg gaacagagcc tggactgttt 120
tagccaatgg ctcccttacct caggatgttg cattcccagc acattctggt tgggtgttgag 180
aactacaaac aagaaagtgg gaagaactgt tctgcaccat ttatgtaaac ttctaggaaa 240
gcaaactaat gtattgtgac agaaggaaga tgagtgattg cttaaacaca agggaggaat 300
gctacacagg gaaggtctgg agagttggat tacaaagaga gacaaggaca cttttgggag 360
agatggatat gttttattatc ttgcttatgg tgatagtttc ataggtccat aaatcccaaa 420
agcatctcat tgtacactaa ctatgtatga tttatttata ttaattatac ctcaataaag 480
ttgtttttta aaaaagttac cacttaatcc tgtaaataga ccaggaaggc aattaattaa 540
tattttctag tttacttttg agaaactaaa gcttagta 578

```

```

<210> 46
<211> 575
<212> PRT
<213> Homo sapiens

```

```

<400> 46

```

```

Ala Gly Thr Thr Thr Ala Thr Gly Gly Ala Cys Cys Ala Gly Cys Cys
1          5          10          15
Thr Thr Cys Cys Cys Thr Gly Thr Gly Ala Ala Ala Thr Thr Thr Gly
20          25          30
Ala Cys Thr Thr Thr Thr Cys Cys Cys Thr Cys Thr Thr Thr Gly Cys
35          40          45
Thr Gly Ala Ala Thr Thr Gly Gly Thr Cys Ala Gly Gly Thr Thr Ala
50          55          60
Ala Cys Ala Ala Thr Gly Gly Thr Thr Ala Cys Cys Cys Cys Thr Gly
65          70          75          80
Gly Ala Thr Thr Ala Cys Ala Gly Gly Ala Ala Gly Gly Gly Cys Ala
85          90          95
Thr Gly Thr Gly Cys Thr Ala Ala Ala Ala Gly Cys Cys Thr Cys Thr
100         105         110
Thr Thr Gly Gly Ala Gly Ala Cys Cys Cys Ala Cys Ala Thr Gly Gly
115         120         125
Cys Cys Cys Thr Cys Ala Gly Ala Thr Gly Ala Gly Cys Ala Ala Thr
130         135         140
Thr Gly Thr Thr Cys Ala Gly Ala Thr Thr Cys Cys Thr Thr Thr Thr
145         150         155         160
Cys Thr Thr Thr Thr Thr Cys Thr Thr Thr Thr Cys Cys Ala Thr Gly
165         170         175

```

Gly Gly Ala Ala Thr Ala Ala Gly Cys Thr Thr Thr Cys Cys Thr Cys  
 180 185 190  
 Thr Cys Thr Cys Cys Ala Ala Ala Gly Thr Ala Cys Ala Thr Gly Thr  
 195 200 205  
 Thr Thr Thr Ala Gly Gly Cys Thr Thr Thr Thr Thr Thr Ala Thr Thr  
 210 215 220  
 Thr Thr Cys Thr Thr Gly Cys Thr Ala Cys Thr Cys Cys Cys Ala Ala  
 225 230 235 240  
 Gly Gly Ala Cys Cys Thr Gly Gly Thr Gly Ala Thr Ala Thr Thr Thr  
 245 250 255  
 Thr Thr Cys Thr Thr Thr Ala Cys Cys Ala Thr Gly Cys Ala Thr Thr  
 260 265 270  
 Ala Ala Ala Cys Ala Gly Ala Ala Thr Cys Thr Gly Thr Gly Ala Gly  
 275 280 285  
 Thr Cys Thr Thr Thr Thr Cys Thr Gly Gly Ala Ala Ala Ala Ala Ala  
 290 295 300  
 Ala Ala Ala Ala Gly Gly Cys Ala Gly Gly Ala Gly Gly Gly Ala Ala  
 305 310 315 320  
 Cys Ala Thr Ala Cys Thr Ala Gly Thr Thr Ala Ala Ala Ala Ala Gly  
 325 330 335  
 Thr Thr Thr Cys Thr Gly Gly Gly Thr Ala Cys Ala Cys Thr Ala Cys  
 340 345 350  
 Cys Ala Ala Gly Ala Thr Gly Thr Ala Cys Cys Thr Ala Thr Thr Thr  
 355 360 365  
 Ala Thr Thr Gly Ala Thr Ala Thr Ala Cys Ala Ala Ala Thr Gly Gly  
 370 375 380  
 Cys Ala Thr Ala Ala Gly Thr Thr Ala Thr Thr Gly Ala Ala Thr Gly  
 385 390 395 400  
 Cys Thr Thr Gly Cys Thr Ala Thr Ala Gly Gly Cys Ala Thr Thr Cys  
 405 410 415  
 Thr Cys Thr Ala Ala Gly Ala Ala Cys Thr Thr Thr Gly Thr Ala Ala  
 420 425 430  
 Gly Ala Ala Thr Thr Gly Ala Cys Thr Thr Ala Cys Ala Thr Gly Ala  
 435 440 445  
 Gly Cys Thr Ala Cys Thr Thr Cys Ala Thr Ala Gly Cys Ala Gly Thr  
 450 455 460  
 Thr Cys Gly Ala Thr Gly Ala Thr Ala Thr Ala Cys Ala Thr Gly Thr  
 465 470 475 480  
 Thr Gly Thr Thr Ala Thr Thr Ala Thr Cys Ala Cys Cys Ala Cys Thr  
 485 490 495

Thr Thr Ala Cys Ala Gly Ala Thr Ala Ala Gly Gly Ala Ala Ala Thr  
 500 505 510  
 Ala Gly Ala Gly Ala Cys Ala Gly Ala Cys Ala Thr Ala Cys Thr Gly  
 515 520 525  
 Ala Ala Thr Gly Ala Cys Ala Thr Gly Cys Thr Cys Ala Ala Cys Gly  
 530 535 540  
 Cys Cys Ala Cys Thr Cys Cys Ala Cys Thr Ala Gly Cys Ala Ala Gly  
 545 550 555 560  
 Thr Gly Gly Cys Ala Gly Ala Ala Cys Cys Cys Ala Ala Gly Cys  
 565 570 575

<210> 47  
 <211> 549  
 <212> DNA  
 <213> Homo sapiens

<400> 47  
 cgtggcaggt tgcaccaact tttaaacc caagcaatta tgagaggttc atttcccttc 60  
 actcacaaca aaactgcatc ttatcacgtt ttaagaaatc ttgcccattt tgctgggatt 120  
 aacttgtctc tcattattct atatttgc attttaagcta ttaagcttga ataatttcca 180  
 tatgggccat ttgtatttgc aaaacaaaca ctacccttc aatgattttg cctggcttct 240  
 gccttcaact gtatttattg cctctttaaa acactaagtt aatagtttta ttgtttcctt 300  
 tgtcagcctc ctctgaaaat gatgacactt ttcaaacagt accacatctt tctactcatt 360  
 tcagttcaat gctcatatag ggcattgtgc atactgactg tccaagctgt gcaactaagca 420  
 ggtgaatctt ccctgccctt ggatcgagta ataacagaac ttaaagaagg ctctgggta 480  
 ggaaaatcat gggactgggt aacagagaga tatggattca aatcctggct tctctactta 540  
 ctactcaa 549

<210> 48  
 <211> 726  
 <212> DNA  
 <213> Homo sapiens

<400> 48  
 acaccaggaa cacacttggg tgcccatcgg cccaggttt cttctgctgg tctgtctgg 60  
 ccatgtgttt gggtttgaaa gtgtccagat tgccctggttc tctgggatct tcacgcaaaa 120  
 gaaacgagca catgatggtg acctggaact ctctcgtctg gcgtcactgc atttttgcaa 180  
 agcctgtaac tgtcctgtct gcattctggg ctccaaggct cagtcctctg atctttcctg 240  
 acctgtcctt ccctagggt gccttcctct ttttccttat tacagtaaag ttctgtatgt 300

actgcagcat attccattta ttgggaattg aatatatttc ttctatgcca ggatttaaaa 360  
 ttaggattgt taatattgtg gtttgtgctc tggttacaga atttctaagg tttgggtgtt 420  
 ctatcccagc accatatttt ctcaaagcct tgctctctgc agtgggtgat tttgcacaat 480  
 gcaaacttct caggtacttt cttctgagct ccaggtcccc atatccaaca tctactcaac 540  
 atcttattct gagatgttca cctcaaacat gtgagaatca acatgtcaac atgtcaatac 600  
 ctcttgagg atttcctaac tccacagatg gtatccgtcc aattgtgcaa gccaaagagca 660  
 aggcgccagc tgggacgttt cccataccca atctatcatc atgtccatt aaattttacc 720  
 agtagt 726

<210> 49  
 <211> 633  
 <212> DNA  
 <213> Homo sapiens

<400> 49  
 cacttgagag gtgatctcat ctataaagag agtatattta aggagctaat ttacacaggt 60  
 agatgacaaa accttacacc aactcttcaa gcacaggaag caccttacac gaaacacttt 120  
 ggtgaatgct catttctact caattttaaa ctaaattcaa ctgaaaatac ctcattttta 180  
 agaggatatt cagttgactc actggggaaa aacaattgga taaactggtt gtttgccctt 240  
 accgagataa aatcactgat aagttgaatg catttttgta atgtgtcaag acagatggct 300  
 ttaatagtca cacatattaa gtatgggata tgtggcgagg aatagagggtt tttatataca 360  
 gaagtaagaa ctgtagaggg gggaaaccat gaaagataga aaaagatgag aagaaagagg 420  
 acagagatga tacactcaag aaaagctgcc tatttcagag tagaatatat tgtaagcccc 480  
 agtatagttg aatgcttaca agaggagaag cttttagcaa ctctttgtt tcttaatgta 540  
 tatattatat ctgaacagaa tgcttctttc ctgtgagcta gacagtagga agtaatctat 600  
 cttcaacctt gtttgggtgtg taaataaaac ttt 633

<210> 50  
 <211> 618  
 <212> DNA  
 <213> Homo sapiens

<400> 50  
 gtgatacggg tcggctgtgt ccccatccaa atctaattct gaattacagc tcccataatt 60  
 tccatgtgtt gtggaagggg cctgggtggga gataaccgaa tcatgggggc ggctcccccc 120  
 atactgttct cttggtagtg aataagtctc ataagatctg atcgttttat aaggggaaac 180  
 cccttttgct tggttcttat tctctctttg ctggctgcc a tgtaagatgc ttctttgctc 240



ttcctttgtg ttctgccatg actgtggggc ctctccagcc atgtggaaat gtgaggaaat 300  
 taaacctctt tcctttataa attgcccagt cttgggtatg tctttatcag cagtgtgaaa 360  
 tggactaata caccagaagt tcaagactag cttgggcaac acagtaagac cctgtctgta 420  
 taaagaaaga ataagaaaaa gaaaaatatc agtggcataa tattagacag aatactataa 480  
 aactgattaa ggcacattaa tatagccaga ataattattc tgattatatc attttgaaag 540  
 accagataat caatataact gattttttatc ccgagtagct cagacttgca agcatgcact 600  
 accatacttg gataattt 618

<210> 51  
 <211> 1612  
 <212> DNA  
 <213> Homo sapiens

<400> 51  
 cagtgagccg agatggtgcc attgcactct agcctggggc aacagagcca gactccatct 60  
 ccaaaaaaaaa aaggccattc tgaggatcaa ggcaccacta gcaacaggga gcccctatggg 120  
 tctcagaccc tctccccaca tctcctggtc cctgccccca cctggcgtag agggaccagc 180  
 cccacggaag gctcttgagg ccaggtaacc atggggaggg gaggaatggg gacaccttcc 240  
 tcttgagtgt cttagggaag agaagcttag gtcagggtggc tgagggtgga aatgagagag 300  
 gggctctctc ctggaggggc tcaccattcc cttggtcacc caccctaact tcatctcccc 360  
 tgatgtgggg aggagcaggg ggcattggatt cctgagcccc agactcaact gttgtgggtt 420  
 acaggggcat caggagagag agcgagcaga acacactcct gcagcatccc ctggcccccc 480  
 gcccctatgat ggagcccaga gaagctggac agcacgtggg ggccgccaac ggcgcccagg 540  
 aggatgtggc cttcaacctc atcatcctgt ccctcaccga ggggctcggc ctcggtgggc 600  
 tgctggggaa tggggcagtc ctctggctgc tcagctcaa tgtctacaga aacctcttgc 660  
 ccatctacct cctggacgtg gcctgcgcgg atctcatctt ccttggctgc cacatggtgg 720  
 ccatcgtccc cgacttgctg caaggccggc tggacttccc gggcttcgtg cagaccagcc 780  
 tggcaacgct gcgcttcttc tgetacatcg tgggcctgag tctcctggcg gcgctcagcg 840  
 tggagcagtg cctggccgcc ctcttcccag cctggtactc gtgcgcgcgc ccacgccacc 900  
 tgaccacctg tgtgtgcgcc ctacactggg ccctctgcct gctgctgcac ctgctgctca 960  
 gcggcgctg caccagttc ttcggggagc ccagccgcca cttgtgccgg acgctgtggc 1020  
 tgggtggcagc ggtgctgctg gctctgctgt gttgcacat gtgtggggcc agccttatgc 1080  
 tgctgctgcg ggtggagcga ggccccagc ggccccacc ccggggcttc cctgggctca 1140

tcctcctcac	cgctcctcctc	ttcctctttct	gcggcctgcc	cttcggcatc	tactggctgt	1200
cccggaacct	gctctggtac	atccccact	acttctacca	cttcagcttc	ctcatggccg	1260
ccgtgcaactg	cgcggccaaag	cccgtcgtct	acttctgcct	gggcagtgcc	cagggcccgca	1320
ggctgcccct	ccggctggtc	ctccagcgag	cgctgggaga	cgaggctgag	ctggggggccg	1380
tcaggagagac	ctcccgccgg	ggcctggtgg	acatagcagc	ctgagccctg	gggccccoga	1440
ccccagctgc	agccccctg	aggcaagagg	gtgacgtggg	gaagggtggtg	gggtcagagg	1500
ctggggccag	ccggacctgg	aggaggcctt	ggtgggtgac	ccggtcatgt	gctgtcaaag	1560
ttgtgaccct	tggtctggag	catgaggctc	ccctgggagg	cagctggaaa	gg	1612

&lt;210&gt; 52

&lt;211&gt; 1527

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

atgacgtcca	cctgcaccaa	cagcacgcgc	gagagtaaca	gcagccacac	gtgcatgccc	60
ctctccaaaa	tgcccatcag	cctggcccac	ggcatcatcc	gctcaaccgt	gctggttatc	120
ttcctcgccg	cctctttcgt	cggcaacata	gtgctggcgc	tagtgttgca	gcgcaagccg	180
cagctgctgc	aggtgaccaa	ccgttttata	tttaacctcc	tcgtcacoga	cctgctgcag	240
atttcgctcg	tggccccctg	ggtgggtggc	acctctgtgc	ctctcttctg	gcccccaac	300
agccacttct	gcacggccct	ggttagcctc	acctacctgt	tcgccttcgc	cagcgtcaac	360
accattgtcg	tggtgtcagt	ggatcgctac	ttgtccatca	tccacctct	ctcctacccg	420
tccaagatga	cccagcgccg	cggttacctg	ctcctctatg	gcacctggat	tgtggccatc	480
ctgcagagca	ctcctccact	ctacggctgg	ggccaggctg	cctttgatga	gcgcaatgct	540
ctctgctcca	tgatctgggg	ggccagcccc	agctacacta	ttctcagcgt	ggtgtccttc	600
atcgtcattc	cactgattgt	catgattgcc	tgctactccg	tggtgttctg	tgcagcccg	660
aggcagcatg	ctctgctgta	caatgtcaag	agacacagct	tggaagtgcg	agtcaaggac	720
tgtgtggaga	atgaggatga	agaggagca	gagaagaagg	aggagtcca	ggatgagagt	780
gagtttcgcc	gccagcatga	aggtgaggtc	aaggccaagg	agggcagaat	ggaagccaag	840
gacggcagcc	tgaaggccaa	ggaaggaagc	acggggacca	gtgagagtag	tgtagaggcc	900
aggggcagcg	aggaggtcag	agagagcagc	acggtggcca	gcgacggcag	catggagggg	960
aaggaaggca	gcaccaaagt	tgaggagaac	agcatgaagg	cagacaagg	tcgcacagag	1020
gtcaaccagt	gcagcattga	cttgggtgaa	gatgacatgg	agtttggtga	agacgacatc	1080

aatttcagtg aggatgacgt cgaggcagtg aacatcccgg agagcctccc acccagtcgt 1140  
cgtaacagca acagcaaccc tcctctgccc aggtgctacc agtgcaaagc tgctaaagtg 1200  
atcttcatca tcattttctc ctatgtgcta tccctggggc cctactgctt tttagcagtc 1260  
ctggccgtgt ggggtggatgt cgaaaccag gtacccagtg gggatgatcac cataatcatc 1320  
tggtttttct tcctgcagtg ctgcatccac ccctatgtct atggctacat gcacaagacc 1380  
attaagaagg aaatccagga catgctgaag aagttcttct gcaaggaaaa gccccgaaa 1440  
gaagatagcc acccagacct gcccggaaca gaggggtggga ctgaaggcaa gattgtccct 1500  
tcctacgatt ctgctacttt tccttga 1527

<210> 53  
<211> 939  
<212> DNA  
<213> Homo sapiens

<400> 53  
atgatggagc ccagagaagc tggacagcac gtggggggccg ccaacagcgc ccaggaggat 60  
gtggccttca acctcatcat cctgtccctc accgaggggc tcggcctcgg tgggctgctg 120  
gggaatgggg cagtcctctg gctgctcagc tccaatgtct acagaaaccc cttcgccatc 180  
tacctcctgg acgtggcctg cgcggatctc atcttccttg gctgccacat ggtggccatc 240  
gtccccgact tgctgcaagg ccggctggac ttcccgggct tcgtgcagac cagcctggca 300  
acgctgcgct tcttctgcta catcgtgggc ctgagtctcc tggcggccgt cagcgtggag 360  
cagtgcctgg ccgccctctt cccagcctgg tactcgtgcc gccgcccacg ccacctgacc 420  
acctgtgtgt gcgccctcac ctgggcccctc tgccctgtgc tgcaacctgct gctcagcggc 480  
gcctgcaccc agttcttcgg ggagcccagc cgccacttgt gccggacgct gtggctggtg 540  
gcagcgggtg tgctggctct gctgtgttgc accatgtgtg gggccagcct tatgctgctg 600  
ctgcgggtgg agcgaggccc ccagcggccc ccaccccggg gcttcctcgg gctcatcctc 660  
ctcaccgtcc tcctcttctt cttctgcggc ctgcccttcg gcattactg gctgtcccgg 720  
aacctgctct ggtacatccc ccactacttc taccatttca gcttcctcat ggccgccgtg 780  
cactgcgcgg ccaagcccggt cgtctacttc tgccctgggca gtgcccaggg ccgcaggctg 840  
cccctccggc tggtcctcca gcgagcgtg ggagacgagg ctgagctggg ggccgtcagg 900  
gagacctccc gccggggcct ggtggacata gcagcctga 939

<210> 54  
<211> 1778

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 54

```

ggagcctcgc cggccgccat gtggagctgc agctggttca acggcacagg gctgggtggag      60
gagctgcctg cctgccagga cctgcagctg gggctgtcac tgttgtoct gctgggcctg      120
gtgggtgggcg tgccagtggg cctgtgctac aacgccctgc tgggtgctggc caacctacac      180
agcaaggcca gcatgaccat gccggacgtg tactttgtca acatggcagt ggcaggcctg      240
gtgctcagcg ccctggcccc tgtgcacctg ctcggecccc cgagctcccg gtgggcgctg      300
tggagtgtgg gcggcgaagt ccacgtggca ctgcagatcc ccttcaatgt gtctcactg      360
gtggccatgt actccaccgc cctgctgagc ctgcaccact acatcgagcg tgcactgccg      420
cggacctaca tggccagcgt gtacaacacg cggcacgtgt gcggcttcgt gtgggggtggc      480
gcgctgctga ccagcttctc ctcgctgctc ttctacatct gcagccatgt gtccacccgc      540
gcgctagagt gcgccaagat gcagaacgca gaagctgccg acgccacgct ggtgttcac      600
ggctacgtgg tgccagcact ggccaccctc tacgcgctgg tgetactctc ccgcgtccgc      660
agggaggaca cgccctgga ccgggacacg ggccggctgg agccctcggc acacaggctg      720
ctgggtggcca ccgtgtgcac gcagtttggg ctctggacgc cacactatct gatcctgctg      780
gggcacacgg tcatcatctc gcgaggggaag cccgtggacg cacactacct ggggctactg      840
cactttgtga aggatttctc caaactcctg gccttctcca gcagctttgt gacaccactt      900
ctctaccgct acatgaacca gagcttcccc agcaagctcc aacggctgat gaaaaagctg      960
ccctgcgggg accggcactg ctccccggac cacatggggg tgcagcagggt gctggcgtag     1020
gcggcccagc cctcctgggg agacgtgact ctgggtggacg cagagcactt agttaccctg     1080
gacgctcccc acatccttcc agaaggagac gagctgctgg aagagaagca ggaggggtgt     1140
ttttcttgaa gtttcctttt tcccacaaat gccactcttg ggccaaggct gtgggtccccg     1200
tggctggcat ctggcttgag tctccccgag gcctgtgcgt ctcccaaaca cgcagctcaa     1260
ggtccacatc cgcaaaagcc tcctcgctt cagctcctc agcattcagt ttgtcaatga     1320
agtgatgaaa gcttagagcc agtatattata ctttgtgggt aaaatacttg attccccctt     1380
gtttgtttta caaaaacaga tgtttcctag aaaaatgaca aatagtaaaa tgaacaaaac     1440
cctacgaaag aatggcaaca gccagggtgg ccgggccctg ccagtgggcg gcgtgtgcta     1500
gcaaggcctg ccgggtgtgc cgcagtcacc acagggttct gagaacattt cacagaagtg     1560
cctgagacgc ggagacatgg ctgggtgtta atggagctat tcaatagcag tgacgcgctc     1620
tcctcagcca ccaaagtcc ctgacaccct cccagcccc cacagataac atcagctgag     1680

```

gttttttttca gtatgaacct gtcctaaatc aattcctcaa agtgtgcaca aaactaaaga 1740  
 atataaataa acaaaagaaa ggcaaaaaaa aaaaaaaa 1778

<210> 55  
 <211> 1014  
 <212> DNA  
 <213> Homo sapiens

<400> 55  
 atggggaacg attctgtcag ctacgagtat ggggattaca gcgacctctc ggaccgccct 60  
 gtggactgcc tggatggcgc ctgcctggcc atcgaccgcg tgcgcgtggc cccgctccca 120  
 ctgtatgccg ccatcttctt ggtgggggtg ccgggcaatg ccatggtggc ctgggtggct 180  
 gggaagggtg cccgccggag ggtgggtgcc acctggttgc tccacctggc cgtggcggat 240  
 ttgctgtgct gtttgtctct gcccatcctg gcagtgccca ttgccgtgg aggccactgg 300  
 ccgtatggtg cagtgggctg tggggcgtg cctccatca tctgctgac catgtatgcc 360  
 agcgtcctgc tctggcagc tctcagtgcc gacctctgt tctggctct cgggcctgcc 420  
 tgggtggtcta cggttcagcg ggcgtgcggg gtgcagggtg cctgtggggc agcctggaca 480  
 ctggccttgc tgctcacctg gccctccgcc atctaccgcc ggctgcacca ggagcacttc 540  
 ccagcccggc tgcagtgtgt ggtggactac ggcggctcct ccagcaccga gaatgcggtg 600  
 actgccatcc ggtttctttt tggttctctg gggccctgg tggccgtggc cagctgccac 660  
 agtgccctcc tgtgctgggc agcccgacgc tgccggccgc tgggcacagc cattgtggtg 720  
 gggttttttg tctgctgggc acctaccac ctgctggggc tgggtgctcac tgtggcggcc 780  
 ccgaactccg cactcctggc cagggccctg cgggctgaac cctcatcgt gggccttgcc 840  
 ctgctcaca gctgcctcaa tccatgctc ttctgtatt ttgggagggc tcaactccgc 900  
 cggtcactgc cagctgcctg tcaactgggc ctgagggagt ccagggcca ggacgaaagt 960  
 gtggacagca agaaatccac cagccatgac ctggtctcgg agatggaggt gtag 1014

<210> 56  
 <211> 1170  
 <212> DNA  
 <213> Homo sapiens

<400> 56  
 atggcacctt ctcatcgggc atcacagggtg gggttttgcc ccaccctga acgccctctg 60  
 tggcgccttc caccacctg tagggccaga aggatgtcgg tctgctaccg tccccaggg 120  
 aacgagacac tgctgagctg gaagacttcg cgggccacag gcacagcctt cctgctgctg 180

gcggcgctgc tggggctgcc tggcaacggc ttcgtggtgt ggagcttggc gggctggcgg 240  
 cctgcacggg ggcgaccgct ggcggccacg cttgtgctgc acctggcgct ggccgacggc 300  
 gcgggtgctgc tgetcacgcc gctctttgtg gccttctga cccggcaggg ctggccgctg 360  
 ggccaggcgg gctgcaaggc ggtgtactac gtgtgcgcgc tcagcatgta cggcagcgtg 420  
 ctgctcaccg gcctgctcag cctgcagcgc tgcctcgag tcacccgccc ctctctggcg 480  
 cctcggtctgc gcagcccggc cctggcccgc cgcctgctgc tggcggtctg gctggccgcc 540  
 ctgttgctcg ccgtcccggc cgcctctac cgcacctgt ggagggaccg cgtatgccag 600  
 ctgtgccacc cgtgcgggt ccacgccgc gccacctga gcctggagac tctgaccgct 660  
 ttcgtgcttc ctttcgggct gatgctcggc tgctacagcg tgacgctggc acggctgcgg 720  
 ggcgcccgt ggggctccgg gcggcacggg gcgcgggtgg gccggctggt gagcgccatc 780  
 gtgcttgcc tggcttgct ctgggcccc taccacgcag tcaaccttct gcaggcggtc 840  
 gcagcgctgg ctccaccgga aggggccttg gcgaagctgg gcggagccgg ccaggcggcg 900  
 cgagcgggaa ctacggcctt ggccttcttc agttctagcg tcaaccgggt gctctacgtc 960  
 ttcacgctg gagatctgct gccccgggca ggtccccgtt tcctcacgcg gctcttcgaa 1020  
 ggctctgggg aggcccgagg gggcgccgc tctagggaa ggacatgga gctccgaact 1080  
 acccctcagc tgaaagtgtt ggggcagggc cgcggcaatg gagaccggg ggggtgggatg 1140  
 gagaaggacg gtccggaatg ggacctttga 1170

<210> 57  
 <211> 1965  
 <212> DNA  
 <213> Homo sapiens

<400> 57  
 ccgcccgcgc gggaggacgc gagcaccag ctttaatccc tggaaagtcc acgaacaatg 60  
 aatccatttc atgcatcttg ttggaacacc tctgccgaac ttttaaaaca atcctggaat 120  
 aaagagtttg cttatcaaac tgccagtgtg gtagatacag tcatcctccc ttccatgatt 180  
 gggattatct gttcaacagg gctggttggc aacatcctca ttgtattcac tataataaga 240  
 tccaggaaaa aaacagtccc tgacatctat atctgcaacc tggctgtggc tgatttggtc 300  
 cacatagttg gaatgccttt tcttattcac caatgggccc gagggggaga gtgggtgttt 360  
 ggggggcctc tctgcacat catcacatcc ctggatactt gtaaccaatt tgctgtagt 420  
 gccatcatga ctgtaatgag tgtggacagg tactttgccc tcgtccaacc atttcgactg 480  
 acacgttgga gaacaaggta caagaccatc cggatcaatt tgggcctttg ggcagcttcc 540

```

tttatcctgg cattgcctgt ctgggtctac tcgaaggcca tcaaatttaa agacgggtgtt    600
gagagttgtg cttttgattt gacatcccct gacgatgtac tctggtatac actttatttg    660
acgataacaa cttttttttt ccctctaccc ttgatttttg tgtgctatat tttaatttta    720
tgctataactt gggagatgta tcaacagaat aaggatgcca gatgctgcaa tcccagtgtg    780
ccaaaacaga gagtgatgaa gttgacaaag atgggtgctgg tgctgggtgg agtctttatc    840
ctgagtgtctg ccccttatca tgtgatacaa ctgggtgaact tacagatgga acagcccaca    900
ctggcctttct atgtgggtta ttacctctcc atctgtctca gctatgccag cagcagcatt    960
aacccttttc tctacatcct gctgagtgga aatttccaga aacgtctgcc tcaaattcaa   1020
agaagagcga ctgagaagga aatcaacaat atgggaaaca ctctgaaatc acacttttag   1080
gaaagtacat ggatcaccat gagtctagac atgattgtct atcttactgg tattattaga   1140
aagggcaggt gtaccgatat gtttatgccc attcttcttg tgtacttggt actcttagca   1200
gcatggaaga gaagtgtaac catgcaaata caatgagctt aatatgctaa ctttagcaag   1260
atgtaaaatg ttgatctata ttgtgggtag ggaatgggat agtctgagat acccaggctt   1320
catgatgggtg tatattattt cagcatatta taaactagtc actaatgaaa atggccatcc   1380
atgaccattg actcaaaact caccaaggaa cctgaccttg cctccacac tgtggcctca   1440
ctgtaacagt ttcctcaagg ttcctaggag ggtatcacct tagagtgaag tctaaaattt   1500
ggctattttt tatctataaa aaatgtcagt tttatatggt ccaatactaa taccctcaac   1560
aactaagccc caccttttag aataagttac cattttattgc acacatgcaa tgtgtaagat   1620
tacatgtaac aaacctgtga aataagtatt attacctttg tttgctaagg ctcagaaagg   1680
agaaatgata ggcctaattg tgcaacagct atctaagagc tgagctaaca ttcagctctg   1740
cctgtttctt ttctactgcc gaccttgaca acctttactt atcatactgg agaaccctag   1800
aacttgaggt ttcttttgct ttctcctgta gccctacaag aggagaacta aagtctgata   1860
gaaatgagtt gatgttttaa gcatcatttt ggattatctt gttctcacac ctgctaactg   1920
tagaaactgg catctggact ttaataataa tactttactt ctgga                      1965

```

```

<210> 58
<211> 1031
<212> DNA
<213> Homo sapiens

```

```

<400> 58
ccatgaccag caatttttcc caacctgttg tgcagctttg ctatgaggat gtgaatggat    60
cttgatttga aactccctat tctcctgggt cccgggtaat tctgtacacg gcgttttagct   120

```

```

ttgggtcttt gctggctgta tttggaaatc tcttagtaat gacttctgtt cttcatttta 180
agcagctgca ctctccaacc aattttctca ttgcctctct ggcoctgtgct gacttcttgg 240
taggtgtgac tgtgatgctt ttcagcatgg tcaggacggg ggagagctgc tggatatttg 300
gagccaaatt ttgtactctt cacagttgct gtgatgtggc attttgttac tcttctgtcc 360
tccacttgtg cttcatctgc atcgacaggt acattgtggt tactgatccc ctgggtctatg 420
ctaccaagtt caccgtgtct gtgtcgggaa tttgcatcag cgtgtcctgg attctgcctc 480
tcacgtacag cggtgctgtg ttctacacag gtgtcaatga tgatgggctg gaggaattag 540
taagtgtctt caactgcgta ggtggctgtc aaattattgt aagtcaaggc tgggtgttga 600
tagattttct gttattcttc atacctacc ttgttatgat aattctttac agtaagattt 660
ttcttatagc taaacaacaa gctataaaaa ttgaaactac tagtagcaaa gtagaatcat 720
cctcagagag ttataaaatc agagtggcca agagagagag gaaagcagct aaaaccctgg 780
gggtcacggg actagcattt gttatttcat ggttacogta tacagttgat atattaattg 840
atgcctttat gggcttcctg acccctgcct atatctatga aatttgctgt tggagtgcct 900
attataactc agccatgaat cctttgattt atgctctatt ttatccttgg tttaggaaag 960
ccataaaact tattttaagt ggagatgttt taaaggctag ttcacatcaacc attagtttat 1020
ttttagaata a 1031

```

```

<210> 59
<211> 972
<212> DNA
<213> Homo sapiens

```

```

<400> 59
atgccactcc ctgtgcccc agcgggggca caaaagactc ctgaggacca tgtctgtctc 60
catcttgacg gtccctcgcc agcaccctct gagccggcca ggatgtttgg gctgttcggc 120
ctctggagaa ccttcgacag tgtggtcttc tacctgacgc tgatcgtggg cctcggggga 180
ccggtaggta acgggctggt gctctggaac ctcggttcc gcatcaagaa gggcccttc 240
tccatctacc tgctgcacct ggccgcgcc gacttctgt tctctctctg ccgtgtgggc 300
ttctccgtgg ctgaggtgc cctgggcgcc caggacacac tctacttctg gctcaccttc 360
ctgtggttcg cggtggggct ctggctgctg gcggccttca gcgtggagcg ctgcctctcc 420
gacctcttc ccgctgcta ccagggtgc cggccagac acgcctcggc cgtcctctgc 480
gccctggtgt ggaccccgac cctgcgggcc gtgcgctgc ccgccaacgc ctgcggcctg 540
ctgcgcaaca gcgcgtgccc cctggtctgc ccgcgtacc acgtggccag cgtcacctgg 600

```



```

ttcctggtgc tggcccgcg cgcctggacg gctggcggtg tctcttttgt ctgggtgacc 660
tgctgctcca ctgccccg cgccaggctc tacggcatcg tctggggcgc gctgctcctg 720
ctctttcttct gtggcctgcc ctcggtcttc tactggagcc tgcagcccct gctgaacttc 780
ctgctgcccc tgttttcccc gctggccacg ctgctggcct gcgtcaacag cagctccaag 840
ccctcatct actcggggtt gggccgacag cccgggaagc gggagccgct gaggtcggtg 900
ctgcggaggg ccctggggga gggcgccgag ctgggtgccg ggggacagtc cctgcccatg 960
ggtctctat aa 972

```

```

<210> 60
<211> 1011
<212> DNA
<213> Homo sapiens

```

```

<400> 60
atgaacaaca atacaacatg tattcaacca tctatgatct cttccatggc tttaccaatc 60
atttacatcc tcctttgtat tgttggtgtt tttggaaaca ctctctctca atggatattt 120
ttaacaaaaa taggtaaaaa aacatcaacg cacatctacc tgtcacacct tgtgactgca 180
aacttacttg tgtgcagtgc catgcctttc atgagtatct atttctgaa aggtttccaa 240
tggaatatc aatctgctca atgcagagtg gtcaattttc tgggaactct atccatgcat 300
gcaagtatgt ttgtcagtct ctttaatttta agttggattg ccataagccg ctatgctacc 360
ttaatgcaaa aggattcctc gcaagagact acttcatgct atgagaaaat attttatggc 420
catttactga aaaaatttcg ccagcccaac tttgctagaa aactatgcat ttacatatgg 480
ggagttgtac tgggcataat cattccagtt accgtatact actcagtcac agaggctaca 540
gaaggagaag agagcctatg ctacaatcgg cagatggaac taggagccat gatctctcag 600
attgcaggtc tcattggaac cacatttatt ggattttcct ttttagtagt actaacatca 660
tactactctt ttgtaagcca tctgagaaaa ataagaacct gtacgtccat tatggagaaa 720
gatttgactt acagttctgt gaaaagacat cttttggtca tccagattct actaatagtt 780
tgcttccttc cttatagtat ttttaaacc cttttttatg ttctacacca aagagataac 840
tgtcagcaat tgaattattt aatagaaaca aaaaacattc tcacctgtct tgcttcggcc 900
agaagtagca cagaccccat tatatttctt ttattagaca aaacattcaa gaagacacta 960
tataatctct ttacaaagtc taattcagca catatgcaat catatggttg a 1011

```

```

<210> 61
<211> 180
<212> PRT

```

&lt;213&gt; Homo sapiens

&lt;400&gt; 61

Tyr Ala Thr Ile Gly Arg Trp Glu Leu Gly Ala Met Ile Ser Gln Ile  
 1 5 10 15  
 Ala Gly Leu Ile Gly Thr Thr Phe Ile Gly Phe Ser Phe Leu Val Val  
 20 25 30  
 Leu Thr Ser Tyr Tyr Ser Phe Val Ser His Leu Arg Lys Ile Arg Thr  
 35 40 45  
 Cys Thr Ser Ile Met Glu Lys Asp Leu Thr Tyr Ser Ser Val Lys Arg  
 50 55 60  
 His Leu Leu Val Ile Gln Ile Leu Leu Ile Val Cys Phe Leu Pro Tyr  
 65 70 75 80  
 Ser Ile Phe Lys Pro Ile Phe Tyr Val Leu His Gln Arg Asp Asn Cys  
 85 90 95  
 Gln Gln Leu Asn Tyr Leu Ile Glu Thr Lys Asn Ile Leu Thr Cys Leu  
 100 105 110  
 Ala Ser Ala Arg Ser Ser Thr Asp Pro Ile Ile Phe Leu Leu Leu Asp  
 115 120 125  
 Lys Thr Phe Lys Lys Thr Leu Tyr Asn Leu Phe Thr Lys Ser Asn Ser  
 130 135 140  
 Ala His Met Gln Ser Tyr Gly Leu Leu Asn Gly Lys Pro His Asn Ile  
 145 150 155 160  
 Lys Lys Ser Ile His Val Thr Leu Leu Gly Thr Leu Asn Tyr Ile Ile  
 165 170 175  
 Asn Met Ser Gln  
 180

&lt;210&gt; 62

&lt;211&gt; 530

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 62

Val Ser Arg Asp Gly Ala Ile Ala Leu Pro Gly Ala Thr Glu Pro Asp  
 1 5 10 15  
 Ser Ile Ser Lys Lys Lys Arg Pro Phe Gly Ser Arg His His Gln Gln  
 20 25 30  
 Gly Ala Pro Trp Val Ser Asp Pro Leu Pro Thr Ser Pro Gly Pro Cys  
 35 40 45  
 Pro His Leu Ala Tyr Arg Asp Gln Pro His Gly Arg Leu Leu Arg Pro  
 50 55 60

Gly Asn His Gly Glu Gly Arg Asn Gly Asp Thr Phe Leu Leu Ser Val  
 65 70 75 80  
 Leu Gly Lys Arg Ser Leu Gly Gln Val Ala Glu Gly Gly Asn Glu Arg  
 85 90 95  
 Gly Val Ser Ser Trp Arg Val Ser Pro Phe Pro Trp Ser Pro Thr Gln  
 100 105 110  
 Leu Ser Ser Pro Leu Met Trp Gly Gly Ala Gly Gly Met Asp Ser Ala  
 115 120 125  
 Pro Asp Ser Thr Val Val Val Tyr Arg Gly Ile Arg Arg Glu Ser Glu  
 130 135 140  
 Gln Asn Thr Leu Leu Gln His Pro Leu Ala Pro Arg Pro Met Met Glu  
 145 150 155 160  
 Pro Arg Glu Ala Gly Gln His Val Gly Ala Ala Asn Gly Ala Gln Glu  
 165 170 175  
 Asp Val Ala Phe Asn Leu Ile Ile Leu Ser Leu Thr Glu Gly Leu Gly  
 180 185 190  
 Leu Gly Gly Leu Leu Gly Asn Gly Ala Val Leu Trp Leu Leu Ser Ser  
 195 200 205  
 Asn Val Tyr Arg Asn Pro Phe Ala Ile Tyr Leu Leu Asp Val Ala Cys  
 210 215 220  
 Ala Asp Leu Ile Phe Leu Gly Cys His Met Val Ala Ile Val Pro Asp  
 225 230 235 240  
 Leu Leu Gln Gly Arg Leu Asp Phe Pro Gly Phe Val Gln Thr Ser Leu  
 245 250 255  
 Ala Thr Leu Arg Phe Cys Tyr Ile Val Gly Leu Ser Leu Leu Ala Ala  
 260 265 270  
 Val Ser Val Glu Gln Cys Leu Ala Ala Leu Phe Pro Ala Trp Tyr Ser  
 275 280 285  
 Cys Arg Arg Pro Arg His Leu Thr Thr Cys Val Cys Ala Leu Thr Trp  
 290 295 300  
 Ala Leu Cys Leu Leu Leu His Leu Leu Leu Ser Gly Ala Cys Thr Gln  
 305 310 315 320  
 Phe Phe Gly Glu Pro Ser Arg His Leu Cys Arg Thr Leu Trp Leu Val  
 325 330 335  
 Ala Ala Val Leu Leu Ala Leu Leu Cys Cys Thr Met Cys Gly Ala Ser  
 340 345 350  
 Leu Met Leu Leu Leu Arg Val Glu Arg Gly Pro Gln Arg Pro Pro Pro  
 355 360 365  
 Arg Gly Phe Pro Gly Leu Ile Leu Leu Thr Val Leu Leu Phe Leu Phe  
 370 375 380

Cys Gly Leu Pro Phe Gly Ile Tyr Trp Leu Ser Arg Asn Leu Leu Trp  
 385 390 395 400  
 Tyr Ile Pro His Tyr Phe Tyr His Phe Ser Phe Leu Met Ala Ala Val  
 405 410 415  
 His Cys Ala Ala Lys Pro Val Val Tyr Phe Cys Leu Gly Ser Ala Gln  
 420 425 430  
 Gly Arg Arg Leu Pro Leu Arg Leu Val Leu Gln Arg Ala Leu Gly Asp  
 435 440 445  
 Glu Ala Glu Leu Gly Ala Val Arg Glu Thr Ser Arg Arg Gly Leu Val  
 450 455 460  
 Asp Ile Ala Ala Ala Leu Gly Pro Pro Thr Pro Ala Ala Ala Pro Val  
 465 470 475 480  
 Arg Gln Glu Gly Asp Val Gly Lys Val Val Gly Ser Glu Ala Gly Ala  
 485 490 495  
 Ser Arg Thr Trp Arg Arg Pro Trp Trp Val Thr Arg Ser Cys Ala Val  
 500 505 510  
 Lys Val Val Thr Leu Gly Leu Glu His Glu Ala Pro Leu Gly Gly Ser  
 515 520 525  
 Trp Lys  
 530

<210> 63  
 <211> 313  
 <212> PRT  
 <213> Homo sapiens

<400> 63

Gly Trp Gln His Arg Leu Ser Arg Leu Cys Glu Lys Asn Gly Gly Thr  
 1 5 10 15  
 Thr Ser His Cys Cys Val Arg Arg Met Ala Ala Thr Pro Leu Met Val  
 20 25 30  
 Val Gly Glu Trp Arg His His Leu Ser Trp Leu Cys Glu Ala Leu Arg  
 35 40 45  
 Ala Pro Val Ser Leu Leu Arg Gly Glu Trp Arg His His Leu Ser Trp  
 50 55 60  
 Leu Cys Asn Ile Ser Ile Thr Leu Lys Tyr Pro Leu Ile Val Gly Ala  
 65 70 75 80  
 Ile Cys Asn Ala Glu Val Val Pro Gln Lys His Phe Asn His Phe His  
 85 90 95  
 Phe His Gly Ser Ala Lys Thr Tyr Ala Gly Pro His Arg Ser Gln Arg  
 100 105 110

Ser His Leu Cys Phe Arg Ala Lys Pro Val Phe Leu Leu Ser Thr Ala  
 115 120 125

Asn Ile Leu Thr Val Ile Ile Leu Ser Gln Leu Val Ala Arg Arg Gln  
 130 135 140

Lys Ser Ser Tyr Asn Tyr Leu Leu Ala Leu Ala Ala Ala Asp Ile Leu  
 145 150 155 160

Val Leu Phe Phe Ile Val Phe Val Asp Phe Leu Leu Glu Asp Phe Ile  
 165 170 175

Leu Asn Met Gln Met Pro Gln Val Pro Asp Lys Ile Ile Glu Val Leu  
 180 185 190

Glu Phe Ser Ser Ile His Thr Ser Ile Trp Ile Thr Val Pro Leu Thr  
 195 200 205

Ile Asp Arg Tyr Ile Ala Val Cys His Pro Leu Lys Tyr His Thr Val  
 210 215 220

Ser Tyr Pro Ala Arg Thr Arg Lys Val Ile Val Ser Val Tyr Ile Thr  
 225 230 235 240

Cys Phe Leu Thr Ser Ile Pro Tyr Tyr Trp Trp Pro Asn Ile Trp Thr  
 245 250 255

Glu Asp Tyr Ile Ser Thr Ser Val His His Val Leu Ile Trp Ile His  
 260 265 270

Cys Phe Thr Val Tyr Leu Val Pro Cys Ser Ile Phe Phe Ile Leu Asn  
 275 280 285

Ser Ile Ile Val Tyr Lys Leu Arg Arg Lys Ser Asn Phe Asp Asn Gly  
 290 295 300

Ser Ile Ala Ser His Gly Cys Val Arg  
 305 310

<210> 64

<211> 173

<212> PRT

<213> Homo sapiens

<400> 64

His Leu Cys Arg Trp Thr Ser Tyr Ala Glu Asp Pro Leu Gln Gly Ser  
 1 5 10 15

Glu Asn Ala Cys Asn Gly Ser Arg Gln Pro Ser Leu Pro Ser Thr Ala  
 20 25 30

Ser Pro Gly Thr Asn Glu Arg Pro Arg Arg Leu Thr Ser Trp Cys His  
 35 40 45

Leu Ser Cys Ala Gly Leu Pro Thr Leu Leu Val Ile Ile Gly Ala Ala  
 50 55 60

Cys His Gly Ser His Val Gln His Phe Leu Tyr Glu Thr Arg Phe Trp  
 43

65                      70                      75                      80  
 Leu His Trp Leu Asn Ser Ala Ile Ser Pro Leu Pro Tyr Ser Phe Cys  
                                  85                      90                      95  
 His Glu Phe Gln Lys Ser Ile Pro Ser Asn Arg Cys Val Leu Glu Arg  
                                  100                      105                      110  
 Leu Ser Glu Ile His Pro Pro Pro Tyr Val His Val Thr Glu Thr Ala  
                                  115                      120                      125  
 Leu Val Phe Leu Met Val Glu Ala Val Gln Val Leu Gly Val Leu Asn  
                                  130                      135                      140  
 Lys Glu Leu Asp Lys Thr His Lys Gln Ser Lys Glu Gly Met Lys Phe  
 145                                   150                      155                      160  
 Ile Glu Asn Glu Ser Thr Leu His Ser Val Gly Ala Gly  
                                  165                      170

<210> 65  
 <211> 187  
 <212> PRT  
 <213> Homo sapiens

<400> 65

Tyr Phe Tyr Pro Lys Leu Leu Cys Ser Leu Leu Glu Phe Pro Arg Lys  
 1                      5                      10                      15  
 Ile Pro Lys Glu Leu Met Val Ser Tyr Leu Leu Leu Tyr Tyr Glu Glu  
                                  20                      25                      30  
 Glu Lys Cys Phe Thr Asn Ile Phe Ile Ile Gly Ala Phe Phe Val Val  
                                  35                      40                      45  
 Ser Glu Ile Ile Lys Thr Arg Ile Asp Phe Tyr Ser Val Tyr Phe Cys  
                                  50                      55                      60  
 Tyr Asn Leu Tyr Pro Phe Leu Leu Ile Ser Val Phe Trp Met Pro Asn  
 65                                   70                      75                      80  
 Leu Glu Tyr Ile Thr Lys Val Thr Phe Ser Phe Ser Leu Ser Ile Gln  
                                  85                      90                      95  
 Asp Asp Leu Lys His Leu Trp Leu Pro Phe Leu Ile Phe Leu Leu Cys  
                                  100                      105                      110  
 Lys Phe Ile Lys Gly Gln Ser Leu Cys Ala Leu Ile Ile Pro Ala Phe  
                                  115                      120                      125  
 Ser Cys Phe Thr Cys Ser Thr Ile Tyr Phe Val Phe Ile Met Phe Ser  
                                  130                      135                      140  
 Phe Thr Leu Cys Thr Ile Ile Asp Tyr Asn Glu Glu Asn Leu Asp Asn  
 145                                   150                      155                      160  
 Leu Leu Leu Lys Arg Phe Phe Arg Gln Ile Ile Gly Phe Phe Cys Ile  
                                  165                      170                      175

Leu Lys Arg Tyr Val Tyr Phe Ala Thr Arg Val  
 180 185

<210> 66  
 <211> 287  
 <212> PRT  
 <213> Homo sapiens

<400> 66

Phe Leu Leu Ser Thr Ala Asn Ile Leu Thr Val Ile Ile Leu Ser Gln  
 1 5 10 15  
 Leu Val Ala Arg Arg Gln Lys Ser Ser Tyr Asn Tyr Leu Leu Ala Leu  
 20 25 30  
 Ala Ala Ala Asp Ile Leu Val Leu Phe Phe Ile Val Phe Val Asp Phe  
 35 40 45  
 Leu Leu Glu Asp Phe Ile Leu Asn Met Gln Met Pro Gln Val Pro Asp  
 50 55 60  
 Lys Ile Ile Glu Val Leu Glu Phe Ser Ser Ile His Thr Ser Ile Trp  
 65 70 75 80  
 Ile Thr Val Pro Leu Thr Ile Asp Arg Tyr Ile Ala Val Cys His Pro  
 85 90 95  
 Leu Lys Tyr His Thr Val Ser Tyr Pro Ala Arg Thr Arg Lys Val Ile  
 100 105 110  
 Val Ser Val Tyr Ile Thr Cys Phe Leu Thr Ser Ile Pro Tyr Tyr Trp  
 115 120 125  
 Trp Pro Asn Ile Trp Thr Glu Asp Tyr Ile Ser Thr Ser Val His His  
 130 135 140  
 Val Leu Ile Trp Ile His Cys Phe Thr Val Tyr Leu Val Pro Cys Ser  
 145 150 155 160  
 Met Phe Phe Ile Leu Asn Ser Ile Ile Val Tyr Lys Leu Arg Arg Lys  
 165 170 175  
 Ser Asn Phe Arg Leu Arg Gly Tyr Ser Thr Gly Lys Thr Thr Ala Ile  
 180 185 190  
 Leu Phe Thr Ile Thr Ser Ile Phe Ala Thr Leu Trp Ala Pro Arg Ile  
 195 200 205  
 Ile Met Ile Leu Tyr His Leu Tyr Gly Ala Pro Ile Gln Asn Arg Trp  
 210 215 220  
 Leu Val His Ile Met Ser Asp Ile Ala Asn Met Leu Ala Leu Leu Asn  
 225 230 235 240  
 Thr Ala Ile Asn Phe Phe Leu Tyr Cys Phe Ile Ser Lys Arg Phe Arg  
 245 250 255

Thr Met Ala Ala Thr Leu Lys Ala Phe Phe Lys Cys Gln Lys Gln  
 260 265 270

Pro Val Gln Phe Tyr Thr Asn His Asn Phe Ser Ile Thr Ser Ser  
 275 280 285

<210> 67

<211> 271

<212> PRT

<213> Homo sapiens

<400> 67

Thr Pro Ala Asn Met Phe Ile Ile Asn Leu Ala Val Ser Asp Phe Leu  
 1 5 10 15

Met Ser Phe Thr Gln Ala Pro Val Phe Phe Thr Ser Ser Leu Tyr Lys  
 20 25 30

Gln Trp Leu Phe Gly Glu Thr Gly Arg Cys Trp Gly Ser Leu Leu Leu  
 35 40 45

Glu Gly Gly Gly Gly Phe Pro Gly Asp Ala Leu Asn Gly Gly Trp Pro  
 50 55 60

Lys Gly Gly Asp Leu Leu Leu Gly Arg Glu Trp Val Ala Ala Leu  
 65 70 75 80

Ser Pro Val Ser Lys Gln Glu Gly Lys Met Gln Cys Trp Ser Gly Leu  
 85 90 95

Cys Gln Pro Trp Pro Asp Val Ala Gly Gly Gly Gly Val Arg Ser  
 100 105 110

Val Leu Leu Phe Leu Gly Glu Gly Gln Ser Arg Val Tyr Pro Val Pro  
 115 120 125

Arg Pro Ser Ser Pro Gly Leu Arg Ala Gly Ala Val Pro Thr Gly Cys  
 130 135 140

Glu Phe Tyr Ala Phe Cys Gly Ala Leu Phe Gly Ile Ser Ser Met Ile  
 145 150 155 160

Thr Leu Thr Ala Ile Ala Leu Asp Arg Tyr Leu Val Ile Thr Arg Pro  
 165 170 175

Leu Ala Thr Phe Gly Val Ala Ser Lys Arg Arg Ala Ala Phe Val Leu  
 180 185 190

Leu Gly Val Trp Leu Tyr Ala Leu Ala Trp Ser Leu Pro Pro Phe Phe  
 195 200 205

Gly Trp Ser Lys Trp Ala Ala Gly Thr Gly Arg Gly Ala Asp Gly Leu  
 210 215 220

Gly Gly Ala His Ser Arg Val Ser Arg Trp Thr Trp Val Ser Gln Leu  
 225 230 235 240

Ala Gly Ala Gly Cys Pro Gly Ala Thr Ala Ser Gly Glu Met Asp Ile



	245		250		255
Gln Gly Asp Met Thr Gly Ser Lys Gly Asn His Cys Pro His Leu					
	260		265		270
<210>	68				
<211>	204				
<212>	PRT				
<213>	Homo sapiens				
<400>	68				
Phe Ser Pro Arg Val Thr Asn Ser Ala Cys Val Phe Leu Leu Ser Pro					
1	5		10		15
Ala Ala Leu Ser Ala Val Cys Ala Pro Leu Cys His Pro Pro Val Pro					
	20		25		30
Ala Leu Ser Leu Gln Pro Val Ser Arg Pro Leu Ala Val Ala Met Thr					
	35		40		45
Ser Thr Cys Thr Asn Ser Thr Arg Glu Ser Asn Ser Ser His Thr Cys					
	50		55		60
Met Pro Leu Ser Lys Met Pro Ile Ser Leu Ala His Gly Ile Ile Arg					
65	70		75		80
Ser Thr Val Leu Val Ile Phe Leu Ala Ala Ser Phe Val Gly Asn Ile					
	85		90		95
Val Leu Ala Leu Val Leu Gln Arg Lys Pro Gln Leu Leu Gln Val Thr					
	100		105		110
Asn Arg Phe Ile Phe Asn Leu Leu Val Thr Asp Leu Leu Gln Ile Ser					
	115		120		125
Leu Val Ala Pro Trp Val Val Ala Thr Ser Val Pro Leu Phe Trp Pro					
	130		135		140
Leu Asn Ser His Phe Cys Thr Ala Leu Val Ser Leu Thr His Leu Phe					
145	150		155		160
Ala Phe Ala Ser Val Asn Thr Ile Val Val Val Ser Val Asp Arg Tyr					
	165		170		175
Leu Ser Ile Ile His Pro Leu Phe Tyr Pro Ser Lys Met Thr Gln Arg					
	180		185		190
Arg Gly Tyr Leu Leu Leu Tyr Gly Thr Trp Ile Val					
	195		200		

<210> 69  
 <211> 177  
 <212> PRT  
 <213> Homo sapiens

<400> 69

Gln Gly His Val Asp Arg Val His Gly Asp Val Pro Thr Cys Thr Pro

1	5							10					15			
Ala	Cys	Ser	Ser 20	Ser	Leu	Pro	Val	Arg 25	Ala	Thr	Ile	Arg	Trp 30	Pro	Arg	
Leu	Arg	Ala 35	Thr	Gly	Thr	Arg	Gly 40	His	Gly	Arg	Gly	Asp 45	Cys	Cys	Gly	
Arg	Ser 50	Leu	Gly	Asp	Ser	Cys 55	Cys	Phe	Ser	Ala	Lys 60	Ala	Leu	Cys	Val	
Trp 65	Ile	Trp	Ala	Leu	Ala 70	Ala	Leu	Ala	Ser	Leu 75	Pro	Ser	Ala	Ile	Phe 80	
Ser	Thr	Thr	Val	Lys 85	Val	Met	Gly	Glu	Glu 90	Leu	Cys	Leu	Val	Arg 95	Phe	
Pro	Asp	Lys	Leu 100	Leu	Gly	Arg	Asp	Arg 105	Gln	Phe	Trp	Leu	Gly 110	Leu	Tyr	
His	Ser	Gln 115	Lys	Val	Leu	Leu	Gly 120	Phe	Val	Leu	Pro	Leu 125	Gly	Ile	Ile	
Ile	Leu 130	Cys	Tyr	Leu	Leu	Leu 135	Val	Arg	Phe	Ile	Ala 140	Asp	Arg	Arg	Ala	
Ala 145	Gly	Thr	Lys	Gly	Gly 150	Ala	Ala	Val	Ala	Gly 155	Gly	Arg	Pro	Thr	Gly 160	
Ala	Ser	Ala	Arg	Arg 165	Leu	Ser	Lys	Val	Thr 170	Lys	Ser	Val	Thr	Ile 175	Val	

Val

```
<210> 70
<211> 356
<212> PRT
<213> Homo sapiens
```

<400> 70

Gly	Lys	Gly	Ala	Gly	Gly	Ala	Pro	Arg	Arg	Arg	Gly	Ala	Thr	Ala	Cys
1				5					10					15	
Leu	Val	Leu	Asn	Leu	Phe	Cys	Ala	Asp	Leu	Leu	Phe	Ile	Ser	Ala	Ile
			20					25					30		
Pro	Leu	Val	Leu	Ala	Val	Arg	Trp	Thr	Glu	Ala	Trp	Leu	Leu	Gly	Pro
		35					40					45			
Val	Ala	Cys	His	Leu	Leu	Phe	Tyr	Val	Met	Thr	Leu	Ser	Gly	Ser	Val
	50					55					60				
Thr	Ile	Leu	Thr	Leu	Ala	Ala	Val	Ser	Leu	Glu	Arg	Met	Val	Cys	Ile
65					70					75				80	
Val	His	Leu	Gln	Arg	Gly	Val	Arg	Gly	Pro	Gly	Arg	Arg	Ala	Arg	Ala
				85					90					95	

Val Leu Leu Ala Leu Ile Trp Gly Tyr Ser Ala Val Ala Ala Leu Pro  
 100 105 110  
 Leu Cys Val Phe Phe Arg Val Val Pro Gln Arg Leu Pro Gly Ala Asp  
 115 120 125  
 Gln Val Ser Ala Pro Leu Cys Val Pro Gly Arg Cys Pro Ala Gln Ala  
 130 135 140  
 Gly Lys Arg Gly Pro Asp Gly Ser Trp Asp Glu Asp Asp Gln Glu Gln  
 145 150 155 160  
 Gln Pro Phe Ile Ala Leu Asn Arg Cys Ala Lys Ser Cys Ala His Gly  
 165 170 175  
 Cys Glu Val Ser Leu Lys Ser His Tyr Asn Ala Val His Thr Pro Ser  
 180 185 190  
 Met Met Val Glu Ser Pro Lys Phe Leu Gln Asn Ala Met Thr Val Ala  
 195 200 205  
 Arg Leu Ile Asn Glu Phe Arg Ser Glu Thr Glu Leu Trp Gly Cys Arg  
 210 215 220  
 Val Ala Lys His Phe Val Leu Asn His Asp Ser Ser Pro Gly Ala Thr  
 225 230 235 240  
 Ile Leu Leu Leu Ser His Pro Phe His Arg Glu Pro Glu Ala His Arg  
 245 250 255  
 Ser Val Leu Ala Gln Ser Ile Leu Ser Val Ile Arg Gly Gly Asp Gly  
 260 265 270  
 Leu Leu Pro Glu Val Tyr Pro Phe Lys Glu Glu Leu Gly Ser Cys Pro  
 275 280 285  
 Gly Leu Gly Tyr Gly Pro Ala Leu Lys Ala Arg Gly Gln Ala Cys Glu  
 290 295 300  
 Thr Arg Gly Pro Gln Met Gln Cys Cys Leu Gly Thr Cys Asp Arg Ile  
 305 310 315 320  
 Pro Phe Arg Gly Leu Gln Phe Phe Gln Leu Gln Asn Gly Arg Ile Thr  
 325 330 335  
 Leu Asp Leu Arg Gly Val Leu Gly Ala Asn Val Gln Ile Thr Glu Ala  
 340 345 350  
 Gly Pro Thr Ser  
 355

<210> 71  
 <211> 187  
 <212> PRT  
 <213> Homo sapiens

<400> 71

Leu Ile Pro Gly Ser Pro Arg Thr Met Asn Pro Phe His Ala Ser Cys  
 1 5 10 15  
 Trp Asn Thr Ser Ala Glu Leu Leu Asn Lys Ser Trp Asn Lys Glu Phe  
 20 25 30  
 Ala Tyr Gln Thr Ala Ser Val Val Asp Thr Val Ile Leu Pro Ser Met  
 35 40 45  
 Ile Gly Ile Ile Cys Ser Thr Gly Leu Val Gly Asn Ile Leu Ile Val  
 50 55 60  
 Phe Thr Ile Ile Arg Gly Met Ala Pro Phe Phe Phe Phe Ser Phe His  
 65 70 75 80  
 Thr Leu Gly Asn Tyr Ser Gln Ser Ser Leu Asn Glu Ser Phe Pro Leu  
 85 90 95  
 His Phe Ala Asn Ala Ile Leu Glu Ile Phe Ala Asp Asn Arg Phe Tyr  
 100 105 110  
 Arg Tyr Phe Arg His Ala Leu Gly Asn Ile Ser Val Ala Asn Leu Asn  
 115 120 125  
 Met Leu Arg Pro Arg Pro Met Leu Thr Ala Gly Ser Val Ser Phe Gly  
 130 135 140  
 Asn Asp Ser His Ser Gln Ser Leu His Ile Arg Ile Ile Trp Gly Ala  
 145 150 155 160  
 Leu Lys Thr Pro Asp Met Gln Phe His Pro Arg Pro Ile Asn Ser Glu  
 165 170 175  
 Ser Leu Arg Val Gly Pro Gly Asp Leu Lys Leu  
 180 185

<210> 72  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens

<400> 72

Val Thr Gly Thr Leu Glu Gly Asp Ser Phe Pro Gly Glu Glu Gly Thr  
 1 5 10 15  
 Val Gly Phe Pro Ala Leu Leu Leu Ser Pro Tyr Gly Leu Arg Ser Gly  
 20 25 30  
 Leu Asn Cys Asp Glu Ser Leu Cys Pro Cys Ser Leu Gly Phe Leu Lys  
 35 40 45  
 Tyr Pro Pro Phe Ile Val Pro Ile Ile Ile Asn Thr Asn Gln Ser Leu  
 50 55 60  
 Leu Cys Leu Lys Leu Leu Ile Ala Tyr Arg Gly Lys Cys Arg Leu Leu  
 65 70 75 80  
 Ser Trp Arg Thr Ser Leu Leu His Leu Ala His Ile Glu Leu Ser Arg  
 50

85 90 95  
 Ala Ile Ser Cys His Ser Leu Ala Cys Leu Leu Leu Leu Trp Val Ser  
 100 105 110

Arg Thr Ala His Glu Phe Pro His Met Pro Phe  
 115 120

<210> 73  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 73

Tyr Ala Phe Leu Leu Met Asp Gln Ala Ile Ser Pro Thr Ser Phe Phe  
 1 5 10 15

Leu Ile Ile Pro Phe Leu Ala Leu Phe Ser Arg Lys Lys Phe Lys Cys  
 20 25 30

Leu Leu Pro Ile Phe Ser Glu Gly Glu Met Pro Val Pro Gly Phe Pro  
 35 40 45

Gly Ser Cys Leu Leu Gly Arg Pro Leu Pro Cys Thr Thr Lys Ser Thr  
 50 55 60

Leu Pro Ser Gln His Pro Leu Leu Ser Pro Gly Gln Leu Leu Cys Val  
 65 70 75 80

Leu Phe Ile Pro Ile Ser Leu Pro Glu Leu Leu Arg Pro Leu Cys Leu  
 85 90 95

Ser Ala Ser Cys Pro Ile Phe Gln Ala Leu Val Cys Trp Leu Ser Ala  
 100 105 110

Ser Lys Asn Asp Phe Lys His Leu Val Phe Leu Ser Thr Glu Leu Gln  
 115 120 125

Thr Leu Lys Cys Arg Ser Ser Ile Thr Ser Asn Ile Glu Val Leu Ile  
 130 135 140

Met Leu Pro Cys Gly Phe Met Leu Phe Gly Ile Asp Phe  
 145 150 155

<210> 74  
 <211> 211  
 <212> PRT  
 <213> Homo sapiens

<400> 74

Lys Gly Gly Cys Thr Lys Glu Gly Asn Lys Tyr Ser Leu Met Ile Tyr  
 1 5 10 15

Asn Val Ser Asp Thr Glu Ile His Arg Leu Gly Leu Trp Glu Ser Gln  
 20 25 30

Ile Leu Ile Cys Gln Gln Cys Val Leu Ile Phe Leu Phe Cys Ile Phe  
 51

35                      40                      45  
 Ala Ser Pro Ile Asn Phe Thr Gly Leu Ile Ile Phe Thr Arg His Tyr  
 50                      55                      60  
 Phe Lys Glu Lys Ala Asn Leu Leu Lys Phe Tyr Ala Ser Leu Ile Val  
 65                      70                      75                      80  
 Met Ser Trp Leu Ile Leu Ala Leu Val Ser Met Phe Trp Lys Ile Leu  
 85                      90                      95  
 Lys Tyr Phe His Leu Val Ala Leu Ser Glu His Cys Arg Thr Glu Val  
 100                      105                      110  
 Ile Pro Leu Gly Phe Ser Asn Lys Ser Glu Leu Leu Gly Thr Leu Asn  
 115                      120                      125  
 Gln Phe Met Arg Thr Phe Cys Tyr Thr Cys Glu Phe Thr Tyr Leu Phe  
 130                      135                      140  
 Leu Cys Ser Arg Met Gly Cys Met Gln Arg Arg Met Asn Ser Thr Thr  
 145                      150                      155                      160  
 Val Met Glu Arg Ser Asp Arg Arg Asn Ser Lys Met Pro Ile Cys Tyr  
 165                      170                      175  
 Leu Ser Gly Thr His Asn Ile Ser Asn Ala Glu Arg Asn Met Leu Gly  
 180                      185                      190  
 Ser Ser Gln Thr Ile Val Ile Lys Tyr Leu Leu Asn Ile His Ile Leu  
 195                      200                      205  
 Gly Ser Cys  
 210

<210> 75  
 <211> 120  
 <212> PRT  
 <213> Homo sapiens

<400> 75

Phe Ile Ile Asn Ile Ser Leu Tyr Ala Ile Ile Cys Phe Leu Ala Ile  
 1                      5                      10                      15  
 Tyr Ala Asp Ser Phe Val Cys Phe Ser Phe Arg Ile Phe Leu Asn Arg  
 20                      25                      30  
 Phe Leu Val Phe His Arg Val Phe Val Cys Leu Lys Phe Cys Leu Phe  
 35                      40                      45  
 Ile Phe Leu Leu Ser Met Phe Ser His Asn Tyr Leu Thr Phe Ser Asn  
 50                      55                      60  
 Ile Ile Leu Thr Ile Leu Val Ser Leu Leu Phe Trp Ile Asn Phe Arg  
 65                      70                      75                      80  
 Ile Phe Phe Phe Glu Ser Pro Lys Lys Ser Leu Trp Ile Leu Asp Asp  
 85                      90                      95

Thr Ser Tyr Glu Thr Thr Tyr Gly Leu Phe Phe Ser Leu Ile Phe Ser  
 100 105 110

Leu Pro Tyr Arg Asn Val Phe Phe  
 115 120

<210> 76  
 <211> 239  
 <212> PRT  
 <213> Homo sapiens

<400> 76

Arg Leu Leu Leu Cys Gly Tyr Asn Glu Ser His Gln Leu Gln Ser Phe  
 1 5 10 15

Pro Lys Lys Asp Arg Lys Ile Cys Leu Ile Lys Met Val Phe Lys Ala  
 20 25 30

Asn Tyr Arg Tyr His Arg Lys Lys Lys Asn Ser Gly Arg Asp Leu  
 35 40 45

Ser Ser Phe Ser Tyr Pro Met Gly Phe Ser Ala Ser Arg Phe Pro Cys  
 50 55 60

Thr Ser Arg Gly Phe Cys Ala Asn Gln Trp Ala Trp Pro Ser Ala His  
 65 70 75 80

Pro Gly Leu Ser Cys Gly Cys His Gly Tyr Glu Asp Lys Asn Asp Pro  
 85 90 95

Trp Phe Tyr Arg Ala Leu Gly Pro Leu Leu Thr Gly Leu Phe Ser Tyr  
 100 105 110

Phe Phe His Leu Leu Gln Asn Pro Met Trp Gln Val Gln Gln Gly Val  
 115 120 125

His Gln Cys Ser Gly Gln Ser Glu Gly Thr His Ala Arg Pro Gly Arg  
 130 135 140

Ala Ala Leu Lys Lys Trp Phe Val Arg Val Trp Arg Lys Gly Leu Leu  
 145 150 155 160

Pro His Ser Ala Leu Leu Gly Pro Lys Ala Leu Ile Thr Pro Gly Ser  
 165 170 175

Gly Ile Ile Leu Ser Leu Met Leu Leu Leu Ser Arg Asp Arg Thr Arg  
 180 185 190

Gly Ala Ser Thr Gln Gly Asn Pro Ala Pro Gln Gln Ala Gly Asp His  
 195 200 205

His Asp Cys Asn Arg Gln Gln Glu Ser Glu Leu Trp His Lys Asn Val  
 210 215 220

Thr Ala Thr Pro Gln Gly Gly Thr Pro Gly Gly Lys Gly His Gly  
 225 230 235

<210> 77  
 <211> 228  
 <212> DNA  
 <213> Homo sapiens

<400> 77  
 attatctcca ttatgctcta tcagtttctt tatagaatat gattactact tagtgataaa 60  
 gcttccttttg ctaagatttc agcctacgaa ccatgatcca aaccctactt ctaaaacata 120  
 taaacatgct ttacaagtat cctatatatg gaaaagtcct tggaattatt tgggtaatta 180  
 acctgggtatt acatgtgtta tttcctgata tactattgca aattgaca 228

<210> 78  
 <211> 239  
 <212> PRT  
 <213> Homo sapiens

<400> 78  
 Leu Ile Tyr Thr Lys Asn His Asn Leu Lys Lys Thr Phe Asn Tyr Leu  
 1 5 10 15  
 Tyr Cys Cys Ile Lys Cys Phe Ile Tyr Glu Ser Lys Asn Leu Phe Phe  
 20 25 30  
 Ala Phe Ile Leu Cys His Arg Lys Lys Thr Ser Cys Leu Cys Leu Leu  
 35 40 45  
 Pro Leu Ile Lys Ala Asn Asn Arg Gln Tyr Met Met Ile Leu Leu Thr  
 50 55 60  
 Arg Ala Asn Lys Ser Asn Ile Leu Thr Met Arg Phe Gly Leu Cys Tyr  
 65 70 75 80  
 Leu Tyr Leu Met Ser Asn Phe Pro Tyr Asn Val Asn Val Ile Pro Pro  
 85 90 95  
 Leu Tyr Pro Phe Val Arg Val Ser Val Phe Cys Ser Cys Met Thr Tyr  
 100 105 110  
 Phe Lys Gly Ser Gln Gly Gln Val Ile His Ile Leu Lys Ile Pro Phe  
 115 120 125  
 Arg Ile Ile Thr Ser Lys Asn Ser Gln Glu Arg Gln Phe Gln Tyr Lys  
 130 135 140  
 Ile Ile Asn Tyr Tyr Trp Asn Phe Lys His Ser Trp Gln Lys Arg Asp  
 145 150 155 160  
 Asn Leu Gln Asn Ile Leu Met Ala Thr Phe Leu Phe Tyr His Leu Leu  
 165 170 175  
 Cys Ser Ile Cys Tyr Asp Gln Arg Asn Tyr Ser Ile Ser Thr Thr Ile  
 180 185 190  
 His Lys Thr Gly Met Glu Glu Val Phe Phe Phe Leu Val Leu Met Ser  
 54



195                      200                      205  
 Lys Lys Met Asn Leu Arg Ser Met Ser Phe Phe Asn Thr Met Phe Thr  
     210                      215                      220  
 Asp Ser Thr His Tyr Phe Met Asp Thr Lys Ser Gln Thr Tyr Ile  
 225                      230                      235  
 <210> 79  
 <211> 203  
 <212> PRT  
 <213> Homo sapiens  
 <400> 79  
 Phe Ile Leu Thr Ile Ile Tyr Tyr Gln Arg Asn Leu Arg Lys Gln Lys  
 1                      5                      10                      15  
 Phe Arg Ile Met Phe Leu Thr Cys Val Ser Phe Phe Thr Val Val Leu  
                     20                      25                      30  
 Glu Leu Ser Tyr Pro Trp Gln Leu Ile Leu Leu Leu Tyr Ser Leu Phe  
                     35                      40                      45  
 Ile Val Leu Thr Leu Lys Leu Arg Thr Ser Asn Val Asn Leu Asp Glu  
                     50                      55                      60  
 Leu Lys Thr Glu Asn Val Cys Lys Tyr Val Lys Tyr Val Tyr Lys Asn  
 65                      70                      75                      80  
 Met Tyr Phe Ser Tyr Phe Lys Ser Phe Ile Leu Tyr Ile Thr His Thr  
                     85                      90                      95  
 His Thr His Thr His Thr Met Arg Ser Leu Leu Thr Thr Gln Tyr Lys  
                     100                      105                      110  
 Ile Ile Phe Leu Arg Asn Ile Val Phe Lys Tyr Cys Phe Ile Pro Tyr  
                     115                      120                      125  
 Lys Ser Asn Leu Trp Leu Phe Tyr Gly Phe His Gln Ala Met Ser Leu  
                     130                      135                      140  
 Thr Asn Phe Ala Asn Lys Gly Thr Gln Gly Met Lys Tyr Leu Leu Thr  
 145                      150                      155                      160  
 Asn Lys Lys Pro Ser Asn Ser Met Tyr Val Ile Gly Lys Ile Lys Ser  
                     165                      170                      175  
 Ser Val Asn Ser Ile His Glu Leu Thr Ser Ile Ser Ala Leu Leu Ser  
                     180                      185                      190  
 Leu Lys Ile Ser Asn Ser Leu Lys Ile Ile Arg  
                     195                      200  
 <210> 80  
 <211> 156  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 80

Ser Lys Ser Leu Leu His Ser Ser Leu Leu Gly Val Leu Val Ile Cys  
 1 5 10 15  
 Thr Ser Ser Phe Pro Ser Ala Gly Lys Ala Ile Asn Gly Phe Gln Leu  
 20 25 30  
 Tyr Ala Ala Pro Leu Gly Cys Ser Phe Asp Leu Ser Gln Val Ser Trp  
 35 40 45  
 Ala Ile Glu Asn Asp Leu Ser Ser Cys Arg Gly Leu Gln Pro Ile Leu  
 50 55 60  
 Asn Ile Ala Gly Ile Asn Leu Phe Leu Leu Lys Glu Gly Ile Leu Leu  
 65 70 75 80  
 Tyr Arg Val Leu Leu Asn Thr Leu Leu Asp Phe Phe Asp Phe Val Phe  
 85 90 95  
 Leu Leu Ser Glu Tyr Leu Lys Tyr Pro Cys Arg Tyr Ser Asn Leu Phe  
 100 105 110  
 Pro Val Tyr Gly Phe Phe Leu Cys Leu Leu Trp Thr Gln Val Phe Val  
 115 120 125  
 Ala Cys Phe Ser Lys Ser Ala Gln Lys Leu Leu Val Ile Gly Phe Gly  
 130 135 140  
 Ser Leu Phe Lys Leu Tyr Pro Pro Glu Ser Tyr Arg  
 145 150 155

&lt;210&gt; 81

&lt;211&gt; 208

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 81

Cys Glu Arg Arg Asp Glu Pro Asn Thr Ala Thr Thr Ala Arg Pro Trp  
 1 5 10 15  
 Gly Ser Val Val Gln Met Gly Arg Cys Thr Glu His Ala Gly His Val  
 20 25 30  
 Cys Gln Pro Ser Leu Cys Ala Leu Ala Thr Val Lys Arg Gly Leu Cys  
 35 40 45  
 Leu Gly Arg Leu Tyr His Trp Ser Cys Ile Pro Asn Val Ser Leu Phe  
 50 55 60  
 Thr Arg Leu Tyr Lys His Ala Gln Tyr Leu Cys Ile Phe Thr Val Thr  
 65 70 75 80  
 Ser Phe Ser Tyr Ile Ile Lys Val Pro Ile Tyr Ile Pro Tyr Met Tyr  
 85 90 95  
 Ile Tyr Asn Thr Gln Met His Thr His Lys His Thr Cys Thr Glu Leu  
 100 105 110

Leu Asp Ser Gln Ser Leu Ile Leu Cys Ile Phe Thr Val Thr Ser Phe  
 115 120 125  
 Ser Tyr Ile Ile Lys Val Pro Ile Tyr Ile Pro Arg Ser Thr Glu Val  
 130 135 140  
 Gln Ser Ser Pro Ser Ser Asp Leu Met Ile Ile Leu Glu His Leu Ile  
 145 150 155 160  
 Phe Leu Tyr Phe Ile Met Tyr Leu Ile Ile Thr Leu Phe Ser Glu Cys  
 165 170 175  
 Gln Leu Ile Arg Arg Met Thr Cys Leu Leu Ser Cys Ser Ser Tyr Pro  
 180 185 190  
 Pro Ile Gly Phe Ile Ser Pro Leu Thr Arg Cys Gly Asp Trp Ala Gly  
 195 200 205

<210> 82  
 <211> 97  
 <212> PRT  
 <213> Homo sapiens

<400> 82

Arg Gln Arg Leu Cys Glu Thr Leu His Asp Arg Gly Leu Cys Ile Leu  
 1 5 10 15  
 Ala His Phe Leu Ala Leu Cys Glu Gln Tyr Leu Ser Ser His Tyr Ile  
 20 25 30  
 Lys Ser Ile Glu Pro Ile Gln Arg Ser Glu His Thr Val Met Ser Trp  
 35 40 45  
 Leu Leu Pro Ser Lys Ala Trp Gly Leu Val Leu Tyr Gly Ser Pro Ala  
 50 55 60  
 His Ile Leu Lys Phe Leu Cys Gly Thr Leu Phe Glu Leu Phe Tyr Phe  
 65 70 75 80  
 Glu Cys Ala Ile Phe Leu Leu Gly Leu Leu Lys Gly Arg Ala Leu Leu  
 85 90 95

Ser

<210> 83  
 <211> 201  
 <212> PRT  
 <213> Homo sapiens

<400> 83

Lys Trp Ile Asn Leu His Met Asp Glu His Asp Leu Leu Leu Ser Arg  
 1 5 10 15  
 Ser Gln Arg Ile His Lys Lys Lys Asn Leu Val Met Leu Leu Asp Asp  
 20 25 30

Val Phe Asp Asn Thr Ile Gln Tyr Leu Ser Met Tyr Pro Tyr Asp Ile  
           35                          40                          45  
 Glu Lys Gly Phe Ser Lys Tyr Phe Asn Leu Asn Arg Phe Thr Lys Arg  
           50                          55                          60  
 Asn His Leu Pro Thr Thr Val Pro Cys Leu Trp Ser Ile Arg Val Ile  
   65                          70                          75                          80  
 Ile Leu Phe Ser Leu Tyr Tyr Lys Arg Glu Cys Thr Leu Tyr Lys Ile  
                           85                          90                          95  
 Asn Asn Ile Asp Tyr Ile Ser Arg Ile Asn Thr Lys Ala Gln Lys Tyr  
                           100                          105                          110  
 Lys Ser Lys Pro Leu Asn Ser Leu Pro Pro Thr Ser Arg Lys Thr Gln  
           115                          120                          125  
 Phe Leu Tyr His Ser Tyr Tyr Lys Ser Phe Leu Ser Ser Phe Ser Tyr  
           130                          135                          140  
 Lys His Ala Glu Ile Tyr Met Trp Thr His Leu Cys Gln Asn Ser Phe  
   145                          150                          155                          160  
 Phe Ser Asp Thr His Phe Phe Pro Pro Thr Pro Gln Tyr Val Arg Thr  
                           165                          170                          175  
 Ile Phe Tyr Lys Ile Gln Met Glu His Arg Ile Glu Gln Asn Met Lys  
           180                          185                          190  
 Thr Ser Asn Ser Leu Val Asp Asp Val  
           195                          200

<210> 84  
 <211> 212  
 <212> PRT  
 <213> Homo sapiens

<400> 84

Ser Pro Leu Cys Ser Tyr Phe Ser Leu Cys Asn Ile Ala Cys Arg Glu  
   1                          5                          10                          15  
 Met Val Asp Met Asn Ile Leu Ser Leu Asp His Ala Phe Gly His Tyr  
           20                          25                          30  
 Phe Gln His Cys Cys Lys Leu Phe Phe Lys Arg Pro Ile Tyr Phe Ile  
           35                          40                          45  
 Asp Met Tyr Ile Phe Phe Lys Lys Ser Lys Leu Thr Val Phe Leu Lys  
   50                          55                          60  
 Ala Gln Asn Pro Leu Val Trp Leu Thr Ser Leu Ile Phe Glu Lys Ala  
   65                          70                          75                          80  
 Leu Cys Phe Ser Asn Leu Leu Val Phe Ala Asn Gln Ile Gly Val Lys  
                           85                          90                          95

Lys Ile Phe His Tyr Thr Phe Lys Gly Leu Leu Leu Thr Glu Leu Glu  
100 105 110

His Leu Phe Met Phe Glu Ser Phe Val Phe His Phe Leu Ile Thr Val  
115 120 125

Leu Ile Ile Cys Leu Phe His Thr Tyr Leu Gln Lys Phe Leu Lys Tyr  
130 135 140

Ile Lys Glu Ser Asn Tyr Arg Ile Glu Lys Tyr Ile Gln Glu Asn Ser  
145 150 155 160

Pro Leu Ser Val Thr Gln Val Ala Ser Ile Leu Val Asn Phe Val Phe  
165 170 175

Ala Phe Tyr Leu Phe Ile Leu Ile Cys Tyr Phe His Lys Ile Phe Cys  
180 185 190

Cys Phe Leu Phe Cys Arg Ile Tyr Phe Val Cys Gly Arg Gly Phe Leu  
195 200 205

Glu Glu Pro Ser  
210

<210> 85

<211> 192

<212> PRT

<213> Homo sapiens

<400> 85

Gln Lys Ser Leu Leu Cys Ser Lys Ser Leu His Gly Phe Leu Leu Tyr  
1 5 10 15

Leu Glu Tyr Asn Pro Lys Ser Phe Pro Arg Ala Leu Ala Leu Tyr Trp  
20 25 30

Pro Ser Leu Leu Leu Pro Thr Ser Tyr His Leu Ser Phe Gly His Tyr  
35 40 45

Ser Pro Thr Thr Ser Thr His Pro Leu Ser Arg Arg Pro Trp Asn Ile  
50 55 60

Leu Ser Thr Phe Pro Ser Gln Gly Val Val Asn Ala Val Ser Ser Thr  
65 70 75 80

Cys Ser His Pro Pro Thr Leu Tyr Gln Leu Ser Ser His Ile His Met  
85 90 95

Phe Asp Leu Lys Leu Cys Val Phe Thr Cys Leu Leu Phe Glu Ala Leu  
100 105 110

Cys Ala Ile Val Glu Ser Glu Val Lys Gly Lys Ile Ser Asn Ile Asp  
115 120 125

Ser Glu Phe Met Tyr Phe Asp Phe Phe Pro Arg Asn Phe Ala Leu Ile  
130 135 140

Phe Ile Leu Lys Ile Leu Tyr Asn Phe Ile Tyr Leu Asp Tyr Val Phe  
59

145                                      150                                      160  
 Trp Cys Pro Leu Lys Ile Leu Thr Lys Gly Gln Val Pro Arg Ser Pro  
    165                                      170                                      175  
 His Pro Phe Ser Ser Thr Gly His Trp Ser Val Pro Pro Asn Cys Lys  
    180                                      185                                      190  
  
 <210> 86  
 <211> 227  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 86  
  
 Glu Lys His Ala His Pro Gly Pro Arg Pro Pro Ala Arg Tyr Lys Asp  
 1                                      5                                      10                                      15  
 Gly Ser Gly Pro Ala Trp Glu Gln Phe Pro Trp Cys Gly Pro Pro Asn  
    20                                      25                                      30  
 Leu Gly Gly Lys Leu Lys Ser Arg Thr Ile Ser Ile His Glu Glu Asn  
    35                                      40                                      45  
 Glu Glu Ala Glu Tyr Leu Phe Asn Glu Ser Ala Val Gly Lys Gly Ile  
    50                                      55                                      60  
 Tyr Leu Tyr Tyr Tyr Leu Val Glu Lys Asn Leu Leu Trp Pro Asn Val  
 65                                      70                                      75                                      80  
 Lys Ser Ile Phe Phe Asp Lys Leu Lys Arg Leu Arg Lys Ile Gly Pro  
    85                                      90                                      95  
 Ile Leu Thr Glu Asn Gly Val Thr Arg Gly Val Ile Ser Glu Asn Leu  
    100                                      105                                      110  
 Ala Pro Ala Pro Phe Cys Tyr Gly Glu Lys Tyr Gln Tyr Tyr Leu Leu  
    115                                      120                                      125  
 Arg Arg Leu Gly Leu Asn Trp Lys Arg Gly Trp Leu Leu Cys Ala Asp  
    130                                      135                                      140  
 Gly Asp Lys Ile Arg Ile Leu Pro Ser Leu Gln Ser Gln Met Gln His  
 145                                      150                                      155                                      160  
 Thr Gln Arg Cys Tyr Leu Gly Ala Gly Lys Gln Gly Asp Ile Lys Met  
    165                                      170                                      175  
 Tyr Val Val Ile Thr Asn Gly Arg Ser Lys Phe His Ser Asp Tyr Phe  
    180                                      185                                      190  
 Lys Pro Asn Ala Gly Leu Leu Tyr Ser Phe Glu Ile Phe Phe Ser Val  
    195                                      200                                      205  
 Asn Cys Asn Arg Gly Lys Asn Gln Pro His Ile Asn Phe Ser Gln Gly  
    210                                      215                                      220  
 Lys Pro Ser  
 225

<210> 87  
 <211> 125  
 <212> PRT  
 <213> Homo sapiens

<400> 87

Glu Gly Tyr Thr Phe Asp Lys Arg Leu Ile Cys Phe Thr Val Trp Val  
 1 5 10 15  
 Glu Asn Val Lys Ser Ile Asn Asn Phe Thr Ser Phe Phe His Leu Arg  
 20 25 30  
 Ser Tyr Tyr Ser Leu Tyr Tyr Ala Tyr Pro Pro Arg Ile Tyr Leu His  
 35 40 45  
 Ser Ala Ser Ser Cys Gln Met Ser Ala Val Lys Val Pro Pro Gly Asn  
 50 55 60  
 Phe Arg Glu Thr Arg Lys Thr Ser Ser Ile Asn Val Tyr Phe Asn Leu  
 65 70 75 80  
 Phe Ser Ile Leu Phe Leu Lys Lys Tyr Gln Ser Leu Lys Arg Trp Phe  
 85 90 95  
 Thr Phe Gly Leu Phe His Cys Leu Pro Met Lys Ala Leu Leu Val Leu  
 100 105 110  
 Phe Tyr Asn Leu Met Asp Phe Ser Pro Arg Phe Tyr Thr  
 115 120 125

<210> 88  
 <211> 174  
 <212> PRT  
 <213> Homo sapiens

<400> 88

Met Leu Trp Leu Phe Val Pro Arg Val Arg Gln Lys Arg Gln Trp Leu  
 1 5 10 15  
 Pro Ser Leu Gly Val Thr Gly Thr Leu Glu Gly Asp Ser Phe Pro Gly  
 20 25 30  
 Glu Glu Gly Thr Val Gly Phe Pro Ala Leu Leu Leu Ser Pro Tyr Gly  
 35 40 45  
 Leu Arg Ser Gly Leu Asn Cys Asp Glu Ser Leu Cys Pro Cys Ser Leu  
 50 55 60  
 Gly Phe Leu Lys Tyr Pro Pro Phe Ile Val Pro Ile Ile Ile Asn Thr  
 65 70 75 80  
 Asn Gln Ser Leu Leu Cys Leu Lys Leu Leu Ile Ala Tyr Arg Gly Lys  
 85 90 95  
 Cys Arg Leu Leu Ser Trp Arg Thr Ser Leu Leu His Leu Ala His Ile  
 100 105 110

Glu Leu Ser Arg Ala Ile Ser Cys His Ser Leu Ala Cys Leu Leu Leu  
 115 120 125  
 Leu Trp Phe Ser Arg Thr Ala His Glu Phe Pro His Met Pro Phe Trp  
 130 135 140  
 Phe Gly His Leu Thr Phe Ser Ala Ala Val Pro Ser Ala Trp Thr Ala  
 145 150 155 160  
 Phe Pro Pro Arg Ser Leu Gly Glu Leu Leu Phe Ile Leu Gln  
 165 170

<210> 89  
 <211> 182  
 <212> PRT  
 <213> Homo sapiens

<400> 89

Tyr Lys Val Val Val Leu Asp Asp Leu Tyr Thr Trp Lys Leu Val Tyr  
 1 5 10 15  
 Phe Phe Asn Lys Ala Ser Asp Ser Lys Val Leu Trp Ala Ile Cys Leu  
 20 25 30  
 Cys Cys Asn Tyr Ser Ala Leu Pro Leu Cys Lys Ser Ser His Arg Tyr  
 35 40 45  
 Val Asn Glu Thr Trp Leu Phe Ser Gln Lys Gln Ala Ala Val Arg Ile  
 50 55 60  
 Trp Leu Ala Asp Ser Leu Gly Phe Ser Leu Leu Lys Ala Trp Ser Arg  
 65 70 75 80  
 Arg Pro Ala Ala Ser Ala Leu Pro Val Asn Leu Leu Glu Ile Gln Asn  
 85 90 95  
 Val Glu Pro Leu Leu Ile Leu Ile Ala Leu Glu Ser Lys Pro Cys Phe  
 100 105 110  
 Ser Asn Phe Ser Ile His Arg Thr His Leu Gly Ile Leu Lys Tyr Arg  
 115 120 125  
 Leu Asp Ser Arg Ala Glu His His Thr Ser Leu Gly Glu Ala Met Leu  
 130 135 140  
 Pro Gly Gln Cys Ser Leu Gly Ser Pro Gln Ser Ile Arg Arg Asp Cys  
 145 150 155 160  
 Tyr Ile Asn Lys Phe Gln Tyr Arg Leu Val Asn Ala Ile Thr Glu Val  
 165 170 175  
 Ser Ser Ala Glu Glu His  
 180

<210> 90  
 <211> 188  
 <212> PRT



&lt;213&gt; Homo sapiens

&lt;400&gt; 90

Arg Val Tyr Leu Leu Trp Lys Leu Thr Leu Ala Ser Phe Gln Asn Thr  
 1 5 10 15  
 Phe Phe Ser Gln Lys His Leu Arg Val Ser Gln Phe Ser Cys Lys Thr  
 20 25 30  
 Gln Leu Pro Leu Leu Gly Asn Cys Leu Gln Asp Tyr Leu Phe Tyr Gly  
 35 40 45  
 Lys Lys Thr Asn Asn Leu Tyr Phe Thr Thr Lys Phe Leu Ser Leu His  
 50 55 60  
 Glu Ser Tyr Ser Leu Glu Ile Gln Leu Phe Pro Lys Leu Lys Ile Glu  
 65 70 75 80  
 Met Ser Ser Pro Phe Ser Gly Glu Pro Phe Pro Val Leu Glu Asp Lys  
 85 90 95  
 Ser Phe Gln Gln Arg Cys Glu Gln Met Asn Tyr Leu Leu Trp Thr Asp  
 100 105 110  
 Phe Thr Asp Gly Leu Asn Cys Phe Ser Lys Pro Cys Gln Leu Phe Cys  
 115 120 125  
 Asn Phe Trp Ser Ser Ile Phe Leu Thr Met Cys Cys Ala Val Leu Ile  
 130 135 140  
 Tyr Leu Ala Lys Val Val Leu Ala Asn Val Met Gln Ala Glu Thr Trp  
 145 150 155 160  
 Lys Lys Lys Ser Val His Phe Ser Leu Phe Tyr Tyr Cys Val Thr Trp  
 165 170 175  
 Pro Ala Leu Ser Trp Met Ile Ala Ile Asn Met Lys  
 180 185

&lt;210&gt; 91

&lt;211&gt; 333

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 91

Met Trp Ser Cys Ser Trp Phe Asn Gly Thr Gly Leu Val Glu Glu Leu  
 1 5 10 15  
 Pro Ala Cys Gln Asp Leu Gln Leu Gly Leu Ser Leu Leu Ser Leu Leu  
 20 25 30  
 Gly Leu Val Val Gly Val Pro Val Gly Leu Cys Tyr Asn Ala Leu Leu  
 35 40 45  
 Val Leu Ala Asn Leu His Ser Lys Ala Ser Met Thr Met Pro Asp Val  
 50 55 60

Tyr Phe Val Asn Met Ala Val Ala Gly Leu Val Leu Ser Ala Leu Ala  
 65 70 75 80  
 Pro Val His Leu Leu Gly Pro Pro Ser Ser Arg Trp Ala Leu Trp Ser  
 85 90 95  
 Val Gly Gly Glu Val His Val Ala Leu Gln Ile Pro Phe Asn Val Ser  
 100 105 110  
 Ser Leu Val Ala Met Tyr Ser Thr Ala Leu Leu Ser Leu Asp His Tyr  
 115 120 125  
 Ile Glu Arg Ala Leu Pro Arg Thr Tyr Met Ala Ser Val Tyr Asn Thr  
 130 135 140  
 Arg His Val Cys Gly Phe Val Trp Gly Gly Ala Leu Leu Thr Ser Phe  
 145 150 155 160  
 Ser Ser Leu Leu Phe Tyr Ile Cys Ser His Val Ser Thr Arg Ala Leu  
 165 170 175  
 Glu Cys Ala Lys Met Gln Asn Ala Glu Ala Ala Asp Ala Thr Leu Val  
 180 185 190  
 Phe Ile Gly Tyr Val Val Pro Ala Leu Ala Thr Leu Tyr Ala Leu Val  
 195 200 205  
 Leu Leu Ser Arg Val Arg Arg Glu Asp Thr Pro Leu Asp Arg Asp Thr  
 210 215 220  
 Gly Arg Leu Glu Pro Ser Ala His Arg Leu Leu Val Ala Thr Val Cys  
 225 230 235 240  
 Thr Gln Phe Gly Leu Trp Thr Pro His Tyr Leu Ile Leu Leu Gly His  
 245 250 255  
 Thr Gly Ile Ile Ser Arg Gly Lys Pro Val Asp Ala His Tyr Leu Gly  
 260 265 270  
 Leu Leu His Phe Val Lys Asp Phe Ser Lys Leu Leu Ala Phe Ser Ser  
 275 280 285  
 Ser Phe Val Thr Pro Leu Leu Tyr Arg Tyr Met Asn Gln Ser Phe Pro  
 290 295 300  
 Ser Lys Leu Gln Arg Leu Met Lys Lys Leu Pro Cys Gly Asp Arg His  
 305 310 315 320  
 Cys Ser Pro Asp His Met Gly Val Gln Gln Val Leu Ala  
 325 330

&lt;210&gt; 92

&lt;211&gt; 97

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 92

Thr Arg Leu Trp Gly Thr Lys Leu Asn Ala Gly Leu Gly Gln Arg Leu  
 64

1	5					10					15				
Met	Pro	Cys	Gly	His	Asp	Val	Met	Leu	Ser	Ile	Leu	Leu	Pro	Ser	Arg
			20					25					30		
Gly	Ser	Arg	Ser	Gly	Ser	Arg	Arg	Gly	Ala	Leu	Leu	Leu	Glu	Gly	Ala
		35					40					45			
Ser	Arg	Asp	Met	Glu	Lys	Val	Asp	Met	Asn	Thr	Ser	Gln	Glu	Gln	Gly
	50					55					60				
Leu	Cys	Gln	Phe	Ser	Glu	Lys	Tyr	Lys	Gln	Val	Tyr	Leu	Ser	Leu	Ala
65					70					75					80
Tyr	Ser	Ile	Ile	Phe	Ile	Leu	Gly	Leu	Pro	Leu	Asn	Gly	Thr	Val	Leu
				85					90					95	

Trp

```
<210> 93
<211> 140
<212> PRT
<213> Homo sapiens
```

<400> 93

Ser	Ala	Arg	Ala	His	Lys	Val	Pro	Glu	Ala	His	Arg	Phe	Ser	Gly	Leu
1				5					10					15	
Gly	Ile	Ala	Tyr	Ser	Asn	Ser	Trp	Ala	Pro	Asn	Ser	Lys	Ser	Pro	Thr
			20					25					30		
Thr	Leu	Ser	His	Leu	Trp	Leu	Gln	Ser	Cys	Phe	Gly	Phe	Pro	Gln	Ile
		35					40					45			
Thr	Lys	Ala	Ser	Arg	Lys	Arg	Leu	Thr	Val	Ser	Leu	Ala	Tyr	Ser	Glu
	50					55					60				
Ser	His	Gln	Ile	Arg	Val	Ser	Gln	Gln	Asp	Phe	Arg	Leu	Phe	Arg	Thr
65					70					75					80
Leu	Phe	Leu	Leu	Met	Val	Ser	Phe	Phe	Ile	Met	Trp	Ser	Pro	Ile	Ile
				85					90					95	
Ile	Thr	Ile	Leu	Leu	Ile	Leu	Ile	Gln	Asn	Phe	Lys	Gln	Asp	Leu	Val
			100					105					110		
Ile	Trp	Pro	Ser	Leu	Phe	Phe	Trp	Val	Val	Gly	Phe	Thr	Phe	Ala	Asn
		115					120					125			
Ser	Ala	Leu	Asn	Pro	Ile	Leu	Tyr	Asn	Met	Thr	Leu				
	130					135					140				

```
<210> 94
<211> 177
<212> PRT
<213> Homo sapiens
```

&lt;400&gt; 94

Asp Ser Val Ser Tyr Glu Tyr Gly Asp Tyr Ser Asp Leu Ser Asp Arg  
 1 5 10 15  
 Pro Val Asp Cys Leu Asp Gly Ala Cys Leu Ala Ile Asp Pro Leu Arg  
 20 25 30  
 Val Ala Pro Leu Pro Leu Tyr Ala Ala Ile Phe Leu Val Gly Val Pro  
 35 40 45  
 Gly Asn Ala Met Val Ala Trp Val Ala Gly Lys Val Ala Arg Arg Arg  
 50 55 60  
 Val Gly Ala Thr Trp Leu Leu His Leu Ala Val Ala Asp Leu Leu Cys  
 65 70 75 80  
 Cys Leu Ser Leu Pro Ile Leu Ala Val Pro Ile Ala Arg Gly Gly His  
 85 90 95  
 Pro Tyr Gly Ala Val Gly Cys Arg Ala Leu Pro Ser Ile Ile Leu Leu  
 100 105 110  
 Thr Met Tyr Ala Ser Val Leu Leu Leu Ala Ala Leu Ser Ala Asp Leu  
 115 120 125  
 Cys Phe Leu Ala Leu Gly Pro Ala Trp Trp Tyr Thr Val Gln Arg Ala  
 130 135 140  
 Cys Gly Val Gln Val Ala Cys Gly Ala Ala Trp Thr Leu Ala Leu Leu  
 145 150 155 160  
 Leu Thr Val Pro Ser Ala Ile Tyr Arg Arg Leu His Gln Glu His Phe  
 165 170 175

Pro

&lt;210&gt; 95

&lt;211&gt; 177

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 95

Ala Pro Leu Ala Phe Arg Gln Gly Ala Leu Gln Ala Asp Gly Val Cys  
 1 5 10 15  
 Ala Leu His Arg His Leu Arg Gly Val Tyr Leu Met Ala Cys Val Ser  
 20 25 30  
 Val Asp His Tyr Pro Ala Val Val Cys Ala His Trp Gly Pro Cys Leu  
 35 40 45  
 Arg Thr Ala Gly Arg Ala Arg Leu Val Cys Val Ala Ile Trp Thr Leu  
 50 55 60  
 Val Leu Leu Gln Thr Met Pro Leu Leu Leu Met Pro Met Thr Lys Pro  
 65 70 75 80

Leu Val Gly Lys Leu Ala Cys Met Glu Tyr Ser Ser Met Glu Ser Val  
                             85                            90                            95  
 Leu Gly Ala Ala Pro His Gly Pro Gly Gly Leu Cys His Trp Leu Leu  
                             100                            105                            110  
 Trp Ala Ser Gly Asp His Pro Val Leu Leu Tyr Glu Asp His Leu Glu  
                             115                            120                            125  
 Ala Val Gln His Ser Trp Glu Asn Pro Val Thr Ser Gly Lys Gly His  
                             130                            135                            140  
 His Arg Arg Gly Ser Pro Gly Gly Pro Ser Asp Gln Gln Glu Arg Thr  
                             145                            150                            155                            160  
 Pro Pro Ala Gly Gln Pro Arg Arg Thr Gln Pro Ala Gly Lys Asp Thr  
                             165                            170                            175

Thr

<210> 96  
 <211> 358  
 <212> PRT  
 <213> Homo sapiens

<400> 96

Met Ser Val Cys Tyr Arg Pro Pro Gly Asn Glu Thr Leu Leu Ser Trp  
 1                            5                            10                            15  
 Lys Thr Ser Arg Ala Thr Gly Thr Ala Phe Leu Leu Leu Ala Ala Leu  
                             20                            25                            30  
 Leu Gly Leu Pro Gly Asn Gly Phe Val Val Trp Ser Leu Ala Gly Trp  
                             35                            40                            45  
 Arg Pro Ala Arg Gly Arg Pro Leu Ala Ala Thr Leu Val Leu His Leu  
                             50                            55                            60  
 Ala Leu Ala Asp Gly Ala Val Leu Leu Leu Thr Pro Leu Phe Val Ala  
                             65                            70                            75                            80  
 Phe Leu Thr Arg Gln Ala Trp Pro Leu Gly Gln Ala Gly Cys Lys Ala  
                             85                            90                            95  
 Val Tyr Tyr Val Cys Ala Leu Ser Met Tyr Ala Ser Val Leu Leu Thr  
                             100                            105                            110  
 Gly Leu Leu Ser Leu Gln Arg Cys Leu Ala Val Thr Arg Pro Phe Leu  
                             115                            120                            125  
 Ala Pro Arg Leu Arg Ser Pro Ala Leu Ala Arg Arg Leu Leu Leu Ala  
                             130                            135                            140  
 Val Trp Leu Ala Ala Leu Leu Leu Ala Val Pro Ala Ala Val Tyr Arg  
                             145                            150                            155                            160

His Leu Trp Arg Asp Arg Val Cys Gln Leu Cys His Pro Ser Pro Val  
 165 170 175  
 His Ala Ala Ala His Leu Ser Leu Glu Thr Leu Thr Ala Phe Val Leu  
 180 185 190  
 Pro Phe Gly Leu Met Leu Gly Cys Tyr Ser Val Thr Leu Ala Arg Leu  
 195 200 205  
 Arg Gly Ala Arg Trp Gly Ser Gly Arg His Gly Ala Arg Val Gly Arg  
 210 215 220  
 Leu Val Ser Ala Ile Val Leu Ala Phe Gly Leu Leu Trp Ala Pro Tyr  
 225 230 235 240  
 His Ala Val Asn Leu Leu Gln Ala Val Ala Ala Leu Ala Pro Pro Glu  
 245 250 255  
 Gly Ala Leu Ala Lys Leu Gly Gly Ala Gly Gln Ala Ala Arg Ala Gly  
 260 265 270  
 Thr Thr Ala Leu Ala Phe Phe Ser Ser Ser Val Asn Pro Val Leu Tyr  
 275 280 285  
 Val Phe Thr Ala Gly Asp Leu Leu Pro Arg Ala Gly Pro Arg Phe Leu  
 290 295 300  
 Thr Arg Leu Phe Glu Gly Ser Gly Glu Ala Arg Gly Gly Gly Arg Ser  
 305 310 315 320  
 Arg Glu Gly Thr Met Glu Leu Arg Thr Thr Pro Gln Leu Lys Val Val  
 325 330 335  
 Gly Gln Gly Arg Gly Asn Gly Asp Pro Gly Gly Gly Met Glu Lys Asp  
 340 345 350  
 Gly Pro Glu Trp Asp Leu  
 355

<210> 97  
 <211> 211  
 <212> PRT  
 <213> Homo sapiens

<400> 97

Met Pro Leu Pro Val Pro Pro Ala Gly Ala Gln Lys Thr Pro Glu Asp  
 1 5 10 15  
 His Val Cys Leu His Leu Ala Gly Pro Ser Pro Ala Pro Ser Glu Pro  
 20 25 30  
 Ala Arg Met Phe Gly Leu Phe Gly Leu Trp Arg Thr Phe Asp Ser Val  
 35 40 45  
 Val Phe Tyr Leu Thr Leu Ile Val Gly Leu Gly Gly Pro Val Gly Asn  
 50 55 60  
 Gly Leu Val Leu Trp Asn Leu Gly Phe Arg Ile Lys Lys Gly Pro Phe  
 68

65					70					75					80
Ser	Ile	Tyr	Leu	Leu	His	Leu	Ala	Ala	Ala	Asp	Phe	Leu	Phe	Leu	Ser
				85					90					95	
Cys	Arg	Val	Gly	Phe	Ser	Val	Ala	Gln	Ala	Ala	Leu	Gly	Ala	Gln	Asp
			100					105					110		
Thr	Leu	Tyr	Phe	Val	Leu	Thr	Phe	Leu	Trp	Phe	Ala	Val	Gly	Leu	Trp
		115					120					125			
Leu	Leu	Ala	Ala	Phe	Ser	Val	Glu	Arg	Cys	Leu	Ser	Asp	Leu	Phe	Pro
		130				135					140				
Ala	Cys	Tyr	Gln	Gly	Cys	Arg	Pro	Arg	His	Ala	Ser	Ala	Val	Leu	Cys
145					150					155					160
Ala	Leu	Val	Trp	Thr	Pro	Thr	Leu	Pro	Ala	Val	Pro	Leu	Pro	Ala	Asn
				165					170					175	
Ala	Cys	Gly	Leu	Leu	Arg	Asn	Ser	Ala	Cys	Pro	Leu	Val	Cys	Pro	Arg
			180					185					190		
Tyr	His	Val	Ala	Ser	Val	Thr	Trp	Phe	Leu	Val	Leu	Ala	Arg	Val	Ala
		195					200					205			
Trp	Thr	Ala													
		210													
<210>	98														
<211>	187														
<212>	PRT														
<213>	Homo sapiens														
<400>	98														
Val	Tyr	Arg	Asn	Pro	Phe	Ala	Ile	Tyr	Leu	Leu	Val	Arg	Gly	Leu	Gln
1				5					10					15	
Gln	Asp	Leu	Ile	Phe	Leu	Gly	Cys	His	Met	Val	Ala	Ile	Val	Pro	Asp
			20					25					30		
Leu	Leu	Gln	Gly	Arg	Leu	Asp	Phe	Pro	Gly	Phe	Val	Gln	Thr	Ser	Leu
		35					40					45			
Ala	Thr	Leu	Arg	Phe	Phe	Cys	Tyr	Ile	Val	Gly	Leu	Ser	Leu	Leu	Ala
		50				55					60				
Ala	Val	Ser	Val	Glu	Gln	Cys	Leu	Ala	Ala	Leu	Phe	Pro	Ala	Trp	Tyr
65					70					75					80
Ser	Cys	Arg	Arg	Pro	Arg	His	Leu	Thr	Thr	Cys	Val	Cys	Ala	Leu	Thr
				85					90					95	
Trp	Ala	Leu	Cys	Leu	Leu	Leu	His	Leu	Leu	Leu	Ser	Ser	Ala	Cys	Thr
			100					105					110		
Gln	Phe	Phe	Gly	Glu	Pro	Ser	Arg	His	Leu	Cys	Arg	Thr	Leu	Trp	Leu
			115				120					125			

Val Ala Ala Val Leu Leu Ala Leu Leu Cys Cys Thr Met Cys Gly Ala  
 130 135 140  
 Ser Leu Met Leu Leu Leu Arg Val Glu Arg Gly Pro Gln Arg Pro Pro  
 145 150 155 160  
 Pro Arg Gly Phe Pro Gly Leu Ile Leu Pro His Arg Pro Pro Leu Pro  
 165 170 175  
 Leu Leu Arg Pro Ala Leu Arg His Leu Leu Ala  
 180 185

<210> 99  
 <211> 216  
 <212> PRT  
 <213> Homo sapiens

<400> 99

Gly Ser Arg Asn Thr Leu Pro His Asn Phe Tyr Gly Cys Leu Tyr Pro  
 1 5 10 15  
 Val Tyr Leu Asn Val Ser Ser Ala Ala Ile Pro Val Tyr Gln Asn  
 20 25 30  
 Arg Glu Val Met Lys Leu Thr Lys Met Val Leu Val Leu Val Val Val  
 35 40 45  
 Phe Ile Leu Ser Ala Ala Pro Tyr His Val Ile Gln Leu Val Asn Leu  
 50 55 60  
 Gln Met Glu Gln Pro Thr Leu Ala Phe Tyr Val Gly Tyr Tyr Leu Ser  
 65 70 75 80  
 Ile Cys Leu Ser Tyr Ala Ser Ser Ser Ile Asn Pro Phe Leu Tyr Ile  
 85 90 95  
 Leu Leu Ser Gly Asn Phe Gln Lys Arg Leu Pro Gln Ile Gln Arg Arg  
 100 105 110  
 Ala Thr Glu Lys Glu Ile Asn Asn Met Gly Asn Thr Leu Lys Ser His  
 115 120 125  
 Phe Glu Ser Thr Trp Ile Thr Met Ser Leu Asp Met Ile Val Tyr Leu  
 130 135 140  
 Thr Gly Ile Ile Arg Lys Gly Arg Cys Thr Asp Met Phe Met Pro Ile  
 145 150 155 160  
 Leu Leu Val Tyr Leu Leu Leu Ala Ala Trp Lys Arg Ser Val Thr Met  
 165 170 175  
 Gln Ile Gln Ala Tyr Ala Asn Phe Ser Lys Met Asn Val Asp Leu Tyr  
 180 185 190  
 Cys Gly Gly Met Gly Ser Glu Ile Pro Arg Leu His Asp Gly Val Tyr  
 195 200 205



Tyr Phe Ser Ile Leu Thr Ser His  
210 215

<210> 100  
<211> 181  
<212> PRT  
<213> Homo sapiens

<400> 100

Gln Phe Ser Ile Ala Ser Leu Ala Cys Ala Asp Phe Leu Val Gly Val  
1 5 10 15

Thr Val Met Leu Phe Ser Met Val Arg Thr Val Glu Ser Cys Trp Tyr  
20 25 30

Phe Gly Ala Lys Phe Cys Thr Leu His Ser Cys Cys Asp Val Ala Phe  
35 40 45

Cys Tyr Ser Ser Val Leu His Leu Cys Phe Ile Cys Ile Asp Arg Tyr  
50 55 60

Ile Val Val Thr Asp Pro Leu Val Tyr Ala Thr Lys Phe Thr Val Ser  
65 70 75 80

Val Ser Gly Ile Cys Ile Ser Val Ser Trp Ile Leu Pro Leu Thr Tyr  
85 90 95

Ser Gly Ala Val Phe Tyr Thr Gly Val Asn Asp Asp Gly Leu Glu Glu  
100 105 110

Leu Val Ser Ala Leu Asn Cys Val Gly Gly Cys Gln Ile Ile Val Ser  
115 120 125

Gln Gly Trp Val Leu Ile Asp Phe Leu Leu Phe Phe Ile Pro Thr Leu  
130 135 140

Val Met Ile Ile Leu Tyr Ser Lys Ile Phe Leu Ile Ala Lys Gln Gln  
145 150 155 160

Ala Ile Lys Ile Glu Thr Thr Ser Ser Lys Val Glu Ser Ser Ser Glu  
165 170 175

Ser Tyr Lys Ile Arg  
180

<210> 101  
<211> 188  
<212> PRT  
<213> Homo sapiens

<400> 101

Tyr Ser Phe Leu Thr Leu Asn Ile His Ala Pro Ala Val Cys Lys Ala  
1 5 10 15

Leu Gly Thr His Thr Lys Asp Glu Asn Gln Gly Val Arg Gln His Val  
20 25 30

Asp Ser Ser Cys Leu Val Glu Arg Leu Gln Ser Glu Ser Pro Tyr Ser  
           35                          40                          45  
 His Ser Val Thr Tyr Arg Gly Glu Arg Lys Trp Gly Glu His Arg Glu  
       50                          55                          60  
 Gly Ile Ser Ile Gly Ser Gly Leu Pro Pro Phe Phe Ser Pro Pro His  
   65                          70                          75                          80  
 Gly Gly Lys Tyr Val Leu Ser Leu Pro Gly Trp Val Met Phe Thr Ala  
                           85                          90                          95  
 Pro Phe Ser Ile Pro Ile Gly Thr Ser Arg Leu Ser Pro Lys Leu Asp  
                          100                         105                         110  
 Leu Gln Ile Thr Ile Leu Arg Lys Leu Asp Val Leu Gln Gln Val Thr  
                          115                         120                         125  
 Gln Glu Leu Ser Leu Ser His Leu Phe Val His Pro Ile Ile Tyr Ser  
      130                         135                         140  
 Phe Ile His Ser Thr Ser Ile Ser Gln Ala Pro Ile Pro Ser Ser Thr  
   145                         150                         155                         160  
 Lys Asp Ser Ile Thr Gly Gln Met Lys Asp Val Thr Ile His Leu Glu  
                          165                         170                         175  
 Lys Phe Ser Leu Ser Trp Ser Glu Cys Thr Gln Arg  
                          180                         185

&lt;210&gt; 102

&lt;211&gt; 204

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 102

Asn Asp Lys Asp Met Arg Met Ser Leu Pro Tyr Ser Leu Asp Ile Cys  
   1                          5                         10                         15  
 Pro Leu Gln Ile Leu Cys Asn Val Ile Pro Asn Val Val Ser Glu Thr  
                          20                         25                         30  
 Trp Trp Lys Val Val Tyr Tyr Glu Gly Gly Ser Leu Met Asn Asp Leu  
      35                         40                         45  
 Ala Pro Ser Ala Trp Met Ser Ser His Glu Ile Val His Thr Arg  
      50                         55                         60  
 Ser Gly Cys Leu Lys Val Asp Leu Leu Ser Leu Ser Leu Ser Leu Ala  
   65                         70                         75                         80  
 Pro Phe His Thr Met Tyr Ser Ser Phe Pro Phe Pro Phe Cys His Tyr  
                          85                         90                         95  
 Lys Leu Ser Glu Ala Pro Thr Arg Ser Gln Ala Asp Val Val His Ser  
                          100                         105                         110  
 Leu Asn His Glu Ser Asn Lys Leu Leu Phe Phe Lys Leu Pro Ser Phe

115                      120                      125  
 Arg Tyr Phe Phe Ile Pro Met Gln Lys Trp Pro Ser Thr Tyr Pro Asp  
     130                      135                      140  
 Leu Ile Ser Ile Gln His Ile Tyr Lys Leu Met Tyr Gln Ile Ile Leu  
     145                      150                      155                      160  
 His Lys Tyr Ile Lys Leu Gln Cys Val Asn Leu Lys Arg Ile Thr Tyr  
                     165                      170                      175  
 Leu Ile Tyr Lys Phe Phe Ile Pro Phe Asn Ser Ile Ser Val Phe Phe  
                     180                      185                      190  
 Thr Ile Phe Asp Tyr Phe Leu Ile Phe Leu Ile Pro  
                     195                      200  
 <210> 103  
 <211> 215  
 <212> PRT  
 <213> Homo sapiens  
 <400> 103  
 Met Arg Ala Phe Trp Lys Lys Thr Val Asn Ser Thr Val Gly Pro Leu  
     1                      5                      10                      15  
 Lys Lys Ser Ile His Arg Leu Asn Pro Leu Val Asp Asn Lys Thr Leu  
                     20                      25                      30  
 Asn Ile Tyr His Ala Phe Val Ile Ile Gln Ile Val Asp Ile Phe Thr  
                     35                      40                      45  
 Arg Ala Thr Asp Glu Leu Gln Pro Ser Phe Tyr Gln Leu His Arg Ile  
                     50                      55                      60  
 Leu Leu Asn Gly Tyr Ile Leu Leu Met Ser Val Leu Ser Ser Glu Tyr  
     65                      70                      75                      80  
 Val Val Ser Asn Phe Ser Ile Leu Glu Thr Met Leu Lys Glu Leu Ala  
                     85                      90                      95  
 Leu Ile Cys Val Ser Tyr Val Tyr Thr Ser Ile Gln Tyr Leu Ser Arg  
                     100                      105                      110  
 Ser Gly Asn Ser Trp Leu Lys Gln Thr Ala Phe Gln Glu Asp Leu Leu  
                     115                      120                      125  
 Ile Tyr Ile Pro Thr Asn Ala Val Leu Glu Cys Leu Phe Leu His Ile  
                     130                      135                      140  
 Pro Met Asn Thr Glu Cys Ser Leu Phe Asn Thr Phe Phe Pro Gln Ser  
     145                      150                      155                      160  
 Thr Arg Leu Leu Leu Lys Tyr Ile Leu Phe Lys Thr Pro Ala Tyr Pro  
                     165                      170                      175  
 Ser Ile Tyr Cys Leu Leu Lys Asp Tyr Leu Val Leu Phe Met Cys Val  
                     180                      185                      190

Gly Gly Lys Leu Ile Val Ile Leu Gln Lys Leu Gln Leu Leu Leu His  
 195 200 205

Gln Pro Asn Ile Ile His Val  
 210 215

<210> 104  
 <211> 194  
 <212> PRT  
 <213> Homo sapiens

<400> 104

Leu Phe Trp Phe Pro Phe Phe Leu Ile Thr Tyr Leu Phe Ser Phe Ala  
 1 5 10 15

Gly Pro Ser His Leu Pro Gly Leu Phe Phe Val Cys Ile Tyr Leu Cys  
 20 25 30

Gly Leu Pro Ser Leu Thr His Leu Pro Ile Ser Tyr Ser Gln Met Tyr  
 35 40 45

Ile Ser Ile Pro Gly Pro Phe Leu Thr Pro Asp Leu Tyr Phe Gln Leu  
 50 55 60

Pro Thr Gln Tyr Leu Leu Gly Tyr Leu Leu Lys Phe Glu Ser Arg His  
 65 70 75 80

Ala Asn Thr Ser Tyr Leu Ser Lys Ala Leu Ser Phe Tyr Cys Phe Ser  
 85 90 95

His Pro Asn Lys Leu Pro Leu Ile Leu Pro Asp Ala Trp Ala Thr Phe  
 100 105 110

Leu Gln Ser Ser Leu Ile Ser Phe Phe Leu Tyr Pro Thr Tyr Ile His  
 115 120 125

Gln His Met Leu Pro Cys Leu His Phe Lys His Asn Gln His Pro Thr  
 130 135 140

Phe Leu Gln Leu Leu Leu Ala Leu Tyr Ser Pro Ile Leu Ser Pro Leu  
 145 150 155 160

Asp Tyr Cys Ile Ser Leu Thr Gly Ile Pro Leu Gln Pro Phe Pro Thr  
 165 170 175

Ser Ser Ser Gln Ser Gln Leu Cys Glu Gln Gln Leu Lys Leu Phe Phe  
 180 185 190

Gln Lys

<210> 105  
 <211> 186  
 <212> PRT  
 <213> Homo sapiens

<400> 105

Asn Pro Gly Lys Ala Ser His Leu Gly Leu Cys Thr Ser Gly Leu Phe  
 1 5 10 15  
 Asp Ala Leu Gly Asn Val Glu Gly His Pro Val Ser Arg Trp Gly Leu  
 20 25 30  
 Glu Gln Ser Leu Asp Cys Phe Ser Gln Trp Leu Leu Thr Ser Gly Cys  
 35 40 45  
 Cys Ile Pro Ser Thr Phe Trp Leu Val Leu Arg Thr Thr Asn Lys Lys  
 50 55 60  
 Val Gly Arg Thr Val Leu His His Leu Cys Lys Leu Leu Gly Lys Gln  
 65 70 75 80  
 Thr Asn Val Leu Gln Lys Glu Asp Glu Leu Leu Lys His Lys Gly Gly  
 85 90 95  
 Met Leu His Arg Glu Gly Leu Glu Ser Trp Ile Thr Lys Arg Asp Lys  
 100 105 110  
 Asp Thr Phe Gly Arg Asp Gly Tyr Val Tyr Tyr Leu Ala Tyr Gly Asp  
 115 120 125  
 Ser Phe Ile Gly Pro Ile Pro Lys Ala Ser His Cys Thr Leu Thr Met  
 130 135 140  
 Tyr Asp Leu Phe Ile Leu Ile Ile Pro Gln Ser Cys Phe Leu Lys Lys  
 145 150 155 160  
 Leu Pro Leu Asn Pro Val Asn Arg Pro Gly Arg Gln Leu Ile Asn Ile  
 165 170 175  
 Phe Phe Thr Phe Glu Lys Leu Lys Leu Ser  
 180 185

<210> 106  
 <211> 184  
 <212> PRT  
 <213> Homo sapiens

<400> 106

Leu Gly Phe Cys His Leu Leu Val Glu Trp Arg Ala Cys His Ser Val  
 1 5 10 15  
 Cys Leu Ser Leu Phe Pro Tyr Leu Ser Gly Asp Asn Asn Asn Met Tyr  
 20 25 30  
 Ile Ile Glu Leu Leu Ser Ser Ser Cys Lys Ser Ile Leu Thr Lys Phe  
 35 40 45  
 Leu Glu Asn Ala Tyr Ser Lys His Ser Ile Thr Tyr Ala Ile Cys Ile  
 50 55 60  
 Ser Ile Asn Arg Tyr Ile Leu Val Val Tyr Pro Glu Thr Phe Leu Val  
 65 70 75 80

Cys Ser Leu Leu Pro Phe Phe Phe Pro Glu Lys Thr His Arg Phe Cys  
                     85                    90                    95  
 Leu Met His Gly Lys Glu Lys Tyr His Gln Val Leu Gly Ser Ser Lys  
                     100                    105                    110  
 Lys Ile Lys Lys Pro Lys Thr Cys Thr Leu Glu Arg Gly Lys Leu Ile  
                     115                    120                    125  
 Pro Met Glu Lys Lys Lys Lys Arg Asn Leu Asn Asn Cys Ser Ser Glu  
                     130                    135                    140  
 Gly His Val Gly Leu Gln Arg Gly Phe His Met Pro Phe Leu Ser Arg  
                     145                    150                    155                    160  
 Gly Asn His Cys Pro Asp Gln Phe Ser Lys Glu Gly Lys Val Lys Phe  
                     165                    170                    175  
 His Arg Glu Gly Trp Ser Ile Asn  
                     180

<210> 107  
 <211> 170  
 <212> PRT  
 <213> Homo sapiens

<400> 107

Val Ala Gly Cys Thr Asn Phe Thr His Lys Gln Leu Glu Val His Phe  
 1                    5                    10                    15  
 Pro Ser Leu Thr Thr Lys Leu His Leu Ile Thr Phe Glu Ile Phe Ala  
                     20                    25                    30  
 Ile Leu Leu Gly Leu Thr Cys Leu Ser Leu Phe Tyr Ile Cys Ile Ser  
                     35                    40                    45  
 Lys Leu Leu Ser Leu Asn Asn Phe His Met Gly His Leu Tyr Leu Gln  
                     50                    55                    60  
 Asn Lys His Tyr Pro Phe Asn Asp Phe Ala Trp Leu Leu Pro Ser Leu  
                     65                    70                    75                    80  
 Val Phe Ile Ala Ser Leu Lys His Val Asn Ser Phe Ile Cys Ser Phe  
                     85                    90                    95  
 Val Ser Leu Leu Lys His Phe Ser Asn Ser Thr Thr Ser Phe Tyr Ser  
                     100                    105                    110  
 Phe Gln Phe Asn Ala His Ile Gly His Cys Ala Tyr Leu Ser Lys Leu  
                     115                    120                    125  
 Cys Thr Lys Gln Val Asn Leu Pro Cys Pro Trp Ile Glu Gln Asn Leu  
                     130                    135                    140  
 Lys Lys Ala Pro Gly Glu Asn His Gly Tyr Trp Gln Arg Asp Met Asp  
                     145                    150                    155                    160  
 Ser Asn Pro Gly Phe Ser Thr Tyr Ser Leu

165

170

<210> 108  
 <211> 239  
 <212> PRT  
 <213> Homo sapiens

<400> 108

Thr	Arg	Asn	Thr	Leu	Gly	Cys	Pro	Ser	Ala	Pro	Gly	Phe	Phe	Cys	Trp
1				5					10					15	
Ser	Cys	Leu	Ala	Met	Cys	Leu	Gly	Leu	Lys	Val	Ser	Arg	Leu	Pro	Gly
		20					25						30		
Ser	Pro	Gly	Ser	Ser	Arg	Lys	Arg	Asn	Glu	His	Met	Met	Val	Thr	Trp
		35					40					45			
Asn	Ser	Pro	Arg	Trp	Arg	His	Cys	Ile	Phe	Ala	Lys	Pro	Val	Thr	Val
	50					55					60				
Leu	Ser	Ala	Phe	Trp	Ala	Pro	Arg	Leu	Ser	Pro	Leu	Ile	Phe	Pro	Asp
65					70					75					80
Leu	Ser	Phe	Pro	Ala	Ala	Phe	Leu	Phe	Phe	Leu	Ile	Thr	Val	Lys	Phe
				85					90					95	
Cys	Met	Tyr	Cys	Ser	Ile	Phe	His	Leu	Leu	Gly	Ile	Glu	Tyr	Ile	Ser
			100					105					110		
Ser	Met	Pro	Gly	Phe	Lys	Ile	Arg	Ile	Val	Asn	Ile	Val	Val	Cys	Ala
		115					120					125			
Leu	Val	Thr	Glu	Phe	Leu	Arg	Phe	Gly	Cys	Ser	Ile	Pro	Ala	Pro	Tyr
		130				135					140				
Phe	Leu	Lys	Ala	Leu	Leu	Ser	Ala	Val	Gly	Asp	Phe	Ala	Gln	Cys	Lys
145					150					155					160
Leu	Leu	Arg	Tyr	Phe	Leu	Leu	Ser	Ser	Arg	Ser	Pro	Tyr	Pro	Thr	Ser
				165					170					175	
Thr	Gln	His	Leu	Ile	Leu	Arg	Cys	Ser	Pro	Gln	Thr	Cys	Glu	Asn	Gln
			180					185					190		
His	Val	Asn	Met	Ser	Ile	Pro	Leu	Ala	Gly	Phe	Pro	Asn	Ser	Thr	Asp
		195					200					205			
Gly	Ile	Arg	Pro	Ile	Val	Gln	Ala	Lys	Ser	Lys	Ala	Pro	Ala	Gly	Thr
	210					215					220				
Phe	Pro	Ile	Pro	Asn	Leu	Ser	Ser	Cys	Pro	Ile	Lys	Phe	Tyr	Gln	
225					230					235					

<210> 109  
 <211> 202  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 109

Lys Phe Tyr Leu His Thr Lys Gln Gly Arg Ile Thr Ser Tyr Cys Leu  
 1 5 10 15  
 Ala His Arg Lys Glu Ala Phe Cys Ser Asp Ile Ile Tyr Thr Leu Arg  
 20 25 30  
 Asn Lys Gly Val Ala Lys Ser Phe Ser Ser Cys Lys His Ser Thr Ile  
 35 40 45  
 Leu Gly Leu Thr Ile Tyr Ser Thr Leu Lys Ala Ala Phe Leu Glu Cys  
 50 55 60  
 Ile Ile Ser Val Leu Phe Leu Leu Ile Phe Phe Tyr Leu Ser Trp Phe  
 65 70 75 80  
 Pro Pro Ser Thr Val Leu Thr Ser Val Tyr Lys Asn Leu Tyr Ser Ser  
 85 90 95  
 Pro His Ile Pro Tyr Leu Ile Cys Val Thr Ile Lys Ala Ile Cys Leu  
 100 105 110  
 Asp Thr Leu Gln Lys Cys Ile Gln Leu Ile Ser Asp Phe Ile Ser Val  
 115 120 125  
 Arg Ala Asn Asn Gln Phe Ile Gln Leu Phe Phe Pro Ser Glu Ser Thr  
 130 135 140  
 Glu Tyr Pro Leu Asn Glu Val Phe Ser Val Glu Phe Ser Leu Lys Leu  
 145 150 155 160  
 Ser Arg Asn Glu His Ser Pro Lys Cys Phe Val Gly Ala Ser Cys Ala  
 165 170 175  
 Arg Val Gly Val Arg Phe Cys His Leu Pro Val Ile Ser Ser Leu Asn  
 180 185 190  
 Ile Leu Ser Leu Met Arg Ser Pro Leu Lys  
 195 200

&lt;210&gt; 110

&lt;211&gt; 192

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

Tyr Gly Ser Ala Val Ser Pro Ser Lys Ser Asn Leu Glu Leu Gln Leu  
 1 5 10 15  
 Pro Phe Pro Cys Val Val Glu Gly Ala Trp Trp Glu Ile Thr Glu Ser  
 20 25 30  
 Trp Gly Arg Leu Pro Pro Tyr Cys Ser Leu Gly Ser Glu Val Ser Asp  
 35 40 45  
 Leu Ile Val Leu Gly Glu Thr Pro Phe Ala Trp Phe Leu Phe Ser Leu  
 50 55 60



Cys Trp Leu Pro Cys Lys Met Leu Leu Cys Ser Ser Phe Val Phe Cys  
 65 70 75 80  
 His Asp Cys Gly Ala Ser Pro Ala Met Trp Lys Cys Glu Glu Ile Lys  
 85 90 95  
 Pro Leu Ser Phe Ile Asn Cys Pro Val Leu Gly Met Ser Leu Ser Ala  
 100 105 110  
 Val Asn Gly Leu Ile His Gln Lys Phe Lys Thr Ser Leu Gly Asn Thr  
 115 120 125  
 Val Arg Pro Cys Leu Tyr Lys Glu Arg Ile Arg Lys Arg Lys Ile Ser  
 130 135 140  
 Val Ala Tyr Thr Glu Tyr Tyr Lys Thr Asp Gly Thr Leu Ile Pro Glu  
 145 150 155 160  
 Leu Phe Leu Tyr His Phe Glu Arg Pro Asp Asn Gln Tyr Asn Phe Leu  
 165 170 175  
 Ser Arg Val Ala Gln Thr Cys Lys His Ala Leu Pro Tyr Leu Asp Asn  
 180 185 190

<210> 111  
 <211> 311  
 <212> PRT  
 <213> Homo sapiens

<400> 111

Met Met Glu Pro Arg Glu Ala Gly Gln His Val Gly Ala Ala Asn Gly  
 1 5 10 15  
 Ala Gln Glu Asp Val Ala Phe Asn Leu Ile Ile Leu Ser Leu Thr Glu  
 20 25 30  
 Gly Leu Gly Leu Gly Gly Leu Leu Gly Asn Gly Ala Val Leu Trp Leu  
 35 40 45  
 Leu Ser Ser Asn Val Tyr Arg Asn Pro Phe Ala Ile Tyr Leu Leu Asp  
 50 55 60  
 Val Ala Cys Ala Asp Leu Ile Phe Leu Gly Cys His Met Val Ala Ile  
 65 70 75 80  
 Val Pro Asp Leu Leu Gln Gly Arg Leu Asp Phe Pro Gly Phe Val Gln  
 85 90 95  
 Thr Ser Leu Ala Thr Leu Arg Phe Cys Tyr Ile Val Gly Leu Ser Leu  
 100 105 110  
 Leu Ala Ala Val Ser Val Glu Gln Cys Leu Ala Ala Leu Phe Pro Ala  
 115 120 125  
 Trp Tyr Ser Cys Arg Arg Pro Arg His Leu Thr Thr Cys Val Cys Ala  
 130 135 140

Leu Thr Trp Ala Leu Cys Leu Leu Leu His Leu Leu Leu Ser Gly Ala  
 145 150 155 160  
 Cys Thr Gln Phe Phe Gly Glu Pro Ser Arg His Leu Cys Arg Thr Leu  
 165 170 175  
 Trp Leu Val Ala Ala Val Leu Leu Ala Leu Leu Cys Cys Thr Met Cys  
 180 185 190  
 Gly Ala Ser Leu Met Leu Leu Leu Arg Val Glu Arg Gly Pro Gln Arg  
 195 200 205  
 Pro Pro Pro Arg Gly Phe Pro Gly Leu Ile Leu Leu Thr Val Leu Leu  
 210 215 220  
 Phe Leu Phe Cys Gly Leu Pro Phe Gly Ile Tyr Trp Leu Ser Arg Asn  
 225 230 235 240  
 Leu Leu Trp Tyr Ile Pro His Tyr Phe Tyr His Phe Ser Phe Leu Met  
 245 250 255  
 Ala Ala Val His Cys Ala Ala Lys Pro Val Val Tyr Phe Cys Leu Gly  
 260 265 270  
 Ser Ala Gln Gly Arg Arg Leu Pro Leu Arg Leu Val Leu Gln Arg Ala  
 275 280 285  
 Leu Gly Asp Glu Ala Glu Leu Gly Ala Val Arg Glu Thr Ser Arg Arg  
 290 295 300  
 Gly Leu Val Asp Ile Ala Ala  
 305 310

<210> 112  
 <211> 508  
 <212> PRT  
 <213> Homo sapiens

<400> 112

Met Thr Ser Thr Cys Thr Asn Ser Thr Arg Glu Ser Asn Ser Ser His  
 1 5 10 15  
 Thr Cys Met Pro Leu Ser Lys Met Pro Ile Ser Leu Ala His Gly Ile  
 20 25 30  
 Ile Arg Ser Thr Val Leu Val Ile Phe Leu Ala Ala Ser Phe Val Gly  
 35 40 45  
 Asn Ile Val Leu Ala Leu Val Leu Gln Arg Lys Pro Gln Leu Leu Gln  
 50 55 60  
 Val Thr Asn Arg Phe Ile Phe Asn Leu Leu Val Thr Asp Leu Leu Gln  
 65 70 75 80  
 Ile Ser Leu Val Ala Pro Trp Val Val Ala Thr Ser Val Pro Leu Phe  
 85 90 95  
 Trp Pro Leu Asn Ser His Phe Cys Thr Ala Leu Val Ser Leu Thr His  
 80

100					105					110					
Leu	Phe	Ala	Phe	Ala	Ser	Val	Asn	Thr	Ile	Val	Val	Val	Ser	Val	Asp
		115					120					125			
Arg	Tyr	Leu	Ser	Ile	Ile	His	Pro	Leu	Ser	Tyr	Pro	Ser	Lys	Met	Thr
	130					135					140				
Gln	Arg	Arg	Gly	Tyr	Leu	Leu	Leu	Tyr	Gly	Thr	Trp	Ile	Val	Ala	Ile
145					150					155					160
Leu	Gln	Ser	Thr	Pro	Pro	Leu	Tyr	Gly	Trp	Gly	Gln	Ala	Ala	Phe	Asp
				165					170					175	
Glu	Arg	Asn	Ala	Leu	Cys	Ser	Met	Ile	Trp	Gly	Ala	Ser	Pro	Ser	Tyr
			180					185					190		
Thr	Ile	Leu	Ser	Val	Val	Ser	Phe	Ile	Val	Ile	Pro	Leu	Ile	Val	Met
		195					200					205			
Ile	Ala	Cys	Tyr	Ser	Val	Val	Phe	Cys	Ala	Ala	Arg	Arg	Gln	His	Ala
	210						215				220				
Leu	Leu	Tyr	Asn	Val	Lys	Arg	His	Ser	Leu	Glu	Val	Arg	Val	Lys	Asp
225					230					235					240
Cys	Val	Glu	Asn	Glu	Asp	Glu	Glu	Gly	Ala	Glu	Lys	Lys	Glu	Glu	Phe
				245					250					255	
Gln	Asp	Glu	Ser	Glu	Phe	Arg	Arg	Gln	His	Glu	Gly	Glu	Val	Lys	Ala
			260					265					270		
Lys	Glu	Gly	Arg	Met	Glu	Ala	Lys	Asp	Gly	Ser	Leu	Lys	Ala	Lys	Glu
		275					280					285			
Gly	Ser	Thr	Gly	Thr	Ser	Glu	Ser	Ser	Val	Glu	Ala	Arg	Gly	Ser	Glu
		290				295					300				
Glu	Val	Arg	Glu	Ser	Ser	Thr	Val	Ala	Ser	Asp	Gly	Ser	Met	Glu	Gly
305					310					315					320
Lys	Glu	Gly	Ser	Thr	Lys	Val	Glu	Glu	Asn	Ser	Met	Lys	Ala	Asp	Lys
				325					330					335	
Gly	Arg	Thr	Glu	Val	Asn	Gln	Cys	Ser	Ile	Asp	Leu	Gly	Glu	Asp	Asp
			340					345					350		
Met	Glu	Phe	Gly	Glu	Asp	Asp	Ile	Asn	Phe	Ser	Glu	Asp	Asp	Val	Glu
		355					360					365			
Ala	Val	Asn	Ile	Pro	Glu	Ser	Leu	Pro	Pro	Ser	Arg	Arg	Asn	Ser	Asn
		370				375					380				
Ser	Asn	Pro	Pro	Leu	Pro	Arg	Cys	Tyr	Gln	Cys	Lys	Ala	Ala	Lys	Val
385					390					395					400
Ile	Phe	Ile	Ile	Ile	Phe	Ser	Tyr	Val	Leu	Ser	Leu	Gly	Pro	Tyr	Cys
				405					410					415	

Phe Leu Ala Val Leu Ala Val Trp Val Asp Val Glu Thr Gln Val Pro  
                   420                                  425                                  430  
 Gln Trp Val Ile Thr Ile Ile Ile Trp Leu Phe Phe Leu Gln Cys Cys  
                   435                                  440                                  445  
 Ile His Pro Tyr Val Tyr Gly Tyr Met His Lys Thr Ile Lys Lys Glu  
                   450                                  455                                  460  
 Ile Gln Asp Met Leu Lys Lys Phe Phe Cys Lys Glu Lys Pro Pro Lys  
                   465                                  470                                  475                                  480  
 Glu Asp Ser His Pro Asp Leu Pro Gly Thr Glu Gly Gly Thr Glu Gly  
                                   485                                  490                                  495  
 Lys Ile Val Pro Ser Tyr Asp Ser Ala Thr Phe Pro  
                                   500                                  505

<210> 113  
 <211> 311  
 <212> PRT  
 <213> Homo sapiens

<400> 113

Met Met Glu Pro Arg Glu Ala Gly Gln His Val Gly Ala Ala Asn Ser  
 1                  5                                  10                                  15  
 Ala Gln Glu Asp Val Ala Phe Asn Leu Ile Ile Leu Ser Leu Thr Glu  
                   20                                  25                                  30  
 Gly Leu Gly Leu Gly Gly Leu Leu Gly Asn Gly Ala Val Leu Trp Leu  
                   35                                  40                                  45  
 Leu Ser Ser Asn Val Tyr Arg Asn Pro Phe Ala Ile Tyr Leu Leu Asp  
                   50                                  55                                  60  
 Val Ala Cys Ala Asp Leu Ile Phe Leu Gly Cys His Met Val Ala Ile  
                   65                                  70                                  75                                  80  
 Val Pro Asp Leu Leu Gln Gly Arg Leu Asp Phe Pro Gly Phe Val Gln  
                                   85                                  90                                  95  
 Thr Ser Leu Ala Thr Leu Arg Phe Cys Tyr Ile Val Gly Leu Ser Leu  
                   100                                  105                                  110  
 Leu Ala Ala Val Ser Val Glu Gln Cys Leu Ala Ala Leu Phe Pro Ala  
                   115                                  120                                  125  
 Trp Tyr Ser Cys Arg Arg Pro Arg His Leu Thr Thr Cys Val Cys Ala  
                   130                                  135                                  140  
 Leu Thr Trp Ala Leu Cys Leu Leu Leu His Leu Leu Leu Ser Gly Ala  
                   145                                  150                                  155                                  160  
 Cys Thr Gln Phe Phe Gly Glu Pro Ser Arg His Leu Cys Arg Thr Leu  
                                   165                                  170                                  175  
 Trp Leu Val Ala Ala Val Leu Leu Ala Leu Leu Cys Cys Thr Met Cys

180					185					190					
Gly	Ala	Ser	Leu	Met	Leu	Leu	Leu	Arg	Val	Glu	Arg	Gly	Pro	Gln	Arg
	195						200					205			
Pro	Pro	Pro	Arg	Gly	Phe	Pro	Gly	Leu	Ile	Leu	Leu	Thr	Val	Leu	Leu
	210					215					220				
Phe	Leu	Phe	Cys	Gly	Leu	Pro	Phe	Gly	Ile	Tyr	Trp	Leu	Ser	Arg	Asn
225					230					235					240
Leu	Leu	Trp	Tyr	Ile	Pro	His	Tyr	Phe	Tyr	His	Phe	Ser	Phe	Leu	Met
				245					250					255	
Ala	Ala	Val	His	Cys	Ala	Ala	Lys	Pro	Val	Val	Tyr	Phe	Cys	Leu	Gly
			260					265					270		
Ser	Ala	Gln	Gly	Arg	Arg	Leu	Pro	Leu	Arg	Leu	Val	Leu	Gln	Arg	Ala
	275						280					285			
Leu	Gly	Asp	Glu	Ala	Glu	Leu	Gly	Ala	Val	Arg	Glu	Thr	Ser	Arg	Arg
	290					295					300				
Gly	Leu	Val	Asp	Ile	Ala	Ala									
305					310										
<210> 114															
<211> 333															
<212> PRT															
<213> Homo sapiens															
<400> 114															
Met	Trp	Ser	Cys	Ser	Trp	Phe	Asn	Gly	Thr	Gly	Leu	Val	Glu	Glu	Leu
1				5					10					15	
Pro	Ala	Cys	Gln	Asp	Leu	Gln	Leu	Gly	Leu	Ser	Leu	Leu	Ser	Leu	Leu
			20					25					30		
Gly	Leu	Val	Val	Gly	Val	Pro	Val	Gly	Leu	Cys	Tyr	Asn	Ala	Leu	Leu
		35					40					45			
Val	Leu	Ala	Asn	Leu	His	Ser	Lys	Ala	Ser	Met	Thr	Met	Pro	Asp	Val
	50					55					60				
Tyr	Phe	Val	Asn	Met	Ala	Val	Ala	Gly	Leu	Val	Leu	Ser	Ala	Leu	Ala
65					70				75					80	
Pro	Val	His	Leu	Leu	Gly	Pro	Pro	Ser	Ser	Arg	Trp	Ala	Leu	Trp	Ser
				85					90					95	
Val	Gly	Gly	Glu	Val	His	Val	Ala	Leu	Gln	Ile	Pro	Phe	Asn	Val	Ser
			100					105					110		
Ser	Leu	Val	Ala	Met	Tyr	Ser	Thr	Ala	Leu	Leu	Ser	Leu	Asp	His	Tyr
	115						120					125			
Ile	Glu	Arg	Ala	Leu	Pro	Arg	Thr	Tyr	Met	Ala	Ser	Val	Tyr	Asn	Thr
	130					135					140				

Arg His Val Cys Gly Phe Val Trp Gly Gly Ala Leu Leu Thr Ser Phe  
 145 150 155 160  
 Ser Ser Leu Leu Phe Tyr Ile Cys Ser His Val Ser Thr Arg Ala Leu  
 165 170 175  
 Glu Cys Ala Lys Met Gln Asn Ala Glu Ala Ala Asp Ala Thr Leu Val  
 180 185 190  
 Phe Ile Gly Tyr Val Val Pro Ala Leu Ala Thr Leu Tyr Ala Leu Val  
 195 200 205  
 Leu Leu Ser Arg Val Arg Arg Glu Asp Thr Pro Leu Asp Arg Asp Thr  
 210 215 220  
 Gly Arg Leu Glu Pro Ser Ala His Arg Leu Leu Val Ala Thr Val Cys  
 225 230 235 240  
 Thr Gln Phe Gly Leu Trp Thr Pro His Tyr Leu Ile Leu Leu Gly His  
 245 250 255  
 Thr Val Ile Ile Ser Arg Gly Lys Pro Val Asp Ala His Tyr Leu Gly  
 260 265 270  
 Leu Leu His Phe Val Lys Asp Phe Ser Lys Leu Leu Ala Phe Ser Ser  
 275 280 285  
 Ser Phe Val Thr Pro Leu Leu Tyr Arg Tyr Met Asn Gln Ser Phe Pro  
 290 295 300  
 Ser Lys Leu Gln Arg Leu Met Lys Lys Leu Pro Cys Gly Asp Arg His  
 305 310 315 320  
 Cys Ser Pro Asp His Met Gly Val Gln Gln Val Leu Ala  
 325 330

<210> 115  
 <211> 337  
 <212> PRT  
 <213> Homo sapiens

<400> 115

Met Gly Asn Asp Ser Val Ser Tyr Glu Tyr Gly Asp Tyr Ser Asp Leu  
 1 5 10 15  
 Ser Asp Arg Pro Val Asp Cys Leu Asp Gly Ala Cys Leu Ala Ile Asp  
 20 25 30  
 Pro Leu Arg Val Ala Pro Leu Pro Leu Tyr Ala Ala Ile Phe Leu Val  
 35 40 45  
 Gly Val Pro Gly Asn Ala Met Val Ala Trp Val Ala Gly Lys Val Ala  
 50 55 60  
 Arg Arg Arg Val Gly Ala Thr Trp Leu Leu His Leu Ala Val Ala Asp  
 65 70 75 80

Val

<400> 116

Met Ala Pro Ser His Arg Ala Ser Gln Val Gly Phe Cys Pro Thr Pro  
85

1	5	10	15
Glu Arg Pro Leu Trp Arg Leu Pro Pro Thr Cys Arg Pro Arg Arg Met	20	25	30
Ser Val Cys Tyr Arg Pro Pro Gly Asn Glu Thr Leu Leu Ser Trp Lys	35	40	45
Thr Ser Arg Ala Thr Gly Thr Ala Phe Leu Leu Leu Ala Ala Leu Leu	50	55	60
Gly Leu Pro Gly Asn Gly Phe Val Val Trp Ser Leu Ala Gly Trp Arg	65	70	75
Pro Ala Arg Gly Arg Pro Leu Ala Ala Thr Leu Val Leu His Leu Ala	85	90	95
Leu Ala Asp Gly Ala Val Leu Leu Leu Thr Pro Leu Phe Val Ala Phe	100	105	110
Leu Thr Arg Gln Ala Trp Pro Leu Gly Gln Ala Gly Cys Lys Ala Val	115	120	125
Tyr Tyr Val Cys Ala Leu Ser Met Tyr Ala Ser Val Leu Leu Thr Gly	130	135	140
Leu Leu Ser Leu Gln Arg Cys Leu Ala Val Thr Arg Pro Phe Leu Ala	145	150	155
Pro Arg Leu Arg Ser Pro Ala Leu Ala Arg Arg Leu Leu Leu Ala Val	165	170	175
Trp Leu Ala Ala Leu Leu Leu Ala Val Pro Ala Ala Val Tyr Arg His	180	185	190
Leu Trp Arg Asp Arg Val Cys Gln Leu Cys His Pro Ser Pro Val His	195	200	205
Ala Ala Ala His Leu Ser Leu Glu Thr Leu Thr Ala Phe Val Leu Pro	210	215	220
Phe Gly Leu Met Leu Gly Cys Tyr Ser Val Thr Leu Ala Arg Leu Arg	225	230	235
Gly Ala Arg Trp Gly Ser Gly Arg His Gly Ala Arg Val Gly Arg Leu	245	250	255
Val Ser Ala Ile Val Leu Ala Phe Gly Leu Leu Trp Ala Pro Tyr His	260	265	270
Ala Val Asn Leu Leu Gln Ala Val Ala Ala Leu Ala Pro Pro Glu Gly	275	280	285
Ala Leu Ala Lys Leu Gly Gly Ala Gly Gln Ala Ala Arg Ala Gly Thr	290	295	300
Thr Ala Leu Ala Phe Phe Ser Ser Ser Val Asn Pro Val Leu Tyr Val	305	310	315
			320



Phe Thr Ala Gly Asp Leu Leu Pro Arg Ala Gly Pro Arg Phe Leu Thr  
                                   325                                  330                                  335

Arg Leu Phe Glu Gly Ser Gly Glu Ala Arg Gly Gly Gly Arg Ser Arg  
                                   340                                  345                                  350

Glu Gly Thr Met Glu Leu Arg Thr Thr Pro Gln Leu Lys Val Val Gly  
                                   355                                  360                                  365

Gln Gly Arg Gly Asn Gly Asp Pro Gly Gly Gly Met Glu Lys Asp Gly  
                                   370                                  375                                  380

Pro Glu Trp Asp Leu  
 385

<210> 117  
 <211> 340  
 <212> PRT  
 <213> Homo sapiens

<400> 117

Met Asn Pro Phe His Ala Ser Cys Trp Asn Thr Ser Ala Glu Leu Leu  
 1                                  5                                  10                                  15

Asn Lys Ser Trp Asn Lys Glu Phe Ala Tyr Gln Thr Ala Ser Val Val  
                                   20                                  25                                  30

Asp Thr Val Ile Leu Pro Ser Met Ile Gly Ile Ile Cys Ser Thr Gly  
                                   35                                  40                                  45

Leu Val Gly Asn Ile Leu Ile Val Phe Thr Ile Ile Arg Ser Arg Lys  
                                   50                                  55                                  60

Lys Thr Val Pro Asp Ile Tyr Ile Cys Asn Leu Ala Val Ala Asp Leu  
 65                                  70                                  75                                  80

Val His Ile Val Gly Met Pro Phe Leu Ile His Gln Trp Ala Arg Gly  
                                   85                                  90                                  95

Gly Glu Trp Val Phe Gly Gly Pro Leu Cys Thr Ile Ile Thr Ser Leu  
                                   100                                  105                                  110

Asp Thr Cys Asn Gln Phe Ala Cys Ser Ala Ile Met Thr Val Met Ser  
                                   115                                  120                                  125

Val Asp Arg Tyr Phe Ala Leu Val Gln Pro Phe Arg Leu Thr Arg Trp  
                                   130                                  135                                  140

Arg Thr Arg Tyr Lys Thr Ile Arg Ile Asn Leu Gly Leu Trp Ala Ala  
 145                                  150                                  155                                  160

Ser Phe Ile Leu Ala Leu Pro Val Trp Val Tyr Ser Lys Val Ile Lys  
                                   165                                  170                                  175

Phe Lys Asp Gly Val Glu Ser Cys Ala Phe Asp Leu Thr Ser Pro Asp  
                                   180                                  185                                  190

Asp Val Leu Trp Tyr Thr Leu Tyr Leu Thr Ile Thr Thr Phe Phe Phe

195					200					205					
Pro	Leu	Pro	Leu	Ile	Leu	Val	Cys	Tyr	Ile	Leu	Ile	Leu	Cys	Tyr	Thr
210						215					220				
Trp	Glu	Met	Tyr	Gln	Gln	Asn	Lys	Asp	Ala	Arg	Cys	Cys	Asn	Pro	Ser
225					230					235					240
Val	Pro	Lys	Gln	Arg	Val	Met	Lys	Leu	Thr	Lys	Met	Val	Leu	Val	Leu
				245					250					255	
Val	Val	Val	Phe	Ile	Leu	Ser	Ala	Ala	Pro	Tyr	His	Val	Ile	Gln	Leu
			260					265					270		
Val	Asn	Leu	Gln	Met	Glu	Gln	Pro	Thr	Leu	Ala	Phe	Tyr	Val	Gly	Tyr
		275					280					285			
Tyr	Leu	Ser	Ile	Cys	Leu	Ser	Tyr	Ala	Ser	Ser	Ser	Ile	Asn	Pro	Phe
	290					295					300				
Leu	Tyr	Ile	Leu	Leu	Ser	Gly	Asn	Phe	Gln	Lys	Arg	Leu	Pro	Gln	Ile
305					310					315					320
Gln	Arg	Arg	Ala	Thr	Glu	Lys	Glu	Ile	Asn	Asn	Met	Gly	Asn	Thr	Leu
				325					330					335	
Lys	Ser	His	Phe												
			340												

<210> 118  
 <211> 342  
 <212> PRT  
 <213> Homo sapiens

<400> 118

Met	Thr	Ser	Asn	Phe	Ser	Gln	Pro	Val	Val	Gln	Leu	Cys	Tyr	Glu	Asp
1				5					10					15	
Val	Asn	Gly	Ser	Cys	Ile	Glu	Thr	Pro	Tyr	Ser	Pro	Gly	Ser	Arg	Val
			20					25					30		
Ile	Leu	Tyr	Thr	Ala	Phe	Ser	Phe	Gly	Ser	Leu	Leu	Ala	Val	Phe	Gly
		35					40					45			
Asn	Leu	Leu	Val	Met	Thr	Ser	Val	Leu	His	Phe	Lys	Gln	Leu	His	Ser
	50					55					60				
Pro	Thr	Asn	Phe	Leu	Ile	Ala	Ser	Leu	Ala	Cys	Ala	Asp	Phe	Leu	Val
65					70					75					80
Gly	Val	Thr	Val	Met	Leu	Phe	Ser	Met	Val	Arg	Thr	Val	Glu	Ser	Cys
				85					90					95	
Trp	Tyr	Phe	Gly	Ala	Lys	Phe	Cys	Thr	Leu	His	Ser	Cys	Cys	Asp	Val
			100					105					110		
Ala	Phe	Cys	Tyr	Ser	Ser	Val	Leu	His	Leu	Cys	Phe	Ile	Cys	Ile	Asp
		115					120					125			

Arg Tyr Ile Val Val Thr Asp Pro Leu Val Tyr Ala Thr Lys Phe Thr  
 130 135 140  
 Val Ser Val Ser Gly Ile Cys Ile Ser Val Ser Trp Ile Leu Pro Leu  
 145 150 155 160  
 Thr Tyr Ser Gly Ala Val Phe Tyr Thr Gly Val Asn Asp Asp Gly Leu  
 165 170 175  
 Glu Glu Leu Val Ser Ala Leu Asn Cys Val Gly Gly Cys Gln Ile Ile  
 180 185 190  
 Val Ser Gln Gly Trp Val Leu Ile Asp Phe Leu Leu Phe Phe Ile Pro  
 195 200 205  
 Thr Leu Val Met Ile Ile Leu Tyr Ser Lys Ile Phe Leu Ile Ala Lys  
 210 215 220  
 Gln Gln Ala Ile Lys Ile Glu Thr Thr Ser Ser Lys Val Glu Ser Ser  
 225 230 235 240  
 Ser Glu Ser Tyr Lys Ile Arg Val Ala Lys Arg Glu Arg Lys Ala Ala  
 245 250 255  
 Lys Thr Leu Gly Val Thr Val Leu Ala Phe Val Ile Ser Trp Leu Pro  
 260 265 270  
 Tyr Thr Val Asp Ile Leu Ile Asp Ala Phe Met Gly Phe Leu Thr Pro  
 275 280 285  
 Ala Tyr Ile Tyr Glu Ile Cys Cys Trp Ser Ala Tyr Tyr Asn Ser Ala  
 290 295 300  
 Met Asn Pro Leu Ile Tyr Ala Leu Phe Tyr Pro Trp Phe Arg Lys Ala  
 305 310 315 320  
 Ile Lys Leu Ile Leu Ser Gly Asp Val Leu Lys Ala Ser Ser Ser Thr  
 325 330 335  
 Ile Ser Leu Phe Leu Glu  
 340

<210> 119  
 <211> 323  
 <212> PRT  
 <213> Homo sapiens

<400> 119

Met Pro Leu Pro Val Pro Pro Ala Gly Ala Gln Lys Thr Pro Glu Asp  
 1 5 10 15  
 His Val Cys Leu His Leu Ala Gly Pro Ser Pro Ala Pro Ser Glu Pro  
 20 25 30  
 Ala Arg Met Phe Gly Leu Phe Gly Leu Trp Arg Thr Phe Asp Ser Val  
 35 40 45

Val Phe Tyr Leu Thr Leu Ile Val Gly Leu Gly Gly Pro Val Gly Asn  
 50 55 60  
 Gly Leu Val Leu Trp Asn Leu Gly Phe Arg Ile Lys Lys Gly Pro Phe  
 65 70 75 80  
 Ser Ile Tyr Leu Leu His Leu Ala Ala Ala Asp Phe Leu Phe Leu Ser  
 85 90 95  
 Cys Arg Val Gly Phe Ser Val Ala Gln Ala Ala Leu Gly Ala Gln Asp  
 100 105 110  
 Thr Leu Tyr Phe Val Leu Thr Phe Leu Trp Phe Ala Val Gly Leu Trp  
 115 120 125  
 Leu Leu Ala Ala Phe Ser Val Glu Arg Cys Leu Ser Asp Leu Phe Pro  
 130 135 140  
 Ala Cys Tyr Gln Gly Cys Arg Pro Arg His Ala Ser Ala Val Leu Cys  
 145 150 155 160  
 Ala Leu Val Trp Thr Pro Thr Leu Pro Ala Val Pro Leu Pro Ala Asn  
 165 170 175  
 Ala Cys Gly Leu Leu Arg Asn Ser Ala Cys Pro Leu Val Cys Pro Arg  
 180 185 190  
 Tyr His Val Ala Ser Val Thr Trp Phe Leu Val Leu Ala Arg Val Ala  
 195 200 205  
 Trp Thr Ala Gly Val Val Leu Phe Val Trp Val Thr Cys Cys Ser Thr  
 210 215 220  
 Arg Pro Arg Pro Arg Leu Tyr Gly Ile Val Leu Gly Ala Leu Leu Leu  
 225 230 235 240  
 Leu Phe Phe Cys Gly Leu Pro Ser Val Phe Tyr Trp Ser Leu Gln Pro  
 245 250 255  
 Leu Leu Asn Phe Leu Leu Pro Val Phe Ser Pro Leu Ala Thr Leu Leu  
 260 265 270  
 Ala Cys Val Asn Ser Ser Ser Lys Pro Leu Ile Tyr Ser Gly Leu Gly  
 275 280 285  
 Arg Gln Pro Gly Lys Arg Glu Pro Leu Arg Ser Val Leu Arg Arg Ala  
 290 295 300  
 Leu Gly Glu Gly Ala Glu Leu Gly Ala Arg Gly Gln Ser Leu Pro Met  
 305 310 315 320  
 Gly Leu Leu

<210> 120  
 <211> 336  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 120

```

Met Asn Asn Asn Thr Thr Cys Ile Gln Pro Ser Met Ile Ser Ser Met
1      5      10      15
Ala Leu Pro Ile Ile Tyr Ile Leu Leu Cys Ile Val Gly Val Phe Gly
20      25      30
Asn Thr Leu Ser Gln Trp Ile Phe Leu Thr Lys Ile Gly Lys Lys Thr
35      40      45
Ser Thr His Ile Tyr Leu Ser His Leu Val Thr Ala Asn Leu Leu Val
50      55      60
Cys Ser Ala Met Pro Phe Met Ser Ile Tyr Phe Leu Lys Gly Phe Gln
65      70      75      80
Trp Glu Tyr Gln Ser Ala Gln Cys Arg Val Val Asn Phe Leu Gly Thr
85      90      95
Leu Ser Met His Ala Ser Met Phe Val Ser Leu Leu Ile Leu Ser Trp
100     105     110
Ile Ala Ile Ser Arg Tyr Ala Thr Leu Met Gln Lys Asp Ser Ser Gln
115     120     125
Glu Thr Thr Ser Cys Tyr Glu Lys Ile Phe Tyr Gly His Leu Leu Lys
130     135     140
Lys Phe Arg Gln Pro Asn Phe Ala Arg Lys Leu Cys Ile Tyr Ile Trp
145     150     155     160
Gly Val Val Leu Gly Ile Ile Ile Pro Val Thr Val Tyr Tyr Ser Val
165     170     175
Ile Glu Ala Thr Glu Gly Glu Glu Ser Leu Cys Tyr Asn Arg Gln Met
180     185     190
Glu Leu Gly Ala Met Ile Ser Gln Ile Ala Gly Leu Ile Gly Thr Thr
195     200     205
Phe Ile Gly Phe Ser Phe Leu Val Val Leu Thr Ser Tyr Tyr Ser Phe
210     215     220
Val Ser His Leu Arg Lys Ile Arg Thr Cys Thr Ser Ile Met Glu Lys
225     230     235     240
Asp Leu Thr Tyr Ser Ser Val Lys Arg His Leu Leu Val Ile Gln Ile
245     250     255
Leu Leu Ile Val Cys Phe Leu Pro Tyr Ser Ile Phe Lys Pro Ile Phe
260     265     270
Tyr Val Leu His Gln Arg Asp Asn Cys Gln Gln Leu Asn Tyr Leu Ile
275     280     285
Glu Thr Lys Asn Ile Leu Thr Cys Leu Ala Ser Ala Arg Ser Ser Thr
290     295     300

```

Asp Pro Ile Ile Phe Leu Leu Leu Asp Lys Thr Phe Lys Lys Thr Leu  
 305 310 315 320

Tyr Asn Leu Phe Thr Lys Ser Asn Ser Ala His Met Gln Ser Tyr Gly  
 325 330 335

<210> 121  
 <211> 28  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Primer

<400> 121  
 ttcaaagctt atgacgtcca cctgcacc

28

<210> 122  
 <211> 35  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Primer

<400> 122  
 ttcactcgag tcaaggaaaa gtagcagaat cgtag

35

<210> 123  
 <211> 35  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Primer

<400> 123  
 gatcgaattc atgatggagc ccagagaagc tggac

35

<210> 124  
 <211> 28  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Primer

<400> 124  
 cgagtcaggc tgctatgtcc accaggcc

28

<210> 125  
 <211> 37  
 <212> DNA  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Primer

&lt;400&gt; 125

gcataagctt ccatgtggag ctgcagctgg ttcaacg

37

&lt;210&gt; 126

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Primer

&lt;400&gt; 126

gcattctcgag cctacgccag cacctgctgc acc

33

&lt;210&gt; 127

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Primer

&lt;400&gt; 127

gcataagctt ccatggggaa cgattctgtc agc

33

&lt;210&gt; 128

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Primer

&lt;400&gt; 128

gcattctcgag cctacacctc catctccgag acc

33

&lt;210&gt; 129

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Primer

&lt;400&gt; 129

gatcaagctt gcatggcacc ttctcatcgg g

31

&lt;210&gt; 130

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial

<220>  
<223> Primer  
  
<400> 130  
gatcctcgag tcaaaggtcc cattccggac c 31  
  
<210> 131  
<211> 13  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Primer  
  
<400> 131  
tccctgtgcc ccc 13  
  
<210> 132  
<211> 23  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Primer  
  
<400> 132  
ttataggaga cccatgggca ggg 23  
  
<210> 133  
<211> 40  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Primer  
  
<400> 133  
ggtgctctgg aacctcggct tccgcatcaa gaagggcccc 40  
  
<210> 134  
<211> 40  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Primer  
  
<400> 134  
ggggcccttc ttgatgcgga agccgaggtt ccagagcacc 40  
  
<210> 135  
<211> 38  
<212> DNA



<213> Artificial

<220>

<223> Primer

<400> 135

ttcaaagctt atgaacaaca atacaacatg tattcaac

38

<210> 136

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 136

tcaaacatat gattgcatat gtg

23

<210> 137

<211> 39

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 137

gcacatatgc aatcatatgg ttgactcgag tgaaaaggg

39

<210> 138

<211> 39

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 138

cccttttcac tcgagtcaac catatgattg catatgtgc

39

<210> 139

<211> 30

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 139

ctccatctgt ctcagctatg ccagcagcag

30

<210> 140

<211> 13

<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 140  
tctcagctat gcc

13

<210> 141  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 141  
tccttctcag tcgctcttc

19

<210> 142  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 142  
cctccacttg tgcttcac

19

<210> 143  
<211> 21  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 143  
aaaatctatc aacacccagc c

21

<210> 144  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 144  
gatcaagctt accatgacca gcaatttttc cc

32

<210> 145

<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 145  
gatcctcgag cttattctaa aaataaacta atgg

34

<210> 146  
<211> 25  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 146  
catgatctgg ggggccagcc ccagc

25

<210> 147  
<211> 25  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 147  
gctggggctg gccccccaga tcatg

25

<210> 148  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 148  
gacatcaatt tcagtgagga tgacgtcgag gcag

34

<210> 149  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 149  
tgcctcgacg tcatcctcac tgaaattgat gtc

33

<210> 150  
<211> 34  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 150  
cgaactccgc actcctggcc agggccctgc gggc

34

<210> 151  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 151  
gcccgaggg ccctggccag gagtgcggag ttc

33

<210> 152  
<211> 48  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 152  
gcgtaatacg actcactata gggagacctg ccagtgtggt agatacag

48

<210> 153  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 153  
ggatgtgatg atggtgcag

19

<210> 154  
<211> 47  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 154  
gcgtaatacg actcactata gggagaccgg atgtgatgat ggtgcag

47

<210> 155  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 155  
tgccagtgtg gtagatacag

20

<210> 156  
<211> 48  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 156  
gcgtaatacag actcactata gggagaccac cagcaatttt tcccaacc

48

<210> 157  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 157  
aataccagca gctctccac

19

<210> 158  
<211> 47  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 158  
gcgtaatacag actcactata gggagaccaa taccagcagc tctccac

47

<210> 159  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 159  
accagcaatt tttcccaacc

20

<210> 160  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 160  
gtgactaact ctgcctgcg

19

<210> 161  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 161  
ttgcgctgca aactagcg

19

<210> 162  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 162  
acagcccaa agccaaacac

20

<210> 163  
<211> 22  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 163  
ccgcaggagc aatgaaaatc ag

22

<210> 164  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 164

tctccaaact cctggccttc

20

<210> 165  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 165  
gcagggcagc tttttcatcc

20

<210> 166  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 166  
acgcccgtg aaccgtatac

20

<210> 167  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 167  
gggtgccacc tggttgctc

19

<210> 168  
<211> 18  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 168  
atggcacctt ctcatcgg

18

<210> 169  
<211> 18  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 169  
acgtagtaca ccgccttg 18

<210> 170  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 170  
agcaggtaga tggagaagg 19

<210> 171  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 171  
gactcctgag gaccatgtc 19

<210> 172  
<211> 21  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 172  
cattggaacc acatttattg g 21

<210> 173  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 173  
aagcaagaca ggtgagaatg 20

<210> 174  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer



<400> 174  
ccatctgtct cagctatgcc 20

<210> 175  
<211> 19  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 175  
tcctttctcag tcgctcttc 19

<210> 176  
<211> 21  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 176  
caaacaacaa acagcagaac c 21

<210> 177  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 177  
tcacagtcac acctaccaag 20

<210> 178  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Primer

<400> 178  
ccatctgtct cagctatgcc 20

<210> 179  
<211> 19  
<212> DNA  
<213> Artificial

<220>

<223> Primer

<400> 179

tccttctcag tcgctcttc

19

<210> 180

<211> 21

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 180

gtctatgcta ccaagttcac c

21

<210> 181

<211> 19

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 181

attcctccag cccatcatc

19

<210> 182

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Primer

<400> 182

Ala Pro Arg Thr Pro Gly Gly Arg Arg

1

5

<210> 183

<211> 19

<212> PRT

<213> Homo sapiens

<400> 183

Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys

1

5

10

15

Trp Gln Val

<210> 184

<211> 17

<212> PRT

<213> Salmon

<400> 184

Asp	Thr	Met	Arg	Cys	Met	Val	Gly	Arg	Val	Tyr	Arg	Pro	Cys	Trp	Glu
1				5					10					15	

Val



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number  
**WO 01/062797 A3**

(51) International Patent Classification<sup>7</sup>: **C07K 14/72**,  
C12N 15/63, 15/11, C07K 1/14, C12Q 1/68, G01N 33/53,  
C07K 16/28, A61K 38/17, 39/395

(21) International Application Number: PCT/US01/05676

(22) International Filing Date: 23 February 2001 (23.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/184,305	23 February 2000 (23.02.2000)	US
60/184,304	23 February 2000 (23.02.2000)	US
60/184,303	23 February 2000 (23.02.2000)	US
60/184,397	23 February 2000 (23.02.2000)	US
60/184,247	23 February 2000 (23.02.2000)	US
60/186,457	2 March 2000 (02.03.2000)	US
60/186,810	3 March 2000 (03.03.2000)	US
60/188,064	9 March 2000 (09.03.2000)	US
60/188,880	13 March 2000 (13.03.2000)	US
60/194,344	3 April 2000 (03.04.2000)	US
60/213,861	23 June 2000 (23.06.2000)	US
60/217,370	11 July 2000 (11.07.2000)	US
60/217,369	11 July 2000 (11.07.2000)	US
60/218,337	14 July 2000 (14.07.2000)	US
60/218,492	20 July 2000 (20.07.2000)	US

(71) Applicant (for all designated States except US): **PHARMACIA & UPJOHN COMPANY** [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VOGELI, Gabriel**

[US/US]; 2576 9th Avenue, Seattle, WA 98119 (US). **WOOD, Linda, S.** [US/US]; 10193 Fox Hollow, Portage, MI 49024 (US). **PARODI, Luis, A.** [SE/SE]; Grevgafan-24, S-115 43 Stockholm (SE). **LIND, Peter** [SE/SE]; Borjegatan 31 C, S-751 29 Uppsala (SE).

(74) Agents: **DELUCA, Mark et al.**; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
24 October 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: G PROTEIN-COUPLED RECEPTORS

(57) Abstract: The present invention provides a gene encoding a G protein-coupled receptor termed nGPCR-x; constructs and recombinant host cells incorporating the genes; the nGPCR-x polypeptides encoded by the gene; antibodies to the nGPCR-x polypeptides; and methods of making and using all of the foregoing.



WO 01/062797 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PO/US 01/05676

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/72 C12N15/63 C12N15/11 C07K1/14 C12Q1/68  
G01N33/53 C07K16/28 A61K38/17 A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N C12Q G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, SEQUENCE SEARCH, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 892 051 A (SMITHKLINE BEECHAM CORP) 20 January 1999 (1999-01-20)  page 12, line 1 - line 55 page 13, line 15 -page 15, line 10 ---	1-48,50, 51, 53-56, 58-62, 64-74, 82-89, 138
E	WO 01 36471 A (ARENA PHARMACEUTICALS INC) 25 May 2001 (2001-05-25)  SEQ ID NO:25; SEQ ID NO:26 --- -/-	1-48,50, 51,53, 54,61,62

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

30 November 2001

Date of mailing of the international search report

15. 5. 02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-9016

Authorized officer

Schönwasser, D

# INTERNATIONAL SEARCH REPORT

International Application No

P US 01/05676

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL [Online]  Accession no. AL161458,  16 March 2000 (2000-03-16)  BURTON J.: "Human DNA sequence ***  SEQUENCE IN PROGRESS *** FROM CLONE  RP11-163L4"  XP002184439  abstract</p>	1-30
A	<p>---  WO 94 12635 A (ICOS CORP)  9 June 1994 (1994-06-09)</p>	1-48, 50, 51, 53-56, 58-62, 64-74, 82-89, 138
A	<p>SEQ ID NO:44</p> <p>---  ALTSCHUL S F ET AL: "Iterated profile  searches with PSI-BLAST-a tool for  discovery in protein databases"  TIBS TRENDS IN BIOCHEMICAL SCIENCES,  ELSEVIER PUBLICATION, CAMBRIDGE, EN,  vol. 23, no. 11,  1 November 1998 (1998-11-01), pages  444-447, XP004143492  ISSN: 0968-0004  the whole document</p>	1-28, 35-39
A	<p>---  STADEL ET AL: "Orphan G protein-coupled  receptors:"  TRENDS IN PHARMACOLOGICAL SCIENCES,  ELSEVIER TRENDS JOURNAL, CAMBRIDGE, GB,  no. 18, 1 November 1997 (1997-11-01),  pages 430-437, XP002073279  ISSN: 0165-6147  the whole document</p> <p style="text-align: center;">-----</p>	1-28, 35-39

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 01/05676**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
**see FURTHER INFORMATION sheet PCT/ISA/210**
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

**see additional sheet**

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**SIEHE PCT/ISA 210**

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Claims Nos.: 49,52,57,75-81 no search; 1,45,47,48,50,51,53-56,58  
incompl. search

Present claims 45, 50 and 53 relate to an extremely large number of methods involving possible nGPCR-x or nucleic acids encoding nGPCR-x. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the nGPCR-x or nucleic acids encoding nGPCR-x of SEQ ID NO:61-120 or SEQ ID NO:1-60, respectively.

Further, claims 49,52 and 57 refer to compounds identified in a certain screening process without giving a true technical characterization of said compounds. Moreover, no such compounds are defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (Art. 5 and 6 PCT). No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the results to be achieved.

Furthermore, claims 75 and 76 refer to an allelic variant of certain claimed G-protein coupled receptors (GPCR) and claims 77-81 refer to polynucleotides encoding polypeptides which are related to mental disorder and which differ in at least one residue from the claimed GPCR polypeptides. No true technical characterization of said compounds is given and no such compounds are defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (Art. 5 and 6 PCT). No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-48,50,51,53-56,58-62,64-74,82-89,  
138 (all partially)

Invention 1:

An isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:61; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:1; vector comprising said SEQ ID NO:1 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; methods of screening a human subject involving said polypeptide, a kit for screening a human subject involving an oligonucleotide of above nucleic acid molecule; methods for identifying an allelic variant of above nucleic acid molecule; methods to identify compounds useful for the treatment of mental disorder or as modulators of binding between above polypeptide and its binding partner; a method of purifying a G-protein by use of above polypeptide; a method of identifying an allelic variant of above nucleic acid molecule.

2. Claims: 1-48,50,51,53-56,58-62,64-74,82-89,  
138 (all partially)

Inventions 2-15:

The inventions 2-15 relate to subject-matter as defined above for "invention 1", whereby each invention refers to one of the polypeptide sequences of SEQ ID NO:62,86,91,94, 96,97,111-116,119 and 120 (and the respective nucleic acid sequences of SEQ ID NO:2,8,31,43,36,37,51-56, 59,60).

3. Claims: 1-5,7-10,12-27,29-38,40-45,47,48,50,51,53,55,56,  
58-60,86-88,138 (all partially)

Invention 16:

An isolated nucleic acid molecule encoding a polypeptide

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

comprising the amino acid sequence of SEQ ID NO:63; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:3; vector comprising said SEQ ID NO:3 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; a method of purifying a G-protein by use of above polypeptide; a method of identifying an allelic variant of above nucleic acid molecule.

4. Claims: 1-5,7-10,12-27,29-38,40-45,47,48,50,51,53,55,56, 58-60,86-88,138 (all partially)

Inventions 17-56:

The inventions 17-56 relate to subject-matter as defined above for "invention 15", whereby each invention refers to one of the polypeptide sequences of SEQ ID NO:64-67,69-90, 92,93,95,98,111-120 (and the respective nucleic acid sequences of SEQ ID NO:4-7,9-30,32,33,35,38,41-50).

5. Claims: 1-48,50,51,53-56,58-74,82-89,95-113,115-117, 138 (all partially)

Invention 57:

An isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:99; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:39; vector comprising said SEQ ID NO:39 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; methods of screening a human subject involving said polypeptide, a kit for screening a human subject involving an oligonucleotide of above nucleic acid molecule; methods for identifying an allelic variant of above nucleic acid

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

molecule; methods to identify compounds useful for the treatment of mental disorder or as modulators of binding between above polypeptide and its binding partner; a method of purifying a G-protein by use of above polypeptide; methods for identifying a compound that binds to or modulates the activity of above polypeptide; a method of screening a human subject to diagnose a disorder affecting the brain, a kit for screening a human subject to diagnose a mental disorder; methods for identifying a compound useful for the treatment of a mental disorder or as a modulator of binding between above polypeptide and its binding partner; a method of identifying an allelic variant of above nucleic acid molecule.

6. Claims: 1-48,50,51,53-56,58-74,82-89,123-137,  
139-141 (all partially)

## Invention 58:

An isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:100; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:40; vector comprising said SEQ ID NO:40 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; methods of screening a human subject involving said polypeptide, a kit for screening a human subject involving an oligonucleotide of above nucleic acid molecule; methods to identify compounds useful for the treatment of mental disorder or as modulators of binding between above polypeptide and its binding partner; a method of purifying a G-protein by use of above polypeptide; methods of identifying a compound that binds to above polypeptide or modulated its activity; a method of screening a human subject to diagnose a disorder affecting the brain involving above nucleic acid molecule, a kit for screening a human subject to diagnose a mental disorder involving above nucleic acid molecule; a method of identifying an allelic variant of above nucleic acid molecule, methods of identifying compounds that modulate the biological activity of above polypeptide or which are useful for the treatment of a mental disorder or as modulators of binding between above polypeptide and its binding partner.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## 7. Claims: 1-48,50,51,53-56,58-74,82-117,138 (all partially)

## Invention 59:

An isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:117; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:57; vector comprising said SEQ ID NO:57 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; methods of screening a human subject involving said polypeptide, a kit for screening a human subject involving an oligonucleotide of above nucleic acid molecule; methods for identifying an allelic variant of above nucleic acid molecule; methods to identify compounds useful for the treatment of mental disorder or as modulators of binding between above polypeptide and a binding partner; a method of purifying above polypeptide; a nucleic acid encoding a polypeptide comprising SEQ ID NO:117; a nucleic acid molecule comprising SEQ ID NO:57; a polypeptide comprising SEQ ID NO:117; methods for identifying a compound that binds to or modulates the activity of above polypeptide; a method of screening a human subject to diagnose a disorder affecting the brain, a kit for screening a human subject to diagnose a mental disorder; methods for identifying a compound useful for the treatment of a mental disorder or as a modulator of binding between above polypeptide and a binding partner; methods of identifying an allelic variant of above nucleic acid molecule or above amino acid.

8. Claims: 1-48,50,51,53-56,58-74,82-89,  
118-141 (all partially)

## Invention 60:

An isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:118; an isolated nucleic acid sequence comprising the sequence SEQ ID NO:58; vector comprising said SEQ ID NO:58 and host cell comprising said vector; compositions comprising said nucleic acid molecule or said vector, methods of producing above polypeptide; said isolated polypeptide; an isolated antibody which binds to an epitope of above polypeptide; compositions comprising said polypeptide or antibody; a method for

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

inducing an immune response in a mammal by using said polypeptide; methods for identifying compounds that bind above polypeptide or above nucleic acid molecule or which modulate the activity of above polypeptide, a method of identifying an animal homologue of said polypeptide; methods of screening a human subject involving said polypeptide, a kit for screening a human subject involving an oligonucleotide of above nucleic acid molecule; methods for identifying an allelic variant of above nucleic acid molecule; methods to identify compounds useful for the treatment of mental disorder or as modulators of binding between above polypeptide and a binding partner; a method of purifying above polypeptide; a nucleic acid molecule comprising a sequence that encodes SEQ ID NO:118; a polypeptide comprising an amino acid sequence homologous to SEQ ID NO:118; methods of identifying a compound that binds to above polypeptide or modulated its activity; a method of screening a human subject to diagnose a disorder affecting the brain involving above nucleic acid molecule, a kit for screening a human subject to diagnose a mental disorder involving above nucleic acid molecule; a method of identifying an allelic variant of above nucleic acid molecule, methods of identifying compounds that modulate the biological activity of above polypeptide or which are useful for the treatment of a mental disorder or as modulators of binding between above polypeptide and its binding partner.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

US 01/05676

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0892051	A	20-01-1999	US 5955308 A	21-09-1999
			EP 0892051 A2	20-01-1999
			JP 11098988 A	13-04-1999
-----				
WO 0136471	A	25-05-2001	AU 1769601 A	30-05-2001
			WO 0136471 A2	25-05-2001
-----				
WO 9412635	A	09-06-1994	AT 164884 T	15-04-1998
			CA 2128208 A1	09-06-1994
			DE 69317883 D1	14-05-1998
			DE 69317883 T2	12-11-1998
			DK 630405 T3	04-01-1999
			EP 0630405 A1	28-12-1994
			EP 0846762 A2	10-06-1998
			ES 2118372 T3	16-09-1998
			JP 7503145 T	06-04-1995
			JP 2000350582 A	19-12-2000
			US 6348574 B1	19-02-2002
			WO 9412635 A2	09-06-1994
			US 5759804 A	02-06-1998
			US 6107475 A	22-08-2000
-----				

